

Akihiko Kamoshita

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6263839/publications.pdf>

Version: 2024-02-01

38
papers

1,345
citations

430874

18
h-index

345221

36
g-index

38
all docs

38
docs citations

38
times ranked

1130
citing authors

#	ARTICLE	IF	CITATIONS
1	Genetic analysis of root vascular traits in a population from two temperate japonica rice ecotypes. <i>Plant Production Science</i> , 2022, 25, 320-336.	2.0	1
2	Recent changes in rice production in rainfed lowland and irrigated ecosystems in Thailand. <i>Plant Production Science</i> , 2021, 24, 15-28.	2.0	10
3	Ecophysiology of drill-seeded rice under reduced nitrogen fertilizer and reduced irrigation during El Niño in Central Colombia. <i>Plant Production Science</i> , 2021, 24, 418-432.	2.0	1
4	Eco-physiological evaluation of Stele Transversal Area 1 for rice root anatomy and shoot growth. <i>Plant Production Science</i> , 2020, 23, 202-210.	2.0	6
5	Salinity intrusion reduces grain yield in coastal paddy fields: case study in two estuaries in the Red River Delta, Vietnam. <i>Paddy and Water Environment</i> , 2020, 18, 399-416.	1.8	3
6	Preliminary Assessment of Rice Production in Coastal Part of Red River Delta Surrounding Xuan Thuy National Park, Vietnam, for Improving Resilience. <i>Science for Sustainable Societies</i> , 2018, , 7-38.	0.5	2
7	Comparative analysis of farmers engaged in participatory research to cope with climate change versus non-participants in Northeast Thailand. <i>Plant Production Science</i> , 2018, 21, 287-301.	2.0	5
8	Salinity intrusion and rice production in Red River Delta under changing climate conditions. <i>Paddy and Water Environment</i> , 2017, 15, 37-48.	1.8	23
9	Near-isogenic lines of IR64 (<i>Oryza sativa</i> subsp. indica cv.) introgressed with DEEPER ROOTING 1 and STELE TRANSVERSAL AREA 1 improve rice yield formation over the background parent across three water management regimes. <i>Plant Production Science</i> , 2017, 20, 249-261.	2.0	6
10	Association between root growth angle and root length density of a near-isogenic line of IR64 rice with DEEPER ROOTING 1 under different levels of soil compaction. <i>Plant Production Science</i> , 2017, 20, 162-175.	2.0	26
11	Residual effects of cultivation methods on weed seed banks and weeds in Cambodia. <i>Weed Biology and Management</i> , 2016, 16, 93-107.	1.4	6
12	Field level damage of deepwater rice by the 2011 Southeast Asian Flood in a flood plain of Tonle Sap Lake, Northwest Cambodia. <i>Paddy and Water Environment</i> , 2015, 13, 455-463.	1.8	6
13	Weed biodiversity and rice production during the irrigation rehabilitation process in Cambodia. <i>Agriculture, Ecosystems and Environment</i> , 2014, 194, 1-6.	5.3	17
14	Water availability, management practices and grain yield for deepwater rice in Northwest Cambodia. <i>Field Crops Research</i> , 2013, 152, 44-56.	5.1	4
15	Improvement of resilience of crop production under waterlogged conditions in humid Asia and Africa for sustainable agricultural development. <i>Field Crops Research</i> , 2013, 152, 1-2.	5.1	1
16	Current Status of Research on Improvement of Drought Resistance in Rice (<i>Oryza sativa</i> L.). <i>Japanese Journal of Crop Science</i> , 2011, 80, 1-12.	0.2	6
17	Farmers' Management Practices and Grain Yield of Rice in Response to Different Water Environments in Kamping Puoy Irrigation Rehabilitation Area in Northwest Cambodia. <i>Plant Production Science</i> , 2011, 14, 377-390.	2.0	9
18	Ecophysiological study on weed seed banks and weeds in Cambodian paddy fields with contrasting water availability. <i>Weed Biology and Management</i> , 2010, 10, 261-272.	1.4	33

#	ARTICLE	IF	CITATIONS
19	High-yielding Crop Management by Enhancing Growth in Reproductive Stage of Direct-Seeded Rainfed Lowland Rice (<i>Oryza sativa</i> L.) in Northeast Thailand. <i>Plant Production Science</i> , 2010, 13, 104-115.	2.0	5
20	Spatial variability in the growth of direct-seeded rainfed lowland rice (<i>Oryza sativa</i> L.) in northeast Thailand. <i>Field Crops Research</i> , 2009, 111, 251-261.	5.1	13
21	Assessment of management of direct seeded rice production under different water conditions in Cambodia. <i>Paddy and Water Environment</i> , 2008, 6, 91-103.	1.8	27
22	Phenotypic and genotypic analysis of drought-resistance traits for development of rice cultivars adapted to rainfed environments. <i>Field Crops Research</i> , 2008, 109, 1-23.	5.1	265
23	Preflowering Abortion Reduces Spikelet Number in Upland Rice (<i>Oryza sativa</i> L.) under Water Stress. <i>Crop Science</i> , 2008, 48, 2389-2395.	1.8	87
24	Varietal Differences in Stem Diameter and Rooting Number of Phytomers in Conjunction with Root System Development of Field-Grown Rice (<i>Oryza sativa</i> L.). <i>Plant Production Science</i> , 2007, 10, 357-360.	2.0	6
25	Evaluation of Water-Saving Rice-Winter Crop Rotation System in a Suburb of Tokyo. <i>Plant Production Science</i> , 2007, 10, 219-231.	2.0	10
26	Historical Changes in Urban Rice Production Systems in Tokyo, Japan. <i>Plant Production Science</i> , 2007, 10, 245-269.	2.0	11
27	Growth of Rice (<i>Oryza Sativa</i> L.) Cultivars Under Upland Conditions With Different Levels of Water Supply3. Root System Development, Soil Moisture Changeand Plant Water Status. <i>Plant Production Science</i> , 2007, 10, 3-13.	2.0	46
28	Genotypic differences in grain yield of transplanted and direct-seeded rainfed lowland rice (<i>Oryza</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50	5.1	39
29	Evaluating the resistance of six rice cultivars to drought: restriction of deep rooting and the use of raised beds. <i>Plant and Soil</i> , 2007, 300, 149-161.	3.7	38
30	Improvement of rice (<i>Oryza sativa</i> L.) growth in upland conditions with deep tillage and mulch. <i>Soil and Tillage Research</i> , 2007, 92, 30-44.	5.6	22
31	Genotypic Variation in Root Growth Angle in Rice (<i>Oryza sativa</i> L.) and its Association with Deep Root Development in Upland Fields with Different Water Regimes. <i>Plant and Soil</i> , 2006, 287, 117-129.	3.7	202
32	Effect of Planting Density on Grain Yield and Water Productivity of Rice (<i>Oryza sativa</i> L.) Grown in Flooded and Non-flooded Fields in Japan. <i>Plant Production Science</i> , 2006, 9, 298-311.	2.0	37
33	Growth of Three Rice (<i>Oryza sativa</i> L.) Cultivars under Upland Conditions with Different Levels of Water Supply. <i>Plant Production Science</i> , 2006, 9, 422-434.	2.0	38
34	Growth of Three Rice Cultivars (<i>Oryza sativa</i> L.) under Upland Conditions with Different Levels of Water Supply. <i>Plant Production Science</i> , 2006, 9, 435-445.	2.0	35
35	Genotypic Variation in Response of Rainfed Lowland Rice to Prolonged Drought and Rewatering. <i>Plant Production Science</i> , 2004, 7, 406-420.	2.0	85
36	Genotypic Variation in Response of Rainfed Lowland Rice to Drought and Rewatering. <i>Plant Production Science</i> , 2000, 3, 173-179.	2.0	46

#	ARTICLE	IF	CITATIONS
37	Genotypic Variation in Response of Rainfed Lowland Rice to Drought and Rewatering. III. Water extraction during the drought period. <i>Plant Production Science</i> , 2000, 3, 189-196.	2.0	72
38	Genotypic Variation in Response of Rainfed Lowland Rice to Drought and Rewatering. <i>Plant Production Science</i> , 2000, 3, 180-188.	2.0	96