

Shi Zhao

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6268592/publications.pdf>

Version: 2024-02-01

16
papers

244
citations

1040056

9
h-index

996975

15
g-index

17
all docs

17
docs citations

17
times ranked

466
citing authors

#	ARTICLE	IF	CITATIONS
1	Exenatide Twice Daily Plus Glargine Versus Aspart 70/30 Twice Daily in Patients With Type 2 Diabetes With Inadequate Glycemic Control on Premixed Human Insulin and Metformin. <i>Endocrine Practice</i> , 2021, 27, 790-797.	2.1	4
2	STAT3 ameliorates cognitive deficits via regulation of NMDAR expression in an Alzheimer's disease animal model. <i>Theranostics</i> , 2021, 11, 5511-5524.	10.0	25
3	Autologous Bone Marrow Mononuclear Cell Transplantation Therapy Improved Symptoms in Patients with Refractory Diabetic Sensorimotor Polyneuropathy via the Mechanisms of Paracrine and Immunomodulation: A Controlled Study. <i>Cell Transplantation</i> , 2020, 29, 096368972094925.	2.5	5
4	Direct medical costs for patients with type 2 diabetes in 16 tertiary hospitals in urban China: A multicenter prospective cohort study. <i>Journal of Diabetes Investigation</i> , 2019, 10, 539-551.	2.4	30
5	The Association of Hypoglycemia Assessed by Continuous Glucose Monitoring With Cardiovascular Outcomes and Mortality in Patients With Type 2 Diabetes. <i>Frontiers in Endocrinology</i> , 2019, 10, 536.	3.5	15
6	Oxidized LDL upregulates macrophage DPP4 expression via TLR4/TRIF/CD36 pathways. <i>EBioMedicine</i> , 2019, 41, 50-61.	6.1	26
7	Efficacy of autologous bone marrow mononuclear cell transplantation therapy in patients with refractory diabetic peripheral neuropathy. <i>Chinese Medical Journal</i> , 2019, 132, 11-16.	2.3	10
8	Purified Repaired Expanded Hexanucleotide GGGGCC Repeat Noncoding RNA-Caused Neuronal Toxicity in Neuro-2a Cells. <i>Neurotoxicity Research</i> , 2018, 33, 693-701.	2.7	4
9	Reduced mRNA and Protein Expression Levels of Tet Methylcytosine Dioxygenase 3 in Endothelial Progenitor Cells of Patients of Type 2 Diabetes With Peripheral Artery Disease. <i>Frontiers in Immunology</i> , 2018, 9, 2859.	4.8	6
10	The hepatic ectonucleotide pyrophosphatase/phosphodiesterase 1 gene mRNA abundance is reduced by insulin and induced by dexamethasone. <i>Brazilian Journal of Medical and Biological Research</i> , 2018, 51, e6980.	1.5	2
11	Admission Glucose and In-hospital Mortality after Acute Myocardial Infarction in Patients with or without Diabetes. <i>Chinese Medical Journal</i> , 2017, 130, 767-775.	2.3	18
12	Biomarkers for Early Diagnostic of Mild Cognitive Impairment in Type-2 Diabetes Patients: A Multicentre, Retrospective, Nested Case-Control Study. <i>EBioMedicine</i> , 2016, 5, 105-113.	6.1	35
13	Increased BMR in overweight and obese patients with type 2 diabetes may result from an increased fat-free mass. <i>Journal of Huazhong University of Science and Technology [Medical Sciences]</i> , 2016, 36, 59-63.	1.0	8
14	Association of height with peripheral arterial disease in type 2 diabetes. <i>Journal of Endocrinological Investigation</i> , 2015, 38, 57-63.	3.3	2
15	Neurotoxic effects of iron overload under high glucose concentration. <i>Neural Regeneration Research</i> , 2013, 8, 3423-33.	3.0	14
16	Roles of vitamin A status and retinoids in glucose and fatty acid metabolism. <i>Biochemistry and Cell Biology</i> , 2012, 90, 142-152.	2.0	40