

Soledad Verdejo-Lucas

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6270352/publications.pdf>

Version: 2024-02-01

11
papers

251
citations

1163117

8
h-index

1281871

11
g-index

11
all docs

11
docs citations

11
times ranked

177
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Perception of the impact of root-knot nematode-induced diseases in horticultural protected crops of south-eastern Spain. <i>Nematology</i> , 2012, 14, 517-527. | 0.6 | 62 |
| 2 | Selection of virulent populations of <i>Meloidogyne javanica</i> by repeated cultivation of <i>Mi</i> resistance gene tomato rootstocks under field conditions. <i>Plant Pathology</i> , 2009, 58, 990-998. | 2.4 | 59 |
| 3 | Thermal time requirements of root-knot nematodes on zucchini-squash and population dynamics with associated yield losses on spring and autumn cropping cycles. <i>European Journal of Plant Pathology</i> , 2014, 140, 481-490. | 1.7 | 26 |
| 4 | Penetration and reproduction of root-knot nematodes on cucurbit species. <i>European Journal of Plant Pathology</i> , 2014, 138, 863-871. | 1.7 | 23 |
| 5 | Root-knot nematodes on zucchini (<i>Cucurbita pepo</i> subsp. <i>pepo</i>): Pathogenicity and management. <i>Crop Protection</i> , 2019, 126, 104943. | 2.1 | 20 |
| 6 | Population dynamics of <i>Meloidogyne javanica</i> and its relationship with the leaf chlorophyll content in zucchini. <i>Crop Protection</i> , 2015, 70, 8-14. | 2.1 | 17 |
| 7 | Management of Soil-Borne Fungi and Root-Knot Nematodes in Cucurbits through Breeding for Resistance and Grafting. <i>Agronomy</i> , 2020, 10, 1641. | 3.0 | 17 |
| 8 | Pathogenic potential, parasitic success and host efficiency of <i>Meloidogyne incognita</i> and <i>M. javanica</i> on cucurbitaceous plant genotypes. <i>European Journal of Plant Pathology</i> , 2019, 153, 1287-1297. | 1.7 | 9 |
| 9 | Differential feeding site development and reproductive fitness of <i>Meloidogyne incognita</i> and <i>M. javanica</i> on zucchini, a source of resistance to <i>M. incognita</i> . <i>Nematology</i> , 2018, 20, 187-199. | 0.6 | 7 |
| 10 | Suitability of Zucchini and Cucumber Genotypes to Populations of <i>Meloidogyne arenaria</i> , <i>M. incognita</i> , and <i>M. javanica</i> . <i>Journal of Nematology</i> , 2015, 47, 79-85. | 0.9 | 7 |
| 11 | Pathogenicity of <i>Meloidogyne incognita</i> and <i>M. javanica</i> on recombinant inbred lines from a crossing of <i>Cucurbita pepo</i> subsp. <i>pepo</i> × <i>C. pepo</i> subsp. <i>ovifera</i> . <i>Plant Pathology</i> , 2019, 68, 1225-1232. | 2.4 | 4 |