
Jeffrey Weiner

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6271412/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Adolescent social isolation increases anxiety-like behavior and ethanol intake and impairs fear extinction in adulthood: Possible role of disrupted noradrenergic signaling. Neuropharmacology, 2015, 97, 149-159.	4.1	121
2	Increased Basolateral Amygdala Pyramidal Cell Excitability May Contribute to the Anxiogenic Phenotype Induced by Chronic Early-Life Stress. Journal of Neuroscience, 2015, 35, 9730-9740.	3.6	109
3	Adolescent Social Isolation as a Model of Heightened Vulnerability to Comorbid Alcoholism and Anxiety Disorders. Alcoholism: Clinical and Experimental Research, 2016, 40, 1202-1214.	2.4	85
4	Social isolation rearing increases dopamine uptake and psychostimulant potency in the striatum. Neuropharmacology, 2016, 101, 471-479.	4.1	83
5	Adolescent Social Isolation Does Not Lead to Persistent Increases in Anxiety―Like Behavior or Ethanol Intake in Female Longâ€Evans Rats. Alcoholism: Clinical and Experimental Research, 2014, 38, 2199-2207.	2.4	46
6	Chronic treatment with prazosin or duloxetine lessens concurrent anxietyâ€like behavior and alcohol intake: evidence of disrupted noradrenergic signaling in anxietyâ€related alcohol use. Brain and Behavior, 2014, 4, 468-483.	2.2	34
7	Chronic Intermittent Ethanol Exposure Selectively Increases Synaptic Excitability in the Ventral Domain of the Rat Hippocampus. Neuroscience, 2019, 398, 144-157.	2.3	33
8	Effect of β3 adrenoceptor activation in the basolateral amygdala on ethanol seeking behaviors. Psychopharmacology, 2014, 231, 293-303.	3.1	22
9	Fear conditioning selectively disrupts noradrenergic facilitation of GABAergic inhibition in the basolateral amygdala. Neuropharmacology, 2017, 113, 231-240.	4.1	20
10	Behavioral and neurophysiological evidence that lateral paracapsular GABAergic synapses in the basolateral amygdala contribute to the acquisition and extinction of fear learning. Neurobiology of Learning and Memory, 2016, 127, 10-16.	1.9	17
11	Chronic Methylphenidate Treatment During Early Life Is Associated with Greater Ethanol Intake in Socially Isolated Rats. Alcoholism: Clinical and Experimental Research, 2014, 38, 2260-2268.	2.4	13
12	The neural, behavioral, and epidemiological underpinnings of comorbid alcohol use disorder and post-traumatic stress disorder. International Review of Neurobiology, 2021, 157, 69-142.	2.0	11
13	Early life stress induces hyperactivity but not increased anxiety-like behavior or ethanol drinking in outbred heterogeneous stock rats. Alcohol, 2021, 91, 41-51.	1.7	1
14	Early Life Stress Induces Anxietyâ€like Hyperactivity but Not Altered Ethanol Drinking in Outbred Heterogeneous Stock Rats. FASEB Journal, 2020, 34, 1-1.	0.5	0