
## **Richard A Croker**

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6278701/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Severity of Severe Acute Respiratory System Coronavirus 2 (SARS-CoV-2) Alpha Variant (B.1.1.7) in England. Clinical Infectious Diseases, 2022, 75, e1120-e1127.	2.9	71
2	OpenSAFELY NHS Service Restoration Observatory 1: primary care clinical activity in England during the first wave of COVID-19. British Journal of General Practice, 2022, 72, e63-e74.	0.7	22
3	Trends and clinical characteristics of COVID-19 vaccine recipients: a federated analysis of 57.9 million patients' primary care records <i>in situ</i> using OpenSAFELY. British Journal of General Practice, 2022, 72, e51-e62.	0.7	75
4	Mortality among Care Home Residents in England during the first and second waves of the COVID-19 pandemic: an observational study of 4.3 million adults over the age of 65. Lancet Regional Health - Europe, The, 2022, 14, 100295.	3.0	38
5	Potentially inappropriate prescribing of DOACs to people with mechanical heart valves: A federated analysis of 57.9 million patients' primary care records in situ using OpenSAFELY. Thrombosis Research, 2022, 211, 150-153.	0.8	6
6	Association between oral anticoagulants and COVID-19-related outcomes: a population-based cohort study. British Journal of General Practice, 2022, 72, e456-e463.	0.7	3
7	Comparison of methods for predicting COVID-19-related death in the general population using the OpenSAFELY platform. Diagnostic and Prognostic Research, 2022, 6, 6.	0.8	2
8	Risk of severe COVID-19 outcomes associated with immune-mediated inflammatory diseases and immune-modifying therapies: a nationwide cohort study in the OpenSAFELY platform. Lancet Rheumatology, The, 2022, 4, e490-e506.	2.2	61
9	Long COVID burden and risk factors in 10 UK longitudinal studies and electronic health records. Nature Communications, 2022, 13, .	5.8	243
10	Effect of pre-exposure use of hydroxychloroquine on COVID-19 mortality: a population-based cohort study in patients with rheumatoid arthritis or systemic lupus erythematosus using the OpenSAFELY platform. Lancet Rheumatology, The, 2021, 3, e19-e27.	2.2	49
11	Use of non-steroidal anti-inflammatory drugs and risk of death from COVID-19: an OpenSAFELY cohort analysis based on two cohorts. Annals of the Rheumatic Diseases, 2021, 80, 943-951.	0.5	66
12	Association between living with children and outcomes from covid-19: OpenSAFELY cohort study of 12 million adults in England. BMJ, The, 2021, 372, n628.	3.0	56
13	Evaluating the impact of a very low-cost intervention to increase practices' engagement with data and change prescribing behaviour: a randomized trial in English primary care. Family Practice, 2021, 38, 373-380.	0.8	7
14	Hydroxychloroquine treatment does not reduce COVID-19 mortality; underdosing to the wrong patients? – Authors' reply. Lancet Rheumatology, The, 2021, 3, e172-e173.	2.2	1
15	Case fatality risk of the SARS-CoV-2 variant of concern B.1.1.7 in England, 16 November to 5 February. Eurosurveillance, 2021, 26, .	3.9	156
16	Trends in antidepressant prescribing in England. Lancet Psychiatry,the, 2021, 8, 278-279.	3.7	14
17	Identifying Care Home Residents in Electronic Health Records - An OpenSAFELY Short Data Report. Wellcome Open Research, 2021, 6, 90.	0.9	18
18	Ethnic differences in SARS-CoV-2 infection and COVID-19-related hospitalisation, intensive care unit admission, and death in 17 million adults in England: an observational cohort study using the OpenSAFELY platform. Lancet, The, 2021, 397, 1711-1724.	6.3	332

RICHARD A CROKER

#	Article	IF	CITATIONS
19	Clinical coding of long COVID in English primary care: a federated analysis of 58 million patient records <i>in situ</i> using OpenSAFELY. British Journal of General Practice, 2021, 71, e806-e814.	0.7	74
20	Risks of covid-19 hospital admission and death for people with learning disability: population based cohort study using the OpenSAFELY platform. BMJ, The, 2021, 374, n1592.	3.0	70
21	Association between warfarin and COVID-19-related outcomes compared with direct oral anticoagulants: population-based cohort study. Journal of Hematology and Oncology, 2021, 14, 172.	6.9	8
22	OpenSAFELY: impact of national guidance on switching anticoagulant therapy during COVID-19 pandemic. Open Heart, 2021, 8, e001784.	0.9	17
23	Factors associated with COVID-19-related death using OpenSAFELY. Nature, 2020, 584, 430-436.	13.7	4,674
24	Prescription of suboptimal statin treatment regimens: a retrospective cohort study of trends and variation in English primary care. British Journal of General Practice, 2020, 70, e525-e533.	0.7	12
25	Risk of COVID-19-related death among patients with chronic obstructive pulmonary disease or asthma prescribed inhaled corticosteroids: an observational cohort study using the OpenSAFELY platform. Lancet Respiratory Medicine,the, 2020, 8, 1106-1120.	5.2	211
26	Suboptimal prescribing behaviour associated with clinical software design features: a retrospective cohort study in English NHS primary care. British Journal of General Practice, 2020, 70, e636-e643.	0.7	10
27	Impact of Electronic Health Record Interface Design on Unsafe Prescribing of Ciclosporin, Tacrolimus, and Diltiazem: Cohort Study in English National Health Service Primary Care. Journal of Medical Internet Research, 2020, 22, e17003.	2.1	13
28	Trends and variation in unsafe prescribing of methotrexate: a cohort study in English NHS primary care. British Journal of General Practice, 2020, 70, e481-e488.	0.7	8
29	Optimising laboratory monitoring of chronic conditions in primary care: a quality improvement framework. BMJ Open Quality, 2019, 8, e000349.	0.4	14
30	Do doctors in dispensing practices with a financial conflict of interest prescribe more expensive drugs? A cross-sectional analysis of English primary care prescribing data. BMJ Open, 2019, 9, e026886.	0.8	14
31	Six months on: NHS England needs to focus on dissemination, implementation and audit of its low-priority initiative. Journal of the Royal Society of Medicine, 2019, 112, 4-5.	1.1	13
32	Why did some practices not implement new antibiotic prescribing guidelines on urinary tract infection? A cohort study and survey in NHS England primary care. Journal of Antimicrobial Chemotherapy, 2019, 74, 1125-1132.	1.3	28
33	Opioid prescribing trends and geographical variation in England, 1998–2018: a retrospective database study. Lancet Psychiatry,the, 2019, 6, 140-150.	3.7	151
34	Measuring the Impact of an Open Web-Based Prescribing Data Analysis Service on Clinical Practice: Cohort Study on NHS England Data. Journal of Medical Internet Research, 2019, 21, e10929.	2.1	14
35	Is use of homeopathy associated with poor prescribing in English primary care? A cross-sectional study. Journal of the Royal Society of Medicine, 2018, 111, 167-174.	1.1	7
36	Trends, geographical variation and factors associated with prescribing of gluten-free foods in English primary care: a cross-sectional study. BMJ Open, 2018, 8, e021312.	0.8	17

**RICHARD A CROKER** 

#	Article	IF	CITATIONS
37	New mechanism to identify cost savings in English NHS prescribing: minimising †price per unit', a cross-sectional study. BMJ Open, 2018, 8, e019643.	0.8	14
38	The clinician impact and financial cost to the NHS of litigation over pregabalin: a cohort study in English primary care. BMJ Open, 2018, 8, e022416.	0.8	5
39	Detecting change in comparison to peers in NHS prescribing data: a novel application of cumulative sum methodology. BMC Medical Informatics and Decision Making, 2018, 18, 62.	1.5	7
40	Pregabalin: what the patent litigation means for doctors and drug companies. BMJ: British Medical Journal, 2018, 361, k2318.	2.4	1
41	Trends and variation in prescribing of low-priority treatments identified by NHS England: a cross-sectional study and interactive data tool in English primary care. Journal of the Royal Society of Medicine, 2018, 111, 203-213.	1.1	23
42	Prescription data for open toe sandals syndrome. BMJ: British Medical Journal, 2017, 356, j194.	2.4	0
43	Study protocol: Comparison of different risk prediction modelling approaches for COVID-19 related death using the OpenSAFELY platform. Wellcome Open Research, 0, 5, 243.	0.9	3
44	A comprehensive high cost drugs dataset from the NHS in England - An OpenSAFELY-TPP Short Data Report. Wellcome Open Research, 0, 6, 360.	0.9	8
45	Rates of serious clinical outcomes in survivors of hospitalisation with COVID-19 in England: a descriptive cohort study within the OpenSAFELY platform. Wellcome Open Research, 0, 7, 142.	0.9	6