

Jennifer E Stevens-Lapsley

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6280380/jennifer-e-stevens-lapsley-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

120
papers

2,866
citations

30
h-index

51
g-index

134
ext. papers

3,547
ext. citations

2.9
avg, IF

5.45
L-index

#	Paper	IF	Citations
120	Prospective telehealth analysis of functional performance, frailty, quality of life, and mental health after COVID-19 hospitalization.. <i>BMC Geriatrics</i> , 2022 , 22, 251	4.1	0
119	Feasibility and initial efficacy of a multimodal swelling intervention after total knee arthroplasty: A prospective pilot study with historical controls.. <i>Knee</i> , 2022 , 35, 25-33	2.6	
118	Gait mechanics are influenced by quadriceps strength, age, and sex after total knee arthroplasty. <i>Journal of Orthopaedic Research</i> , 2021 , 39, 1523-1532	3.8	1
117	Reliability, responsiveness, and validity of handheld dynamometry for assessing quadriceps strength in total knee arthroplasty. <i>Disability and Rehabilitation</i> , 2021 , 43, 3070-3077	2.4	1
116	Development of reference charts for monitoring quadriceps strength with handheld dynamometry after total knee arthroplasty. <i>Disability and Rehabilitation</i> , 2021 , 1-8	2.4	0
115	Modifiable factors associated with health-related quality of life among lung cancer survivors following curative intent therapy.. <i>Lung Cancer</i> , 2021 , 163, 42-50	5.9	0
114	Determination of Pain Phenotypes in Knee Osteoarthritis Using Latent Profile Analysis. <i>Pain Medicine</i> , 2021 , 22, 653-662	2.8	1
113	Examination of exclusion criteria in total knee arthroplasty rehabilitation trials: influence on the application of evidence in day-to-day practice. <i>Journal of Evaluation in Clinical Practice</i> , 2021 , 27, 1335-1342	2.5	0
112	Older adults in skilled nursing facilities demonstrate low physical activity during and after discharge. <i>Disability and Rehabilitation</i> , 2021 , 1-6	2.4	
111	A multi-center trial of exercise and testosterone therapy in women after hip fracture: Design, methods and impact of the COVID-19 pandemic. <i>Contemporary Clinical Trials</i> , 2021 , 104, 106356	2.3	2
110	Reliability and precision of single frequency bioelectrical impedance assessment of lower extremity swelling following total knee arthroplasty. <i>Physiotherapy Theory and Practice</i> , 2021 , 37, 197-203	1.5	1
109	Systems Approach Is Needed for In-Hospital Mobility: A Qualitative Metasynthesis of Patient and Clinician Perspectives. <i>Archives of Physical Medicine and Rehabilitation</i> , 2021 , 102, 984-998	2.8	5
108	Mixed-Methods Approach to Understanding Determinants of Practice Change in Skilled Nursing Facility Rehabilitation: Adapting to and Sustaining Value With Postacute Reform. <i>Journal of Geriatric Physical Therapy</i> , 2021 , 44, 108-118	3.2	0
107	Trunk movement compensation identified by inertial measurement units is associated with deficits in physical performance, muscle strength and functional capacity in people with hip osteoarthritis. <i>Clinical Biomechanics</i> , 2021 , 88, 105436	2.2	0
106	Neighbors-based prediction of physical function after total knee arthroplasty. <i>Scientific Reports</i> , 2021 , 11, 16719	4.9	
105	Lateral Corticospinal Tract and Dorsal Column Damage: Predictive Relationships With Motor and Sensory Scores at Discharge From Acute Rehabilitation After Spinal Cord Injury. <i>Archives of Physical Medicine and Rehabilitation</i> , 2021 ,	2.8	1
104	Association of Leisure-Time Physical Activity With Health-Related Quality of Life Among US Lung Cancer Survivors. <i>JNCI Cancer Spectrum</i> , 2021 , 5,	4.6	2

103	Implementation of a Remote Fidelity Oversight Program for a Multi-Site Exercise Intervention. <i>Innovation in Aging</i> , 2021 , 5, 295-295	0.1	
102	Physical Therapist Management of Total Knee Arthroplasty. <i>Physical Therapy</i> , 2020 , 100, 1603-1631	3.3	20
101	Home Health Rehabilitation Utilization Among Medicare Beneficiaries Following Critical Illness. <i>Journal of the American Geriatrics Society</i> , 2020 , 68, 1512-1519	5.6	7
100	Movement pattern biofeedback training after total knee arthroplasty: Randomized clinical trial protocol. <i>Contemporary Clinical Trials</i> , 2020 , 91, 105973	2.3	1
99	The Impact of Home Health Physical Therapy on Medicare Beneficiaries With a Primary Diagnosis of Dementia. <i>Journal of the American Geriatrics Society</i> , 2020 , 68, 867-871	5.6	7
98	Biobehavioral Intervention Targeting Physical Activity Behavior Change for Older Veterans after Nontraumatic Amputation: A Randomized Controlled Trial. <i>PM and R</i> , 2020 , 12, 957-966	2.2	2
97	Person-Centered Care and Physical Therapy: A "People-Like-Me" Approach. <i>Physical Therapy</i> , 2020 , 100, 99-106	3.3	7
96	Preoperative Activities of Daily Living Dependency is Associated With Higher 30-Day Readmission Risk for Older Adults After Total Joint Arthroplasty. <i>Clinical Orthopaedics and Related Research</i> , 2020 , 478, 231-237	2.2	2
95	Multiparticipant Rehabilitation in Skilled Nursing Facilities: An Observational Comparison Study. <i>Journal of the American Medical Directors Association</i> , 2020 , 21, 1920-1925	5.9	1
94	Reference chart for knee flexion following total knee arthroplasty: a novel tool for monitoring postoperative recovery. <i>BMC Musculoskeletal Disorders</i> , 2020 , 21, 482	2.8	1
93	Application of High-Intensity Functional Resistance Training in a Skilled Nursing Facility: An Implementation Study. <i>Physical Therapy</i> , 2020 , 100, 1746-1758	3.3	2
92	Psychometric properties of the 4-meter walk test after total knee arthroplasty. <i>Disability and Rehabilitation</i> , 2020 , 1-7	2.4	3
91	Benefits of direct patient discharge to outpatient physical therapy after total knee arthroplasty. <i>Disability and Rehabilitation</i> , 2020 , 42, 660-666	2.4	4
90	Development of a reference chart to monitor postoperative swelling following total knee arthroplasty. <i>Disability and Rehabilitation</i> , 2020 , 42, 1767-1774	2.4	5
89	Simulated hip abductor strengthening reduces peak joint contact forces in patients with total hip arthroplasty. <i>Journal of Biomechanics</i> , 2019 , 93, 18-27	2.9	10
88	Progressive Multicomponent Intervention for Older Adults in Home Health Settings Following Acute Hospitalization: Randomized Clinical Trial Protocol. <i>Physical Therapy</i> , 2019 , 99, 1141-1149	3.3	0
87	The relationship between lower extremity swelling, quadriceps strength, and functional performance following total knee arthroplasty. <i>Knee</i> , 2019 , 26, 382-391	2.6	12
86	Incorporating Specific Functional Strength Integration Techniques to Improve Functional Performance for Veterans After Total Hip Arthroplasty: Protocol for a Randomized Clinical Trial. <i>Physical Therapy</i> , 2019 , 99, 1453-1460	3.3	1

85	Conducting Clinical Research in Post-acute and Long-term Nursing Home Care Settings: Regulatory Challenges. <i>Journal of the American Medical Directors Association</i> , 2019 , 20, 798-803	5.9	10
84	Characteristics of young and lower functioning patients following total knee arthroplasty: a retrospective study. <i>BMC Musculoskeletal Disorders</i> , 2019 , 20, 483	2.8	2
83	Physical Function and Pre-Amputation Characteristics Explain Daily Step Count after Dysvascular Amputation. <i>PM and R</i> , 2019 , 11, 1050-1058	2.2	6
82	Side-to-Side Variability in Muscle Function With Parkinson Disease. <i>Topics in Geriatric Rehabilitation</i> , 2019 , 35, 231-236	0.7	
81	Peripheral Nociception Is Associated with Voluntary Activation Deficits and Quadriceps Weakness Following Total Knee Arthroplasty. <i>Journal of Bone and Joint Surgery - Series A</i> , 2019 , 101, 1539-1545	5.6	1
80	Trajectories of functional performance and muscle strength recovery differ after total knee and total hip replacement: a performance-based, longitudinal study. <i>International Journal of Rehabilitation Research</i> , 2019 , 42, 211-216	1.8	2
79	Influence of Diabetes Mellitus on the Recovery Trajectories of Function, Strength, and Self-Report Measures After Total Knee Arthroplasty. <i>Arthritis Care and Research</i> , 2019 , 71, 1059-1067	4.7	2
78	Impaired Physical Performance Predicts Hospitalization Risk for Participants in the Program of All-Inclusive Care for the Elderly. <i>Physical Therapy</i> , 2019 , 99, 28-36	3.3	6
77	Stiff knee gait may increase risk of second total knee arthroplasty. <i>Journal of Orthopaedic Research</i> , 2019 , 37, 397-402	3.8	14
76	Improving Physical Activity Through Adjunct Telerehabilitation Following Total Knee Arthroplasty: Randomized Controlled Trial Protocol. <i>Physical Therapy</i> , 2019 , 99, 37-45	3.3	8
75	Predictors of Functional Change in a Skilled Nursing Facility Population. <i>Journal of Geriatric Physical Therapy</i> , 2019 , 42, 189-195	3.2	11
74	Dementia, Comorbidity, and Physical Function in the Program of All-Inclusive Care for the Elderly. <i>Journal of Geriatric Physical Therapy</i> , 2019 , 42, E1-E6	3.2	13
73	Involvement of Acute Care Physical Therapists in Care Transitions for Older Adults Following Acute Hospitalization: A Cross-sectional National Survey. <i>Journal of Geriatric Physical Therapy</i> , 2019 , 42, E73-E80	3.2	5
72	The impact of hip implant alignment on muscle and joint loading during dynamic activities. <i>Clinical Biomechanics</i> , 2018 , 53, 93-100	2.2	16
71	Relationships Among Perceived Functional Capacity, Self-Efficacy, and Disability After Dysvascular Amputation. <i>PM and R</i> , 2018 , 10, 1056-1061	2.2	6
70	Clinical Use of Neuromuscular Electrical Stimulation for Neuromuscular Rehabilitation: What Are We Overlooking?. <i>Archives of Physical Medicine and Rehabilitation</i> , 2018 , 99, 806-812	2.8	56
69	A Feasibility Study for Improved Physical Activity After Total Knee Arthroplasty. <i>Journal of Aging and Physical Activity</i> , 2018 , 26, 7-13	1.6	21
68	Performance on a Clinical Quadriceps Activation Battery Is Related to a Laboratory Measure of Activation and Recovery After Total Knee Arthroplasty. <i>Archives of Physical Medicine and Rehabilitation</i> , 2018 , 99, 99-106	2.8	5

67	Behavior-Change Intervention Targeting Physical Function, Walking, and Disability After Dysvascular Amputation: A Randomized Controlled Pilot Trial. <i>Archives of Physical Medicine and Rehabilitation</i> , 2018 , 99, 2160-2167	2.8	16
66	Response to Letter to the Editor on "Arthrofibrosis Associated With Total Knee Arthroplasty". <i>Journal of Arthroplasty</i> , 2018 , 33, 968-969	4.4	1
65	Home-Health-Care Physical Therapy Improves Early Functional Recovery of Medicare Beneficiaries After Total Knee Arthroplasty. <i>Journal of Bone and Joint Surgery - Series A</i> , 2018 , 100, 1728-1734	5.6	19
64	Arthrofibrosis Associated With Total Knee Arthroplasty. <i>Journal of Arthroplasty</i> , 2017 , 32, 2604-2611	4.4	97
63	Physical activity behavior change for older veterans after dysvascular amputation. <i>Contemporary Clinical Trials</i> , 2017 , 55, 10-15	2.3	6
62	Magnitude of Deformity Correction May Influence Recovery of Quadriceps Strength After Total Knee Arthroplasty. <i>Journal of Arthroplasty</i> , 2017 , 32, 2730-2737	4.4	5
61	Preparing for the Future of Post-Acute Care: A Metasynthesis of Qualitative Studies.. <i>Home Health Care Management and Practice</i> , 2017 , 29, 70-80	0.9	1
60	Influence of Hip Abductor Strength on Functional Outcomes Before and After Total Knee Arthroplasty: Post Hoc Analysis of a Randomized Controlled Trial. <i>Physical Therapy</i> , 2017 , 97, 896-903	3.3	14
59	Early High-Intensity Versus Low-Intensity Rehabilitation After Total Knee Arthroplasty: A Randomized Controlled Trial. <i>Arthritis Care and Research</i> , 2017 , 69, 1360-1368	4.7	37
58	Use of post-acute care after hospital discharge in urban and rural hospitals 2017 , 5, 16-22		15
57	Progressive multi-component home-based physical therapy for deconditioned older adults following acute hospitalization: a pilot randomized controlled trial. <i>Clinical Rehabilitation</i> , 2016 , 30, 776-85	3.3	14
56	Does Tourniquet Use in TKA Affect Recovery of Lower Extremity Strength and Function? A Randomized Trial. <i>Clinical Orthopaedics and Related Research</i> , 2016 , 474, 69-77	2.2	113
55	Performance-Based Versus Self-Reported Outcomes Using the Hip Disability and Osteoarthritis Outcome Score After Total Hip Arthroplasty. <i>American Journal of Physical Medicine and Rehabilitation</i> , 2016 , 95, 132-8	2.6	29
54	Multi-Joint Compensatory Effects of Unilateral Total Knee Arthroplasty During High-Demand Tasks. <i>Annals of Biomedical Engineering</i> , 2016 , 44, 2529-2541	4.7	19
53	Role of Physical Therapists in Reducing Hospital Readmissions: Optimizing Outcomes for Older Adults During Care Transitions From Hospital to Community. <i>Physical Therapy</i> , 2016 , 96, 1125-34	3.3	41
52	Effects of neuromuscular reeducation on hip mechanics and functional performance in patients after total hip arthroplasty: A case series. <i>Clinical Biomechanics</i> , 2016 , 32, 49-55	2.2	10
51	Physical activity, ambulation, and comorbidities in people with diabetes and lower-limb amputation. <i>Journal of Rehabilitation Research and Development</i> , 2016 , 53, 1069-1078		19
50	Assessing Radiographic Knee Osteoarthritis: An Online Training Tutorial for the Kellgren-Lawrence Grading Scale. <i>MedEdPORTAL: the Journal of Teaching and Learning Resources</i> , 2016 , 12, 10503	1.2	5

49	Current Telerehabilitation Research With Older Adults at Home: An Integrative Review. <i>Journal of Gerontological Nursing</i> , 2016 , 42, 15-20	1.2	23
48	Explaining modified 2-min walk test outcomes in male Veterans with traumatic or nontraumatic lower-limb amputation. <i>Journal of Rehabilitation Research and Development</i> , 2016 , 53, 1035-1044		3
47	Men and Women Demonstrate Differences in Early Functional Recovery After Total Knee Arthroplasty. <i>Archives of Physical Medicine and Rehabilitation</i> , 2016 , 97, 1154-62	2.8	15
46	Functional Outcomes After the Prosthetic Training Phase of Rehabilitation After Dysvascular Lower Extremity Amputation. <i>PM and R</i> , 2015 , 7, 1118-1126	2.2	37
45	Reduced rate of knee extensor torque development in older adults with knee osteoarthritis is associated with intrinsic muscle contractile deficits. <i>Experimental Gerontology</i> , 2015 , 72, 16-21	4.5	18
44	Effects of Weight-Bearing Biofeedback Training on Functional Movement Patterns Following Total Knee Arthroplasty: A Randomized Controlled Trial. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2015 , 45, 647-55	4.2	33
43	Rethinking Hospital-Associated Deconditioning: Proposed Paradigm Shift. <i>Physical Therapy</i> , 2015 , 95, 1307-15	3.3	62
42	Poster 252 Modifiable Gait Parameters Associated with Functional Mobility in Lower Limb Amputees. <i>PM and R</i> , 2015 , 7, S174-S174	2.2	
41	Physical activity after total knee arthroplasty: A critical review. <i>World Journal of Orthopedics</i> , 2015 , 6, 614-22	2.2	38
40	Risk factors for delayed inpatient functional recovery after total knee arthroplasty. <i>BioMed Research International</i> , 2015 , 2015, 167643	3	15
39	Physical Function and Hospital Readmissions. <i>JAMA Internal Medicine</i> , 2015 , 175, 1722	11.5	1
38	Measuring movement symmetry using tibial-mounted accelerometers for people recovering from total knee arthroplasty. <i>Clinical Biomechanics</i> , 2015 , 30, 732-7	2.2	9
37	Poster 42 Early Physical Function for Patients with Type 2 Diabetes Following Transtibial Amputation. <i>PM and R</i> , 2014 , 6, S196	2.2	
36	Strength and functional deficits in individuals with hip osteoarthritis compared to healthy, older adults. <i>Disability and Rehabilitation</i> , 2014 , 36, 307-12	2.4	48
35	Corticospinal and intracortical excitability of the quadriceps in patients with knee osteoarthritis. <i>Experimental Brain Research</i> , 2014 , 232, 3991-9	2.3	28
34	Quadriceps/hamstrings co-activation increases early after total knee arthroplasty. <i>Knee</i> , 2014 , 21, 1115-9.6		32
33	Predicting functional performance and range of motion outcomes after total knee arthroplasty. <i>American Journal of Physical Medicine and Rehabilitation</i> , 2014 , 93, 579-85	2.6	56
32	Preliminary investigation of rate of torque development deficits following total knee arthroplasty. <i>Knee</i> , 2014 , 21, 382-6	2.6	19

31	Muscle strength and functional recovery during the first year after THA. <i>Clinical Orthopaedics and Related Research</i> , 2014 , 472, 654-64	2.2	99
30	Management of symptoms of Restless Legs Syndrome with use of a traction straight leg raise: a preliminary case series. <i>Manual Therapy</i> , 2013 , 18, 299-302		10
29	Muscle activation and coactivation during five-time-sit-to-stand movement in patients undergoing total knee arthroplasty. <i>Journal of Electromyography and Kinesiology</i> , 2013 , 23, 1485-93	2.5	31
28	Factors predicting weight-bearing asymmetry 1month after unilateral total knee arthroplasty: a cross-sectional study. <i>Gait and Posture</i> , 2013 , 37, 363-7	2.6	17
27	Corticospinal and intracortical excitability of the quadriceps in active older and younger healthy adults. <i>Archives of Gerontology and Geriatrics</i> , 2013 , 56, 279-84	4	31
26	Single-step test for unilateral limb ability following total knee arthroplasty. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2013 , 43, 66-73	4.2	12
25	Minimally invasive total knee arthroplasty: surgical implications for recovery. <i>Journal of Knee Surgery</i> , 2013 , 26, 195-201	2.4	11
24	Does delay in total knee arthroplasty impact postoperative performance? A case-based illustration. <i>Journal of Knee Surgery</i> , 2013 , 26 Suppl 1, S81-8	2.4	
23	Slower walking speed forecasts increased postoperative morbidity and 1-year mortality across surgical specialties. <i>Annals of Surgery</i> , 2013 , 258, 582-8; discussion 588-90	7.8	135
22	Linear and curvilinear relationship between knee range of motion and physical functioning in people with knee osteoarthritis: a cross-sectional study. <i>PLoS ONE</i> , 2013 , 8, e76173	3.7	6
21	Minimally invasive total knee arthroplasty improves early knee strength but not functional performance: a randomized controlled trial. <i>Journal of Arthroplasty</i> , 2012 , 27, 1812-1819.e2	4.4	28
20	Quadriceps muscle weakness, activation deficits, and fatigue with Parkinson disease. <i>Neurorehabilitation and Neural Repair</i> , 2012 , 26, 533-41	4.7	56
19	Predicting poor physical performance after total knee arthroplasty. <i>Journal of Orthopaedic Research</i> , 2012 , 30, 1805-10	3.8	26
18	Relationship between intensity of quadriceps muscle neuromuscular electrical stimulation and strength recovery after total knee arthroplasty. <i>Physical Therapy</i> , 2012 , 92, 1187-96	3.3	58
17	Restoration of physical function in patients following total knee arthroplasty: an update on rehabilitation practices. <i>Current Opinion in Rheumatology</i> , 2012 , 24, 208-14	5.3	41
16	Importance of attenuating quadriceps activation deficits after total knee arthroplasty. <i>Exercise and Sport Sciences Reviews</i> , 2012 , 40, 95-101	6.7	30
15	Early neuromuscular electrical stimulation to improve quadriceps muscle strength after total knee arthroplasty: a randomized controlled trial. <i>Physical Therapy</i> , 2012 , 92, 210-26	3.3	165
14	Walking Speed in the Total Joint Arthroplasty Population. <i>Topics in Geriatric Rehabilitation</i> , 2012 , 28, 104-109	0.7	3

13	Muscle strength loss in the lower limb after total knee arthroplasty. <i>American Journal of Physical Medicine and Rehabilitation</i> , 2012 , 91, 220-6; quiz 227-30	2.6	32
12	Continuous femoral nerve blocks: decreasing local anesthetic concentration to minimize quadriceps femoris weakness. <i>Anesthesiology</i> , 2012 , 116, 665-72	4.3	60
11	Comparison of self-reported knee injury and osteoarthritis outcome score to performance measures in patients after total knee arthroplasty. <i>PM and R</i> , 2011 , 3, 541-9; quiz 549	2.2	105
10	Weight-bearing asymmetry during sit-stand transitions related to impairment and functional mobility after total knee arthroplasty. <i>Archives of Physical Medicine and Rehabilitation</i> , 2011 , 92, 1624-9	2.8	40
9	Early high-intensity rehabilitation following total knee arthroplasty improves outcomes. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2011 , 41, 932-41	4.2	94
8	Outcomes before and after total knee arthroplasty compared to healthy adults. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2010 , 40, 559-67	4.2	247
7	Osteoarthritis in women: effects of estrogen, obesity and physical activity. <i>Women's Health</i> , 2010 , 6, 601-15	3	28
6	Weight-bearing asymmetry in relation to measures of impairment and functional mobility for people with knee osteoarthritis. <i>Archives of Physical Medicine and Rehabilitation</i> , 2010 , 91, 1524-8	2.8	74
5	Quadriceps and hamstrings muscle dysfunction after total knee arthroplasty. <i>Clinical Orthopaedics and Related Research</i> , 2010 , 468, 2460-8	2.2	106
4	Impact of body mass index on functional performance after total knee arthroplasty. <i>Journal of Arthroplasty</i> , 2010 , 25, 1104-9	4.4	50
3	Functional performance and muscle strength phenotypes in men and women with Danon disease. <i>Muscle and Nerve</i> , 2010 , 42, 908-14	3.4	12
2	Continuous peripheral nerve blocks: is local anesthetic dose the only factor, or do concentration and volume influence infusion effects as well?. <i>Anesthesiology</i> , 2010 , 112, 347-54	4.3	76
1	Surgical Management and Rehabilitation of Knee Arthroplasty 2010 , 1-63		