

Maohua Miao

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6281327/publications.pdf>

Version: 2024-02-01

18
papers

486
citations

1163117

8
h-index

888059

17
g-index

18
all docs

18
docs citations

18
times ranked

689
citing authors

#	ARTICLE	IF	CITATIONS
1	Preconceptional paternal alcohol consumption and the risk of child behavioral problems: a prospective cohort study. <i>Scientific Reports</i> , 2022, 12, 1508.	3.3	5
2	Association of pre-pregnancy body mass index and gestational weight gain with neonatal anogenital distance in a Chinese birth cohort. <i>Reproductive Health</i> , 2022, 19, .	3.1	3
3	Effects of prenatal exposure to polybrominated diphenyl ethers (PBDEs) on the second to fourth digit ratio in children aged 4 years. <i>International Journal of Hygiene and Environmental Health</i> , 2021, 231, 113639.	4.3	8
4	Risk of intellectual disability and maternal history of spontaneous abortion: a nationwide cohort study. <i>Developmental Medicine and Child Neurology</i> , 2021, 63, 831-838.	2.1	3
5	Association between prenatal exposure to polybrominated diphenyl ethers and anogenital distance in girls at ages 0–4 years. <i>International Journal of Hygiene and Environmental Health</i> , 2021, 233, 113706.	4.3	7
6	Prenatal exposure to bisphenol analogues and digit ratio in children at ages 4 and 6 years: A birth cohort study. <i>Environmental Pollution</i> , 2021, 278, 116820.	7.5	7
7	Effects of low lead exposure on sperm quality and sperm DNA methylation in adult men. <i>Cell and Bioscience</i> , 2021, 11, 150.	4.8	7
8	Association Between Neonatal Thyroid Function and Anogenital Distance from Birth to 48 Months of Age. <i>Frontiers in Endocrinology</i> , 2021, 12, 736505.	3.5	0
9	Prenatal exposure to residential PM2.5 and its chemical constituents and weight in preschool children: A longitudinal study from Shanghai, China. <i>Environment International</i> , 2021, 154, 106580.	10.0	14
10	Prenatal exposure to residential PM2.5 and anogenital distance in infants at birth: A birth cohort study from Shanghai, China. <i>Environmental Pollution</i> , 2020, 264, 114684.	7.5	7
11	Maternal plasma concentrations of perfluoroalkyl and polyfluoroalkyl substances during pregnancy and anogenital distance in male infants. <i>Human Reproduction</i> , 2019, 34, 1356-1368.	0.9	32
12	Prenatal polybrominated diphenyl ethers exposure and anogenital distance in boys from a Shanghai birth cohort. <i>International Journal of Hygiene and Environmental Health</i> , 2019, 222, 513-523.	4.3	40
13	Prenatal plasma concentrations of Perfluoroalkyl and polyfluoroalkyl substances and neuropsychological development in children at four years of age. <i>Environmental Health</i> , 2019, 18, 53.	4.0	33
14	Preconceptional paternal antiepileptic drugs use and risk of congenital anomalies in offspring: a nationwide cohort study. <i>European Journal of Epidemiology</i> , 2019, 34, 651-660.	5.7	9
15	Association Between Paternal Alcohol Consumption Before Conception and Anogenital Distance of Offspring. <i>Alcoholism: Clinical and Experimental Research</i> , 2018, 42, 735-742.	2.4	20
16	Maternal exposure to bisphenol A and anogenital distance throughout infancy: A longitudinal study from Shanghai, China. <i>Environment International</i> , 2018, 121, 269-275.	10.0	63
17	Determinants of plasma concentrations of perfluoroalkyl and polyfluoroalkyl substances in pregnant women from a birth cohort in Shanghai, China. <i>Environment International</i> , 2018, 119, 165-173.	10.0	98
18	In utero exposure to bisphenol-A and anogenital distance of male offspring. <i>Birth Defects Research Part A: Clinical and Molecular Teratology</i> , 2011, 91, 867-872.	1.6	130