

# Pamela Cameron

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6284763/publications.pdf>

Version: 2024-02-01

13  
papers

718  
citations

840776

11  
h-index

1125743

13  
g-index

13  
all docs

13  
docs citations

13  
times ranked

1192  
citing authors

#	ARTICLE	IF	CITATIONS
1	Inhibition of Lipopolysaccharide-Induced Macrophage IL-12 Production by <i>Leishmania mexicana</i> Amastigotes: The Role of Cysteine Peptidases and the NF- $\kappa$ B Signaling Pathway. <i>Journal of Immunology</i> , 2004, 173, 3297-3304.	0.8	164
2	Screening methods used to determine the anti-microbial properties of Aloe vera inner gel. <i>Methods</i> , 2007, 42, 315-320.	3.8	112
3	EspP, a Type V-secreted serine protease of enterohaemorrhagic <i>Escherichia coli</i> O157:H7, influences intestinal colonization of calves and adherence to bovine primary intestinal epithelial cells. <i>FEMS Microbiology Letters</i> , 2007, 271, 258-264.	1.8	75
4	In Vitro Susceptibilities of <i>Shigella flexneri</i> and <i>Streptococcus pyogenes</i> to Inner Gel of Aloe barbadensis Miller. <i>Antimicrobial Agents and Chemotherapy</i> , 2003, 47, 1137-1139.	3.2	70
5	Silver Nanoparticles Decrease the Viability of <i>Cryptosporidium parvum</i> Oocysts. <i>Applied and Environmental Microbiology</i> , 2016, 82, 431-437.	3.1	64
6	The inner gel component of Aloe vera suppresses bacterial-induced pro-inflammatory cytokines from human immune cells. <i>Methods</i> , 2007, 42, 388-393.	3.8	51
7	The role of intracellular Ca <sup>2+</sup> in the regulation of proteinase-activated receptor-2 mediated nuclear factor kappa B signalling in keratinocytes. <i>British Journal of Pharmacology</i> , 2005, 145, 535-544.	5.4	49
8	Conditional expression of MAP kinase phosphatase-2 protects against genotoxic stress-induced apoptosis by binding and selective dephosphorylation of nuclear activated c-jun N-terminal kinase. <i>Cellular Signalling</i> , 2005, 17, 1254-1264.	3.6	46
9	Verotoxin activates mitogen-activated protein kinase in human peripheral blood monocytes: role in apoptosis and proinflammatory cytokine release. <i>British Journal of Pharmacology</i> , 2003, 140, 1320-1330.	5.4	41
10	Crowdfunding genomics and bioinformatics. <i>Genome Biology</i> , 2013, 14, 134.	9.6	21
11	Essential Role for Verotoxin in Sustained Stress-Activated Protein Kinase and Nuclear Factor Kappa B Signaling, Stimulated by <i>Escherichia coli</i> O157:H7 in Vero Cells. <i>Infection and Immunity</i> , 2002, 70, 5370-5380.	2.2	13
12	Phosphorylation of the epidermal growth factor receptor (EGFR) is essential for interleukin-8 release from intestinal epithelial cells in response to challenge with <i>Escherichia coli</i> O157:H7 flagellin. <i>Microbiology (United Kingdom)</i> , 2011, 157, 2339-2347.	1.8	9
13	Verotoxin-2 Activates Mitogen-Activated Protein Kinases in Bovine Adherent Peripheral Blood Mononuclear Cells. <i>Journal of Comparative Pathology</i> , 2012, 147, 20-23.	0.4	3