
Pamela Cameron

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6284763/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Silver Nanoparticles Decrease the Viability of Cryptosporidium parvum Oocysts. Applied and Environmental Microbiology, 2016, 82, 431-437. | 3.1 | 64 |
| 2 | Crowdfunding genomics and bioinformatics. Genome Biology, 2013, 14, 134. | 9.6 | 21 |
| 3 | Verotoxin-2 Activates Mitogen-Activated Protein Kinases in Bovine Adherent Peripheral Blood Mononuclear Cells. Journal of Comparative Pathology, 2012, 147, 20-23. | 0.4 | 3 |
| 4 | Phosphorylation of the epidermal growth factor receptor (EGFR) is essential for interleukin-8 release from intestinal epithelial cells in response to challenge with Escherichia coli O157 : H7 flagellin. Microbiology (United Kingdom), 2011, 157, 2339-2347. | 1.8 | 9 |
| 5 | The inner gel component of Aloe vera suppresses bacterial-induced pro-inflammatory cytokines from human immune cells. Methods, 2007, 42, 388-393. | 3.8 | 51 |
| 6 | Screening methods used to determine the anti-microbial properties of Aloe vera inner gel. Methods, 2007, 42, 315-320. | 3.8 | 112 |
| 7 | EspP, a Type V-secreted serine protease of enterohaemorrhagicEscherichia coliO157:H7, influences intestinal colonization of calves and adherence to bovine primary intestinal epithelial cells. FEMS Microbiology Letters, 2007, 271, 258-264. | 1.8 | 75 |
| 8 | The role of intracellular Ca2+ in the regulation of proteinase-activated receptor-2 mediated nuclear factor kappa B signalling in keratinocytes. British Journal of Pharmacology, 2005, 145, 535-544. | 5.4 | 49 |
| 9 | Conditional expression of MAP kinase phosphatase-2 protects against genotoxic stress-induced apoptosis by binding and selective dephosphorylation of nuclear activated c-jun N-terminal kinase. Cellular Signalling, 2005, 17, 1254-1264. | 3.6 | 46 |
| 10 | Inhibition of Lipopolysaccharide-Induced Macrophage IL-12 Production by <i>Leishmania mexicana</i> Amastigotes: The Role of Cysteine Peptidases and the NF-ήB Signaling Pathway. Journal of Immunology, 2004, 173, 3297-3304. | 0.8 | 164 |
| 11 | Verotoxin activates mitogen-activated protein kinase in human peripheral blood monocytes: role in apoptosis and proinflammatory cytokine release. British Journal of Pharmacology, 2003, 140, 1320-1330. | 5.4 | 41 |
| 12 | In Vitro Susceptibilities of Shigella flexneri and Streptococcus pyogenes to Inner Gel of Aloe barbadensis Miller. Antimicrobial Agents and Chemotherapy, 2003, 47, 1137-1139. | 3.2 | 70 |
| 13 | Essential Role for Verotoxin in Sustained Stress-Activated Protein Kinase and Nuclear Factor Kappa B Signaling, Stimulated by Escherichia coli O157:H7 in Vero Cells. Infection and Immunity, 2002, 70, | 2.2 | 13 |