

# Sara Ceccarelli

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/628497/sara-ceccarelli-publications-by-year.pdf>

**Version:** 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

22  
papers

771  
citations

17  
h-index

22  
g-index

22  
ext. papers

897  
ext. citations

5.8  
avg, IF

3.44  
L-index

| #  | Paper   | IF   | Citations |
|----|---|------|-----------|
| 22 | Focal adhesion kinase depletion reduces human hepatocellular carcinoma growth by repressing enhancer of zeste homolog 2. <i>Cell Death and Differentiation</i> , <b>2017</b> , 24, 889-902                      | 12.7 | 36        |
| 21 | Human B-cell memory is shaped by age- and tissue-specific T-independent and GC-dependent events. <i>European Journal of Immunology</i> , <b>2017</b> , 47, 327-344  | 6.1  | 34        |
| 20 | Increased serum IgM, immunodeficiency, and autoimmunity: A clinical series. <i>International Journal of Immunopathology and Pharmacology</i> , <b>2015</b> , 28, 547-56   | 3    | 6         |
| 19 | Targeting FGF19 binding to its receptor system: a novel therapeutic approach for hepatocellular carcinoma. <i>Hepatology</i> , <b>2015</b> , 62, 1324   | 11.2 | 1         |
| 18 | Serum Bile Acid Levels in Children With Nonalcoholic Fatty Liver Disease. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , <b>2015</b> , 61, 85-90  | 2.8  | 31        |
| 17 | LPS-induced TNF- $\alpha$ factor mediates pro-inflammatory and pro-fibrogenic pattern in non-alcoholic fatty liver disease. <i>Oncotarget</i> , <b>2015</b> , 6, 41434-52                                       | 3.3  | 78        |
| 16 | MicroRNAs as controlled systems and controllers in non-alcoholic fatty liver disease. <i>World Journal of Gastroenterology</i> , <b>2014</b> , 20, 15079-86   | 5.6  | 40        |
| 15 | Plasma high mobility group box 1 protein reflects fibrosis in pediatric nonalcoholic fatty liver disease. <i>Expert Review of Molecular Diagnostics</i> , <b>2014</b> , 14, 763-71                              | 3.8  | 19        |
| 14 | Plasma levels of homocysteine and cysteine increased in pediatric NAFLD and strongly correlated with severity of liver damage. <i>International Journal of Molecular Sciences</i> , <b>2014</b> , 15, 21202-14  | 6.3  | 59        |
| 13 | Toll-like receptor-mediated signaling cascade as a regulator of the inflammation network during alcoholic liver disease. <i>World Journal of Gastroenterology</i> , <b>2014</b> , 20, 16443-51                  | 5.6  | 24        |
| 12 | Gut-liver axis and fibrosis in nonalcoholic fatty liver disease: an input for novel therapies. <i>Digestive and Liver Disease</i> , <b>2013</b> , 45, 543-51  | 3.3  | 60        |
| 11 | Dual role of microRNAs in NAFLD. <i>International Journal of Molecular Sciences</i> , <b>2013</b> , 14, 8437-55   | 6.3  | 51        |
| 10 | EZH2 down-regulation exacerbates lipid accumulation and inflammation in in vitro and in vivo NAFLD. <i>International Journal of Molecular Sciences</i> , <b>2013</b> , 14, 24154-68                             | 6.3  | 39        |
| 9  | Retinoids counteract insulin resistance and liver steatosis: whats the potential mechanism?. <i>Hepatology</i> , <b>2013</b> , 58, 1185   | 11.2 | 0         |
| 8  | Association between Serum Atypical Fibroblast Growth Factors 21 and 19 and Pediatric Nonalcoholic Fatty Liver Disease. <i>PLoS ONE</i> , <b>2013</b> , 8, e67160  | 3.7  | 69        |
| 7  | Markers of activated inflammatory cells correlate with severity of liver damage in children with nonalcoholic fatty liver disease. <i>International Journal of Molecular Medicine</i> , <b>2012</b> , 30, 49-56 | 4.4  | 46        |
| 6  | T regulatory cell number and function: the autoimmune traits in liver diseases. <i>Journal of Hepatology</i> , <b>2012</b> , 57, 1398-9; author reply 1399-400  | 13.4 |           |

|   |  |      |    |
|---|--|------|----|
| 5 | Causative role of gut microbiota in non-alcoholic fatty liver disease pathogenesis. <i>Frontiers in Cellular and Infection Microbiology</i> , <b>2012</b> , 2, 132   | 5.9  | 36 |
| 4 | Emodin prevents intrahepatic fat accumulation, inflammation and redox status imbalance during diet-induced hepatosteatosis in rats. <i>International Journal of Molecular Sciences</i> , <b>2012</b> , 13, 2276-89 | 6.3  | 40 |
| 3 | Hepatic stellate cell proliferation: a potential role of protein kinase R. <i>Hepatology</i> , <b>2011</b> , 54, 1484-5; author reply 1485-6   | 11.2 | 1  |
| 2 | TLR ligation triggers somatic hypermutation in transitional B cells inducing the generation of IgM memory B cells. <i>Journal of Immunology</i> , <b>2010</b> , 185, 7293-301                                      | 5.3  | 59 |
| 1 | Targeted expression of RALT in mouse skin inhibits epidermal growth factor receptor signalling and generates a Waved-like phenotype. <i>EMBO Reports</i> , <b>2005</b> , 6, 755-61                                 | 6.5  | 42 |