

# Joshua S Waxman

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6287163/publications.pdf>

Version: 2024-02-01

40  
papers

1,565  
citations

394421

19  
h-index

315739

38  
g-index

47  
all docs

47  
docs citations

47  
times ranked

1873  
citing authors

#	ARTICLE	IF	CITATIONS
1	Somite morphogenesis is required for axial blood vessel formation during zebrafish embryogenesis. <i>ELife</i> , 2022, 11, .	6.0	5
2	Stx4 is required to regulate cardiomyocyte Ca <sup>2+</sup> handling during vertebrate cardiac development. <i>Human Genetics and Genomics Advances</i> , 2022, 3, 100115.	1.7	1
3	Atrial and Sinoatrial Node Development in the Zebrafish Heart. <i>Journal of Cardiovascular Development and Disease</i> , 2021, 8, 15.	1.6	10
4	Retinoic acid signaling restricts the size of the first heart field within the anterior lateral plate mesoderm. <i>Developmental Biology</i> , 2021, 473, 119-129.	2.0	22
5	Ccdc103 promotes myeloid cell proliferation and migration independent of motile cilia. <i>DMM Disease Models and Mechanisms</i> , 2021, 14, .	2.4	4
6	Patterning of vertebrate cardiac progenitor fields by retinoic acid signaling. <i>Genesis</i> , 2021, 59, e23458.	1.6	5
7	Origin and evolutionary landscape of Nr2f transcription factors across Metazoa. <i>PLoS ONE</i> , 2021, 16, e0254282.	2.5	6
8	Elevated Hoxb5b Expands Vagal Neural Crest Pool and Blocks Enteric Neuronal Development in Zebrafish. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 803370.	3.7	7
9	Pbx4 limits heart size and fosters arch artery formation through partitioning second heart field progenitors and restricting proliferation. <i>Development (Cambridge)</i> , 2020, 147, .	2.5	10
10	Retinoic Acid Signaling and Heart Development. <i>Sub-Cellular Biochemistry</i> , 2020, 95, 119-149.	2.4	8
11	Reiterative Mechanisms of Retinoic Acid Signaling during Vertebrate Heart Development. <i>Journal of Developmental Biology</i> , 2019, 7, 11.	1.7	23
12	HDAC1-mediated repression of the retinoic acid-responsive gene rippy3 promotes second heart field development. <i>PLoS Genetics</i> , 2019, 15, e1008165.	3.5	16
13	Enhancing regeneration after acute kidney injury by promoting cellular dedifferentiation in zebrafish. <i>DMM Disease Models and Mechanisms</i> , 2019, 12, .	2.4	21
14	Nr2f-dependent allocation of ventricular cardiomyocyte and pharyngeal muscle progenitors. <i>PLoS Genetics</i> , 2019, 15, e1007962.	3.5	21
15	In Silico Identification and Experimental Validation of (â <sup>*</sup> )-Muqubilin A, a Marine Norterpene Peroxide, as PPAR $\alpha$ / $\beta$ -RXR $\alpha$ Agonist and RAR $\alpha$ Positive Allosteric Modulator. <i>Marine Drugs</i> , 2019, 17, 110.	4.6	11
16	Direct activation of chordoblasts by retinoic acid is required for segmented centra mineralization during zebrafish spine development. <i>Development (Cambridge)</i> , 2018, 145, .	2.5	29
17	Nr2f1a balances atrial chamber and atrioventricular canal size via BMP signaling-independent and -dependent mechanisms. <i>Developmental Biology</i> , 2018, 434, 7-14.	2.0	24
18	Wnt signaling balances specification of the cardiac and pharyngeal muscle fields. <i>Mechanisms of Development</i> , 2017, 143, 32-41.	1.7	18

#	ARTICLE	IF	CITATIONS
19	Retinoic acid and meiosis induction in adult versus embryonic gonads of medaka. <i>Scientific Reports</i> , 2016, 6, 34281.	3.3	27
20	Cyp26 Enzymes Facilitate Second Heart Field Progenitor Addition and Maintenance of Ventricular Integrity. <i>PLoS Biology</i> , 2016, 14, e2000504.	5.6	38
21	Rdh10a Provides a Conserved Critical Step in the Synthesis of Retinoic Acid during Zebrafish Embryogenesis. <i>PLoS ONE</i> , 2015, 10, e0138588.	2.5	17
22	Excessive feedback of Cyp26a1 promotes cell non-autonomous loss of retinoic acid signaling. <i>Developmental Biology</i> , 2015, 405, 47-55.	2.0	34
23	Input overload: Contributions of retinoic acid signaling feedback mechanisms to heart development and teratogenesis. <i>Developmental Dynamics</i> , 2015, 244, 513-523.	1.8	36
24	Retinoic acid negatively regulates dact3b expression in the hindbrain of zebrafish embryos. <i>Gene Expression Patterns</i> , 2014, 16, 122-129.	0.8	5
25	Cyp26 enzymes are required to balance the cardiac and vascular lineages within the anterior lateral plate mesoderm. <i>Development (Cambridge)</i> , 2014, 141, 1638-1648.	2.5	34
26	Tcf7l1 proteins cell autonomously restrict cardiomyocyte and promote endothelial specification in zebrafish. <i>Developmental Biology</i> , 2013, 380, 199-210.	2.0	7
27	Transgenic retinoic acid sensor lines in zebrafish indicate regions of available embryonic retinoic acid. <i>Developmental Dynamics</i> , 2013, 242, 989-1000.	1.8	27
28	Depletion of Retinoic Acid Receptors Initiates a Novel Positive Feedback Mechanism that Promotes Teratogenic Increases in Retinoic Acid. <i>PLoS Genetics</i> , 2013, 9, e1003689.	3.5	52
29	Distinct phases of Wnt/ $\beta$ -catenin signaling direct cardiomyocyte formation in zebrafish. <i>Developmental Biology</i> , 2012, 361, 364-376.	2.0	34
30	Zebrafish retinoic acid receptors function as context-dependent transcriptional activators. <i>Developmental Biology</i> , 2011, 352, 128-140.	2.0	40
31	Restraint of Fgf8 signaling by retinoic acid signaling is required for proper heart and forelimb formation. <i>Developmental Biology</i> , 2011, 358, 44-55.	2.0	41
32	John Morrill: Scientist, Educator, Friend (Nov. 20, 1929 - Aug. 9, 2010). <i>Molecular Reproduction and Development</i> , 2010, 77, n/a-n/a.	2.0	0
33	Increased Hox activity mimics the teratogenic effects of excess retinoic acid signaling. <i>Developmental Dynamics</i> , 2009, 238, 1207-1213.	1.8	49
34	Combinatorial roles for zebrafish retinoic acid receptors in the hindbrain, limbs and pharyngeal arches. <i>Developmental Biology</i> , 2009, 325, 60-70.	2.0	51
35	Hoxb5b Acts Downstream of Retinoic Acid Signaling in the Forelimb Field to Restrict Heart Field Potential in Zebrafish. <i>Developmental Cell</i> , 2008, 15, 923-934.	7.0	139
36	Comparison of the expression patterns of newly identified zebrafish retinoic acid and retinoid X receptors. <i>Developmental Dynamics</i> , 2007, 236, 587-595.	1.8	58

#	ARTICLE	IF	CITATIONS
37	Regulation of the early expression patterns of the zebrafish Dishevelled-interacting proteins Dapper1 and Dapper2. <i>Developmental Dynamics</i> , 2005, 233, 194-200.	1.8	9
38	Zebrafish Dapper1 and Dapper2 play distinct roles in Wnt-mediated developmental processes. <i>Development (Cambridge)</i> , 2004, 131, 5909-5921.	2.5	74
39	Dapper, a Dishevelled-Associated Antagonist of $\beta$ -Catenin and JNK Signaling, Is Required for Notochord Formation. <i>Developmental Cell</i> , 2002, 2, 449-461.	7.0	238
40	Zebrafish wnt8 Encodes Two Wnt8 Proteins on a Bicistronic Transcript and Is Required for Mesoderm and Neurectoderm Patterning. <i>Developmental Cell</i> , 2001, 1, 103-114.	7.0	313