

J Simon C Arthur

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6288433/j-simon-c-arthur-publications-by-year.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

160
papers

14,552
citations

60
h-index

118
g-index

169
ext. papers

16,301
ext. citations

7
avg. IF

6.36
L-index

#	Paper	IF	Citations
160	Nrf2 activation reprograms macrophage intermediary metabolism and suppresses the type I interferon response.. <i>IScience</i> , 2022 , 25, 103827	6.1	4
159	Salt inducible kinases 2 and 3 are required for thymic T cell development. <i>Scientific Reports</i> , 2021 , 11, 21550	4.9	0
158	The devil is in the detail: cell-specific role of PPAR α in ILC2 activation by IL-33. <i>Mucosal Immunology</i> , 2021 , 14, 544-546	9.2	1
157	Generation of a chemical genetic model for JAK3. <i>Scientific Reports</i> , 2021 , 11, 10093	4.9	3
156	Mitogen and Stress-activated Protein Kinase 1 Negatively Regulates Hippocampal Neurogenesis. <i>Neuroscience</i> , 2021 , 452, 228-234	3.9	6
155	Salt-inducible kinases are required for the IL-33-dependent secretion of cytokines and chemokines in mast cells. <i>Journal of Biological Chemistry</i> , 2021 , 296, 100428	5.4	2
154	Experience Recruits MSK1 to Expand the Dynamic Range of Synapses and Enhance Cognition. <i>Journal of Neuroscience</i> , 2020 , 40, 4644-4660	6.6	6
153	p38 MAPK signalling regulates cytokine production in IL-33 stimulated Type 2 Innate Lymphoid cells. <i>Scientific Reports</i> , 2020 , 10, 3479	4.9	13
152	Loss of Mef2D function enhances TLR induced IL-10 production in macrophages. <i>Bioscience Reports</i> , 2020 , 40,	4.1	3
151	The E3 ligase HOIL-1 catalyses ester bond formation between ubiquitin and components of the Myddosome in mammalian cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019 , 116, 13293-13298	11.5	43
150	Genetic variations in A20 DUB domain provide a genetic link to citrullination and neutrophil extracellular traps in systemic lupus erythematosus. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, 1363-1370	27.4	32
149	Distinct signals and immune cells drive liver pathology and glomerulonephritis in ABIN1[D485N] mice. <i>Life Science Alliance</i> , 2019 , 2,	5.8	7
148	IL-33 regulates cytokine production and neutrophil recruitment via the p38 MAPK-activated kinases MK2/3. <i>Immunology and Cell Biology</i> , 2019 , 97, 54-71	5	23
147	STAT3 activation by E6 is essential for the differentiation-dependent HPV18 life cycle. <i>PLoS Pathogens</i> , 2018 , 14, e1006975	7.6	33
146	Differential control of Toll-like receptor 4-induced interleukin-10 induction in macrophages and B cells reveals a role for p90 ribosomal S6 kinases. <i>Journal of Biological Chemistry</i> , 2018 , 293, 2302-2317	5.4	12
145	ABIN2 Function Is Required To Suppress DSS-Induced Colitis by a Tpl2-Independent Mechanism. <i>Journal of Immunology</i> , 2018 , 201, 3373-3382	5.3	8
144	Dimethyl fumarate is an allosteric covalent inhibitor of the p90 ribosomal S6 kinases. <i>Nature Communications</i> , 2018 , 9, 4344	17.4	20

143	Phosphorylated Histone 3 at Serine 10 Identifies Activated Spinal Neurons and Contributes to the Development of Tissue Injury-Associated Pain. <i>Scientific Reports</i> , 2017 , 7, 41221	4.9	8
142	Roles of the TRAF6 and Pellino E3 ligases in MyD88 and RANKL signaling. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017 , 114, E3481-E3489	11.5	61
141	Mir-132/212 is required for maturation of binocular matching of orientation preference and depth perception. <i>Nature Communications</i> , 2017 , 8, 15488	17.4	25
140	MSK1 regulates transcriptional induction of Arc/Arg3.1 in response to neurotrophins. <i>FEBS Open Bio</i> , 2017 , 7, 821-834	2.7	8
139	Loss of Functionally Redundant p38 Isoforms in T Cells Enhances Regulatory T Cell Induction. <i>Journal of Biological Chemistry</i> , 2017 , 292, 1762-1772	5.4	13
138	Inhibition of SIK2 and SIK3 during differentiation enhances the anti-inflammatory phenotype of macrophages. <i>Biochemical Journal</i> , 2017 , 474, 521-537	3.8	34
137	Mitogen- and Stress-Activated Protein Kinase 1 Regulates Status Epilepticus-Evoked Cell Death in the Hippocampus. <i>ASN Neuro</i> , 2017 , 9, 1759091417726607	5.3	7
136	P38 delta MAPK promotes breast cancer progression and lung metastasis by enhancing cell proliferation and cell detachment. <i>Oncogene</i> , 2017 , 36, 6649-6657	9.2	45
135	A Large Polysaccharide Produced by <i>Helicobacter hepaticus</i> Induces an Anti-inflammatory Gene Signature in Macrophages. <i>Cell Host and Microbe</i> , 2017 , 22, 733-745.e5	23.4	59
134	Identifying Inhibitors of Inflammation: A Novel High-Throughput MALDI-TOF Screening Assay for Salt-Inducible Kinases (SIKs). <i>SLAS Discovery</i> , 2017 , 22, 1193-1202	3.4	34
133	Beta Interferon Production Is Regulated by p38 Mitogen-Activated Protein Kinase in Macrophages via both MSK1/2- and Tristetraprolin-Dependent Pathways. <i>Molecular and Cellular Biology</i> , 2017 , 37,	4.8	12
132	The Kinase Function of MSK1 Regulates BDNF Signaling to CREB and Basal Synaptic Transmission, But Is Not Required for Hippocampal Long-Term Potentiation or Spatial Memory. <i>ENeuro</i> , 2017 , 4,	3.9	16
131	The PP4R1 sub-unit of protein phosphatase PP4 is essential for inhibition of NF- κ B by merkel polyomavirus small tumour antigen. <i>Oncotarget</i> , 2017 , 8, 25418-25432	3.3	24
130	Suppression of IRAK1 or IRAK4 Catalytic Activity, but Not Type 1 IFN Signaling, Prevents Lupus Nephritis in Mice Expressing a Ubiquitin Binding-Defective Mutant of ABIN1. <i>Journal of Immunology</i> , 2016 , 197, 4266-4273	5.3	32
129	A dominant mutation in MAPKAPK3, an actor of p38 signaling pathway, causes a new retinal dystrophy involving Bruch's membrane and retinal pigment epithelium. <i>Human Molecular Genetics</i> , 2016 , 25, 916-26	5.6	8
128	Targeted deletion of miR-132/-212 impairs memory and alters the hippocampal transcriptome. <i>Learning and Memory</i> , 2016 , 23, 61-71	2.8	60
127	The Chromatin Modifier MSK1/2 Suppresses Endocrine Cell Fates during Mouse Pancreatic Development. <i>PLoS ONE</i> , 2016 , 11, e0166703	3.7	6
126	Emerging Roles of the Mitogen and Stress Activated Kinases MSK1 and MSK2. <i>Frontiers in Cell and Developmental Biology</i> , 2016 , 4, 56	5.7	46

125	Micro-CT Imaging Reveals Mekk3 Heterozygosity Prevents Cerebral Cavernous Malformations in Ccm2-Deficient Mice. <i>PLoS ONE</i> , 2016 , 11, e0160833	3.7	12
124	The loop structure and the RNA helicase p72/DDX17 influence the processing efficiency of the mice miR-132. <i>Scientific Reports</i> , 2016 , 6, 22848	4.9	12
123	Dimethyl fumarate blocks pro-inflammatory cytokine production via inhibition of TLR induced M1 and K63 ubiquitin chain formation. <i>Scientific Reports</i> , 2016 , 6, 31159	4.9	60
122	Cerebral cavernous malformations arise from endothelial gain of MEKK3-KLF2/4 signalling. <i>Nature</i> , 2016 , 532, 122-6	50.4	164
121	Development of Selective Covalent Janus Kinase 3 Inhibitors. <i>Journal of Medicinal Chemistry</i> , 2015 , 58, 6589-606	8.3	70
120	Mitogen and stress-activated kinases 1/2 regulate ischemia-induced hippocampal progenitor cell proliferation and neurogenesis. <i>Neuroscience</i> , 2015 , 285, 292-302	3.9	18
119	Accelerated apoptotic death and in vivo turnover of erythrocytes in mice lacking functional mitogen- and stress-activated kinase MSK1/2. <i>Scientific Reports</i> , 2015 , 5, 17316	4.9	23
118	Subverting Toll-Like Receptor Signaling by Bacterial Pathogens. <i>Frontiers in Immunology</i> , 2015 , 6, 607	8.4	30
117	The cerebral cavernous malformation pathway controls cardiac development via regulation of endocardial MEKK3 signaling and KLF expression. <i>Developmental Cell</i> , 2015 , 32, 168-80	10.2	98
116	CREB phosphorylation at Ser133 regulates transcription via distinct mechanisms downstream of cAMP and MAPK signalling. <i>Biochemical Journal</i> , 2014 , 458, 469-79	3.8	88
115	The catalytic subunit of the system L1 amino acid transporter (slc7a5) facilitates nutrient signalling in mouse skeletal muscle. <i>PLoS ONE</i> , 2014 , 9, e89547	3.7	55
114	Transcriptional regulation of IL-10 and its cell-specific role in vivo. <i>Critical Reviews in Immunology</i> , 2014 , 34, 315-45	1.8	26
113	An H3K9/S10 methyl-phospho switch modulates Polycomb and Pol II binding at repressed genes during differentiation. <i>Molecular Biology of the Cell</i> , 2014 , 25, 904-15	3.5	29
112	Mitogen-activated protein kinases in innate immunity. <i>Nature Reviews Immunology</i> , 2013 , 13, 679-92	36.5	993
111	Cooperative control of holliday junction resolution and DNA repair by the SLX1 and MUS81-EME1 nucleases. <i>Molecular Cell</i> , 2013 , 52, 221-33	17.6	115
110	MSK1 and MSK2 inhibit lipopolysaccharide-induced prostaglandin production via an interleukin-10 feedback loop. <i>Molecular and Cellular Biology</i> , 2013 , 33, 1456-67	4.8	33
109	Cross talk between the Akt and p38 β pathways in macrophages downstream of Toll-like receptor signaling. <i>Molecular and Cellular Biology</i> , 2013 , 33, 4152-65	4.8	54
108	Mitogen- and stress-activated protein kinase 1 modulates photic entrainment of the suprachiasmatic circadian clock. <i>European Journal of Neuroscience</i> , 2013 , 37, 130-40	3.5	15

107	IFN α autocrine feedback is required to sustain TLR induced production of MCP-1 in macrophages. <i>FEBS Letters</i> , 2013 , 587, 1496-503	3.8	17
106	X-ray crystal structure of ERK5 (MAPK7) in complex with a specific inhibitor. <i>Journal of Medicinal Chemistry</i> , 2013 , 56, 4413-21	8.3	27
105	RAS-MAPK-MSK1 pathway modulates ataxin 1 protein levels and toxicity in SCA1. <i>Nature</i> , 2013 , 498, 325-331	50.4	101
104	Longitudinal assessment of endothelial function in the microvasculature of mice in-vivo. <i>Microvascular Research</i> , 2013 , 85, 86-92	3.7	9
103	PDK1 regulates VDJ recombination, cell-cycle exit and survival during B-cell development. <i>EMBO Journal</i> , 2013 , 32, 1008-22	13	27
102	PGE(2) induces macrophage IL-10 production and a regulatory-like phenotype via a protein kinase A-SIK-CRTC3 pathway. <i>Journal of Immunology</i> , 2013 , 190, 565-77	5.3	155
101	Two phases of inflammatory mediator production defined by the study of IRAK2 and IRAK1 knock-in mice. <i>Journal of Immunology</i> , 2013 , 191, 2717-30	5.3	64
100	Activation of the canonical IKK complex by K63/M1-linked hybrid ubiquitin chains. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013 , 110, 15247-52	11.5	292
99	miR-132/212 knockout mice reveal roles for these miRNAs in regulating cortical synaptic transmission and plasticity. <i>PLoS ONE</i> , 2013 , 8, e62509	3.7	103
98	Dectin-1 regulates IL-10 production via a MSK1/2 and CREB dependent pathway and promotes the induction of regulatory macrophage markers. <i>PLoS ONE</i> , 2013 , 8, e60086	3.7	61
97	Synthesis and structure-activity relationships of a novel series of pyrimidines as potent inhibitors of TBK1/IKK λ kinases. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2012 , 22, 7169-73	2.9	33
96	p38 α and p38 β kinases regulate the Toll-like receptor 4 (TLR4)-induced cytokine production by controlling ERK1/2 protein kinase pathway activation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012 , 109, 11200-5	11.5	79
95	Met acts through Abl to regulate p53 transcriptional outcomes and cell survival in the developing liver. <i>Journal of Hepatology</i> , 2012 , 57, 1292-8	13.4	14
94	Inhibition of JAKs in macrophages increases lipopolysaccharide-induced cytokine production by blocking IL-10-mediated feedback. <i>Journal of Immunology</i> , 2012 , 189, 2784-92	5.3	92
93	MSK1 regulates homeostatic and experience-dependent synaptic plasticity. <i>Journal of Neuroscience</i> , 2012 , 32, 13039-51	6.6	56
92	Mitogen- and stress-activated kinases regulate progenitor cell proliferation and neuron development in the adult dentate gyrus. <i>Journal of Neurochemistry</i> , 2012 , 123, 676-88	6	16
91	Selective kinase inhibitors as tools for neuroscience research. <i>Neuropharmacology</i> , 2012 , 63, 1227-37	5.5	13
90	MSK1 regulates environmental enrichment-induced hippocampal plasticity and cognitive enhancement. <i>Learning and Memory</i> , 2012 , 19, 550-60	2.8	23

89	Characterization of the cellular action of the MSK inhibitor SB-747651A. <i>Biochemical Journal</i> , 2012 , 441, 347-57	3.8	41
88	Pellino1 is required for interferon production by viral double-stranded RNA. <i>Journal of Biological Chemistry</i> , 2012 , 287, 34825-35	5.4	25
87	Phosphorylation of CRTC3 by the salt-inducible kinases controls the interconversion of classically activated and regulatory macrophages. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012 , 109, 16986-91	11.5	153
86	Phosphorylation of FOXO3a on Ser-7 by p38 promotes its nuclear localization in response to doxorubicin. <i>Journal of Biological Chemistry</i> , 2012 , 287, 1545-55	5.4	97
85	Identification of the protein kinases that activate the E3 ubiquitin ligase Pellino 1 in the innate immune system. <i>Biochemical Journal</i> , 2012 , 441, 339-46	3.8	44
84	The IkappaB kinase family phosphorylates the Parkinson's disease kinase LRRK2 at Ser935 and Ser910 during Toll-like receptor signaling. <i>PLoS ONE</i> , 2012 , 7, e39132	3.7	154
83	Comparison of the specificity of Trk inhibitors in recombinant and neuronal assays. <i>Neuropharmacology</i> , 2011 , 61, 148-55	5.5	15
82	Phosphorylation of cAMP-specific PDE4A5 (phosphodiesterase-4A5) by MK2 (MAPKAPK2) attenuates its activation through protein kinase A phosphorylation. <i>Biochemical Journal</i> , 2011 , 435, 755-69	3.8	48
81	The role of mitogen- and stress-activated protein kinase 1 and 2 in chronic skin inflammation in mice. <i>Experimental Dermatology</i> , 2011 , 20, 140-5	4	18
80	Mice lacking MSK1 and MSK2 show reduced skin tumor development in a two-stage chemical carcinogenesis model. <i>Cancer Investigation</i> , 2011 , 29, 240-5	2.1	25
79	Polyubiquitin binding to optineurin is required for optimal activation of TANK-binding kinase 1 and production of interferon β . <i>Journal of Biological Chemistry</i> , 2011 , 286, 35663-35674	5.4	125
78	Polyubiquitin binding to ABIN1 is required to prevent autoimmunity. <i>Journal of Experimental Medicine</i> , 2011 , 208, 1215-28	16.6	113
77	Genetic analysis of specific and redundant roles for p38alpha and p38beta MAPKs during mouse development. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 12764-9	11.5	40
76	Stress induced gene expression: a direct role for MAPKAP kinases in transcriptional activation of immediate early genes. <i>Nucleic Acids Research</i> , 2011 , 39, 2503-18	20.1	47
75	p38 α and p38 β mitogen-activated protein kinases determine cholinergic transdifferentiation of sympathetic neurons. <i>Journal of Neuroscience</i> , 2011 , 31, 12059-67	6.6	20
74	Role of p38 mitogen-activated protein kinase isoforms in murine skin inflammation induced by 12-O-tetradecanoylphorbol 13-acetate. <i>Acta Dermato-Venereologica</i> , 2011 , 91, 271-8	2.2	10
73	Evidence of p38 α and p38 β involvement in cell transformation processes. <i>Carcinogenesis</i> , 2011 , 32, 1093-9	4.6	26
72	GPR55 ligands promote receptor coupling to multiple signalling pathways. <i>British Journal of Pharmacology</i> , 2010 , 160, 604-14	8.6	156

71	The p38 MAPK pathway is essential for skeletogenesis and bone homeostasis in mice. <i>Journal of Clinical Investigation</i> , 2010 , 120, 2457-73	15.9	295
70	Regulation of miRNA transcription in macrophages in response to <i>Candida albicans</i> . <i>PLoS ONE</i> , 2010 , 5, e13669	3.7	93
69	MSK1 regulates the transcription of IL-1ra in response to TLR activation in macrophages. <i>Biochemical Journal</i> , 2010 , 425, 595-602	3.8	35
68	p38gamma regulates interaction of nuclear PSF and RNA with the tumour-suppressor hDlg in response to osmotic shock. <i>Journal of Cell Science</i> , 2010 , 123, 2596-604	5.3	14
67	Regulation of the miR-212/132 locus by MSK1 and CREB in response to neurotrophins. <i>Biochemical Journal</i> , 2010 , 428, 281-91	3.8	173
66	ERK5 pathway regulates the phosphorylation of tumour suppressor hDlg during mitosis. <i>Biochemical and Biophysical Research Communications</i> , 2010 , 399, 84-90	3.4	11
65	The activation of p38 alpha, and not p38 beta, mitogen-activated protein kinase is required for ischemic preconditioning. <i>Journal of Molecular and Cellular Cardiology</i> , 2010 , 48, 1324-8	5.8	27
64	Indirect inhibition of Toll-like receptor and type I interferon responses by ITAM-coupled receptors and integrins. <i>Immunity</i> , 2010 , 32, 518-30	32.3	110
63	RIG-I-mediated activation of p38 MAPK is essential for viral induction of interferon and activation of dendritic cells: dependence on TRAF2 and TAK1. <i>Journal of Biological Chemistry</i> , 2009 , 284, 10774-82	5.4	83
62	p38delta Mitogen-activated protein kinase is essential for skin tumor development in mice. <i>Cancer Research</i> , 2009 , 69, 4648-55	10.1	66
61	Generation of a conditional CREB Ser133Ala knockin mouse. <i>Genesis</i> , 2009 , 47, 688-96	1.9	20
60	The kinase p38 alpha serves cell type-specific inflammatory functions in skin injury and coordinates pro- and anti-inflammatory gene expression. <i>Nature Immunology</i> , 2008 , 9, 1019-27	19.1	214
59	The kinases MSK1 and MSK2 act as negative regulators of Toll-like receptor signaling. <i>Nature Immunology</i> , 2008 , 9, 1028-36	19.1	248
58	The forced swimming-induced behavioural immobility response involves histone H3 phospho-acetylation and c-Fos induction in dentate gyrus granule neurons via activation of the N-methyl-D-aspartate/extracellular signal-regulated kinase/mitogen- and stress-activated kinase signalling pathway. <i>European Journal of Neuroscience</i> , 2008 , 27, 2701-13	3.5	160
57	Roles for TAB1 in regulating the IL-1-dependent phosphorylation of the TAB3 regulatory subunit and activity of the TAK1 complex. <i>Biochemical Journal</i> , 2008 , 409, 711-22	3.8	55
56	MSK activation and physiological roles. <i>Frontiers in Bioscience - Landmark</i> , 2008 , 13, 5866-79	2.8	126
55	ERK5 regulation in naïve T-cell activation and survival. <i>European Journal of Immunology</i> , 2008 , 38, 2534-47	16.1	19
54	MSK regulate TCR-induced CREB phosphorylation but not immediate early gene transcription. <i>European Journal of Immunology</i> , 2007 , 37, 2583-95	6.1	23

53	The MAPK-activated kinase Rsk controls an acute Toll-like receptor signaling response in dendritic cells and is activated through two distinct pathways. <i>Nature Immunology</i> , 2007 , 8, 1227-35	19.1	103
52	The selectivity of protein kinase inhibitors: a further update. <i>Biochemical Journal</i> , 2007 , 408, 297-315	3.8	2080
51	The nuclear kinase mitogen- and stress-activated protein kinase 1 regulates hippocampal chromatin remodeling in memory formation. <i>Journal of Neuroscience</i> , 2007 , 27, 12732-42	6.6	195
50	Identification of novel phosphorylation sites in MSK1 by precursor ion scanning MS. <i>Biochemical Journal</i> , 2007 , 402, 491-501	3.8	44
49	MAPK activation by radio waves. <i>Biochemical Journal</i> , 2007 , 405, e5-6	3.8	5
48	Glutamate induces histone H3 phosphorylation but not acetylation in striatal neurons: role of mitogen- and stress-activated kinase-1. <i>Journal of Neurochemistry</i> , 2007 , 101, 697-708	6	52
47	CXCL12 and C5a trigger cell migration via a PAK1/2-p38alpha MAPK-MAPKAP-K2-HSP27 pathway. <i>Cellular Signalling</i> , 2006 , 18, 1897-905	4.9	107
46	Posttranslational regulation of tristetraprolin subcellular localization and protein stability by p38 mitogen-activated protein kinase and extracellular signal-regulated kinase pathways. <i>Molecular and Cellular Biology</i> , 2006 , 26, 2408-18	4.8	218
45	Evaluation of approaches to generation of tissue-specific knock-in mice. <i>Journal of Biological Chemistry</i> , 2006 , 281, 28772-81	5.4	29
44	Activation of the mitogen- and stress-activated kinase 1 by arsenic trioxide. <i>Journal of Biological Chemistry</i> , 2006 , 281, 22446-52	5.4	48
43	Post-translational control of Nur77. <i>Biochemical Society Transactions</i> , 2006 , 34, 1107-9	5.1	28
42	Nur77 is phosphorylated in cells by RSK in response to mitogenic stimulation. <i>Biochemical Journal</i> , 2006 , 393, 715-24	3.8	74
41	Pim kinases phosphorylate multiple sites on Bad and promote 14-3-3 binding and dissociation from Bcl-XL. <i>BMC Cell Biology</i> , 2006 , 7, 1		147
40	m-Calpain is required for preimplantation embryonic development in mice. <i>BMC Developmental Biology</i> , 2006 , 6, 3	3.1	109
39	C-terminal phosphorylation controls the stability and function of p27kip1. <i>EMBO Journal</i> , 2006 , 25, 5159-70	15.0	64
38	Signaling downstream of p38 in psoriasis. <i>Journal of Investigative Dermatology</i> , 2006 , 126, 1689-91	4.3	17
37	Signaling pathways and genes that inhibit pathogen-induced macrophage apoptosis--CREB and NF-kappaB as key regulators. <i>Immunity</i> , 2005 , 23, 319-29	32.3	253
36	The phosphorylation of CapZ-interacting protein (CapZIP) by stress-activated protein kinases triggers its dissociation from CapZ. <i>Biochemical Journal</i> , 2005 , 389, 127-35	3.8	50

35	MSK1 activity is controlled by multiple phosphorylation sites. <i>Biochemical Journal</i> , 2005 , 387, 507-17	3.8	132
34	MSKs are required for the transcription of the nuclear orphan receptors Nur77, Nurr1 and Nor1 downstream of MAPK signalling. <i>Biochemical Journal</i> , 2005 , 390, 749-59	3.8	98
33	p38gamma regulates the localisation of SAP97 in the cytoskeleton by modulating its interaction with GKAP. <i>EMBO Journal</i> , 2005 , 24, 1134-45	13	194
32	MAP kinase-mediated phosphorylation of distinct pools of histone H3 at S10 or S28 via mitogen- and stress-activated kinase 1/2. <i>Journal of Cell Science</i> , 2005 , 118, 2247-59	5.3	93
31	Generation and characterization of p38beta (MAPK11) gene-targeted mice. <i>Molecular and Cellular Biology</i> , 2005 , 25, 10454-64	4.8	204
30	Parsing molecular and behavioral effects of cocaine in mitogen- and stress-activated protein kinase-1-deficient mice. <i>Journal of Neuroscience</i> , 2005 , 25, 11444-54	6.6	242
29	Insulin-stimulated glucose uptake does not require p38 mitogen-activated protein kinase in adipose tissue or skeletal muscle. <i>Diabetes</i> , 2005 , 54, 3161-8	0.9	20
28	Mitogen- and stress-activated protein kinase 1 mediates cAMP response element-binding protein phosphorylation and activation by neurotrophins. <i>Journal of Neuroscience</i> , 2004 , 24, 4324-32	6.6	173
27	The in vivo role of PtdIns(3,4,5)P3 binding to PDK1 PH domain defined by knockin mutation. <i>EMBO Journal</i> , 2004 , 23, 2071-82	13	120
26	A novel UBA and UBX domain protein that binds polyubiquitin and VCP and is a substrate for SAPKs. <i>Biochemical Journal</i> , 2004 , 384, 391-400	3.8	55
25	Mitogen and stress response kinase-1 (MSK1) mediates excitotoxic induced death of hippocampal neurones. <i>Journal of Neurochemistry</i> , 2003 , 86, 25-32	6	32
24	MSK2 and MSK1 mediate the mitogen- and stress-induced phosphorylation of histone H3 and HMG-14. <i>EMBO Journal</i> , 2003 , 22, 2788-97	13	402
23	The kinase MSK1 is required for induction of c-fos by lysophosphatidic acid in mouse embryonic stem cells. <i>BMC Molecular Biology</i> , 2003 , 4, 6	4.5	44
22	Knockout of ERK5 causes multiple defects in placental and embryonic development. <i>BMC Developmental Biology</i> , 2003 , 3, 11	3.1	101
21	A second binding site revealed by C-terminal truncation of calpain small subunit, a penta-EF-hand protein. <i>Proteins: Structure, Function and Bioinformatics</i> , 2003 , 53, 649-55	4.2	6
20	In vivo role of the PIF-binding docking site of PDK1 defined by knock-in mutation. <i>EMBO Journal</i> , 2003 , 22, 4202-11	13	149
19	MSK1 and MSK2 are required for the mitogen- and stress-induced phosphorylation of CREB and ATF1 in fibroblasts. <i>Molecular and Cellular Biology</i> , 2002 , 22, 2871-81	4.8	377
18	Phosphorylation of the protein kinase mutated in Peutz-Jeghers cancer syndrome, LKB1/STK11, at Ser431 by p90(RSK) and cAMP-dependent protein kinase, but not its farnesylation at Cys(433), is essential for LKB1 to suppress cell growth. <i>Journal of Biological Chemistry</i> , 2001 , 276, 19469-82	5.4	202

17	Reduced cell migration and disruption of the actin cytoskeleton in calpain-deficient embryonic fibroblasts. <i>Journal of Biological Chemistry</i> , 2001 , 276, 48382-8	5.4	197
16	Roles of individual EF-hands in the activation of m-calpain by calcium. <i>Biochemical Journal</i> , 2000 , 348, 37	3.8	17
15	The role of 3-phosphoinositide-dependent protein kinase 1 in activating AGC kinases defined in embryonic stem cells. <i>Current Biology</i> , 2000 , 10, 439-48	6.3	404
14	Calpain zymography with casein or fluorescein isothiocyanate casein. <i>Methods in Molecular Biology</i> , 2000 , 144, 109-16	1.4	40
13	Fluorescence measurements of Ca ²⁺ binding to domain VI of calpain. <i>Methods in Molecular Biology</i> , 2000 , 144, 121-7	1.4	1
12	Disruption of the murine calpain small subunit gene, <i>Capn4</i> : calpain is essential for embryonic development but not for cell growth and division. <i>Molecular and Cellular Biology</i> , 2000 , 20, 4474-81	4.8	286
11	MSK1 is required for CREB phosphorylation in response to mitogens in mouse embryonic stem cells. <i>FEBS Letters</i> , 2000 , 482, 44-8	3.8	157
10	Crystallization and X-ray crystallographic analysis of m-calpain, a Ca ²⁺ -dependent protease. <i>Acta Crystallographica Section D: Biological Crystallography</i> , 1999 , 55, 1484-6		11
9	Structure of the mouse calpain small subunit gene. <i>BBA - Proteins and Proteomics</i> , 1998 , 1388, 247-52		13
8	m-Calpain subunits remain associated in the presence of calcium. <i>FEBS Letters</i> , 1998 , 436, 367-71	3.8	18
7	The effects of truncations of the small subunit on m-calpain activity and heterodimer formation. <i>Biochemical Journal</i> , 1997 , 326 (Pt 1), 31-8	3.8	49
6	Autolysis, Ca ²⁺ requirement, and heterodimer stability in m-calpain. <i>Journal of Biological Chemistry</i> , 1997 , 272, 11268-75	5.4	69
5	Structure of a calpain Ca(2+)-binding domain reveals a novel EF-hand and Ca(2+)-induced conformational changes. <i>Nature Structural Biology</i> , 1997 , 4, 532-8		162
4	Interaction of aspartic acid-104 and proline-287 with the active site of m-calpain. <i>Biochemical Journal</i> , 1996 , 319 (Pt 2), 535-41	3.8	9
3	Investigation of the interaction of m-calpain with phospholipids: calpain-phospholipid interactions. <i>BBA - Proteins and Proteomics</i> , 1996 , 1293, 201-6		50
2	Ca(2+)-binding domain VI of rat calpain is a homodimer in solution: hydrodynamic, crystallization and preliminary X-ray diffraction studies. <i>Protein Science</i> , 1996 , 5, 535-7	6.3	25
1	Active site residues in m-calpain: identification by site-directed mutagenesis. <i>FEBS Letters</i> , 1995 , 368, 397-400	3.8	43