

Toshiya Inada

List of Publications by Citations

Source: <https://exaly.com/author-pdf/6298874/toshiya-inada-publications-by-citations.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

203
papers

4,779
citations

34
h-index

58
g-index

211
ext. papers

5,339
ext. citations

4.1
avg, IF

4.87
L-index

#	Paper	IF	Citations
203	Psychotropic dose equivalence in Japan. <i>Psychiatry and Clinical Neurosciences</i> , 2015 , 69, 440-7	6.2	251
202	Brain neuronal CB2 cannabinoid receptors in drug abuse and depression: from mice to human subjects. <i>PLoS ONE</i> , 2008 , 3, e1640	3.7	194
201	Genome-wide association study of schizophrenia in a Japanese population. <i>Biological Psychiatry</i> , 2011 , 69, 472-8	7.9	145
200	Brain cannabinoid CB2 receptor in schizophrenia. <i>Biological Psychiatry</i> , 2010 , 67, 974-82	7.9	139
199	Functional expression of brain neuronal CB2 cannabinoid receptors are involved in the effects of drugs of abuse and in depression. <i>Annals of the New York Academy of Sciences</i> , 2008 , 1139, 434-49	6.5	139
198	Genome-wide association for methamphetamine dependence: convergent results from 2 samples. <i>Archives of General Psychiatry</i> , 2008 , 65, 345-55		118
197	Comparative Analyses of Copy-Number Variation in Autism Spectrum Disorder and Schizophrenia Reveal Etiological Overlap and Biological Insights. <i>Cell Reports</i> , 2018 , 24, 2838-2856	10.6	110
196	Meta-analysis of association between genetic variants in COMT and schizophrenia: an update. <i>Schizophrenia Research</i> , 2009 , 110, 140-8	3.6	103
195	Involvement of SMARCA2/BRM in the SWI/SNF chromatin-remodeling complex in schizophrenia. <i>Human Molecular Genetics</i> , 2009 , 18, 2483-94	5.6	89
194	Identification of YWHAЕ, a gene encoding 14-3-3epsilon, as a possible susceptibility gene for schizophrenia. <i>Human Molecular Genetics</i> , 2008 , 17, 3212-22	5.6	88
193	SIRT1 gene is associated with major depressive disorder in the Japanese population. <i>Journal of Affective Disorders</i> , 2010 , 126, 167-73	6.6	87
192	Mitochondrial DNA 3644T-->C mutation associated with bipolar disorder. <i>Genomics</i> , 2004 , 84, 1041-50	4.3	86
191	Genomewide high-density SNP linkage analysis of 236 Japanese families supports the existence of schizophrenia susceptibility loci on chromosomes 1p, 14q, and 20p. <i>American Journal of Human Genetics</i> , 2005 , 77, 937-44	11	83
190	Pathway-based association analysis of genome-wide screening data suggest that genes associated with the gamma-aminobutyric acid receptor signaling pathway are involved in neuroleptic-induced, treatment-resistant tardive dyskinesia. <i>Pharmacogenetics and Genomics</i> , 2008 , 18, 317-23	1.9	82
189	Association study of clock gene (CLOCK) and schizophrenia and mood disorders in the Japanese population. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2009 , 259, 293-7	5.1	71
188	Preliminary genome-wide association study of bipolar disorder in the Japanese population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2009 , 150B, 1110-7	3.5	61
187	Resequencing and association analysis of the KALRN and EPHB1 genes and their contribution to schizophrenia susceptibility. <i>Schizophrenia Bulletin</i> , 2012 , 38, 552-60	1.3	59

186	Relationship between catechol-O-methyltransferase polymorphism and treatment-resistant schizophrenia. <i>American Journal of Medical Genetics Part A</i> , 2003 , 120B, 35-9		59
185	Extrapyramidal symptom profiles in Japanese patients with schizophrenia treated with olanzapine or haloperidol. <i>Schizophrenia Research</i> , 2002 , 57, 227-38	3.6	56
184	Serotonin 1A receptor gene and major depressive disorder: an association study and meta-analysis. <i>Journal of Human Genetics</i> , 2009 , 54, 629-33	4.3	53
183	The dysbindin gene (DTNBP1) is associated with methamphetamine psychosis. <i>Biological Psychiatry</i> , 2008 , 63, 191-6	7.9	53
182	A nonsynonymous polymorphism in the human fatty acid amide hydrolase gene did not associate with either methamphetamine dependence or schizophrenia. <i>Neuroscience Letters</i> , 2005 , 376, 182-7	3.3	50
181	Association of the HSPG2 gene with neuroleptic-induced tardive dyskinesia. <i>Neuropsychopharmacology</i> , 2010 , 35, 1155-64	8.7	49
180	A polymorphism of the metabotropic glutamate receptor mGluR7 (GRM7) gene is associated with schizophrenia. <i>Schizophrenia Research</i> , 2008 , 101, 9-16	3.6	49
179	Association study between brain-derived neurotrophic factor gene polymorphisms and methamphetamine abusers in Japan. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2005 , 132B, 70-3	3.5	48
178	Olanzapine versus haloperidol in the treatment of patients with chronic schizophrenia: results of the Japan multicenter, double-blind olanzapine trial. <i>Psychiatry and Clinical Neurosciences</i> , 2001 , 55, 403-14	6.2	48
177	BDNF is not associated with schizophrenia: data from a Japanese population study and meta-analysis. <i>Schizophrenia Research</i> , 2009 , 112, 72-9	3.6	45
176	Extrapyramidal symptom profiles assessed with the Drug-Induced Extrapyramidal Symptom Scale: comparison with Western scales in the clinical double-blind studies of schizophrenic patients treated with either olanzapine or haloperidol. <i>International Clinical Psychopharmacology</i> , 2003 , 18, 39-48	2.2	43
175	Association between gene polymorphisms of SLC22A3 and methamphetamine use disorder. <i>Alcoholism: Clinical and Experimental Research</i> , 2006 , 30, 1644-9	3.7	42
174	Serotonin 1A receptor gene, schizophrenia and bipolar disorder: an association study and meta-analysis. <i>Psychiatry Research</i> , 2011 , 185, 20-6	9.9	39
173	The dopamine D3 receptor (DRD3) gene and risk of schizophrenia: case-control studies and an updated meta-analysis. <i>Schizophrenia Research</i> , 2010 , 116, 61-7	3.6	38
172	A functional glutathione S-transferase P1 gene polymorphism is associated with methamphetamine-induced psychosis in Japanese population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2005 , 135B, 5-9	3.5	38
171	No association was found between a functional SNP in ZDHHC8 and schizophrenia in a Japanese case-control population. <i>Neuroscience Letters</i> , 2005 , 374, 21-4	3.3	34
170	Positive association of AKT1 haplotype to Japanese methamphetamine use disorder. <i>International Journal of Neuropsychopharmacology</i> , 2006 , 9, 77-81	5.8	34
169	A population-specific uncommon variant in GRIN3A associated with schizophrenia. <i>Biological Psychiatry</i> , 2013 , 73, 532-9	7.9	33

168	The CLOCK gene and mood disorders: a case-control study and meta-analysis. <i>Chronobiology International</i> , 2011 , 28, 825-33	3.6	33
167	Relationship between three serotonin receptor subtypes (HTR3A, HTR2A and HTR4) and treatment-resistant schizophrenia in the Japanese population. <i>Neuroscience Letters</i> , 2008 , 435, 95-8	3.3	32
166	Cytochrome P450 II D6 gene polymorphisms and the neuroleptic-induced extrapyramidal symptoms in Japanese schizophrenic patients. <i>Psychiatric Genetics</i> , 2003 , 13, 163-8	2.9	32
165	Genetic association analysis of serotonin 2A receptor gene (HTR2A) with bipolar disorder and major depressive disorder in the Japanese population. <i>Neuroscience Research</i> , 2009 , 64, 231-4	2.9	30
164	Failure to replicate the association between NRG1 and schizophrenia using Japanese large sample. <i>Schizophrenia Research</i> , 2008 , 101, 1-8	3.6	30
163	No association of haplotype-tagging SNPs in TRAR4 with schizophrenia in Japanese patients. <i>Schizophrenia Research</i> , 2005 , 78, 127-30	3.6	30
162	Rolipram, a selective c-AMP phosphodiesterase inhibitor suppresses oro-facial dyskinesic movements in rats. <i>Life Sciences</i> , 1995 , 56, PL443-7	6.8	30
161	Dopamine transporter gene polymorphism and psychiatric symptoms seen in schizophrenic patients at their first episode. <i>American Journal of Medical Genetics Part A</i> , 1996 , 67, 406-8		30
160	Serotonin 1A receptor gene is associated with Japanese methamphetamine-induced psychosis patients. <i>Neuropharmacology</i> , 2010 , 58, 452-6	5.5	29
159	Genetic variants of D2 but not D3 or D4 dopamine receptor gene are associated with rapid onset and poor prognosis of methamphetamine psychosis. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2009 , 33, 625-9	5.5	29
158	Association analysis of group II metabotropic glutamate receptor genes (GRM2 and GRM3) with mood disorders and fluvoxamine response in a Japanese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2009 , 33, 875-9	5.5	29
157	A promoter haplotype of the inositol monophosphatase 2 gene (IMPA2) at 18p11.2 confers a possible risk for bipolar disorder by enhancing transcription. <i>Neuropsychopharmacology</i> , 2007 , 32, 1727-37	8.7	29
156	Identification of functional polymorphisms in the promoter region of the human PICK1 gene and their association with methamphetamine psychosis. <i>American Journal of Psychiatry</i> , 2007 , 164, 1105-14	11.9	29
155	Cytochrome P450 2D6 polymorphism and character traits. <i>Psychiatric Genetics</i> , 2003 , 13, 111-3	2.9	29
154	Association between a polymorphism in the promoter region of the dopamine D2 receptor gene and schizophrenia in Japanese subjects: replication and evaluation for antipsychotic-related features. <i>International Journal of Neuropsychopharmacology</i> , 1999 , 2, 181-186	5.8	29
153	Association of ANK3 with bipolar disorder confirmed in East Asia. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2011 , 156B, 312-5	3.5	28
152	An association study between catechol-O-methyl transferase gene polymorphism and methamphetamine psychotic disorder. <i>Psychiatric Genetics</i> , 2006 , 16, 133-8	2.9	28
151	Common variants in MAGI2 gene are associated with increased risk for cognitive impairment in schizophrenic patients. <i>PLoS ONE</i> , 2012 , 7, e36836	3.7	27

150	Gene-wide association study between the methylenetetrahydrofolate reductase gene (MTHFR) and schizophrenia in the Japanese population, with an updated meta-analysis on currently available data. <i>Schizophrenia Research</i> , 2010 , 124, 216-22	3.6	26
149	Search for a Susceptibility Locus to Tardive Dyskinesia. <i>Human Psychopharmacology</i> , 1997 , 12, 35-39	2.3	26
148	Association analysis of chromosome 5 GABAA receptor cluster in Japanese schizophrenia patients. <i>Biological Psychiatry</i> , 2005 , 58, 440-5	7.9	26
147	Haplotypes in the expression quantitative trait locus of interleukin-1 β gene are associated with schizophrenia. <i>Schizophrenia Research</i> , 2012 , 140, 185-91	3.6	25
146	Association analyses between brain-expressed fatty-acid binding protein (FABP) genes and schizophrenia and bipolar disorder. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2010 , 153B, 484-493	3.5	25
145	Gene-gene interaction analysis of personality traits in a Japanese population using an electrochemical DNA array chip analysis. <i>Neuroscience Letters</i> , 2007 , 414, 209-12	3.3	24
144	Regional brain cerebral glucose metabolism and temperament: a positron emission tomography study. <i>Neuroscience Letters</i> , 2006 , 396, 33-7	3.3	24
143	Replication and cross-phenotype study based upon schizophrenia GWASs data in the Japanese population: support for association of MHC region with psychosis. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2014 , 165B, 421-7	3.5	23
142	Genetic association analysis of functional polymorphisms in neuronal nitric oxide synthase 1 gene (NOS1) and mood disorders and fluvoxamine response in major depressive disorder in the Japanese population. <i>Neuropsychobiology</i> , 2010 , 61, 57-63	4	23
141	Association analysis of GRM2 and HTR2A with methamphetamine-induced psychosis and schizophrenia in the Japanese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2010 , 34, 639-44	5.5	23
140	Association analysis of SOD2 variants with methamphetamine psychosis in Japanese and Taiwanese populations. <i>Human Genetics</i> , 2006 , 120, 243-52	6.3	23
139	Association of SNPs linked to increased expression of SLC1A1 with schizophrenia. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2012 , 159B, 30-7	3.5	22
138	Gender difference in relationship between anxiety-related personality traits and cerebral brain glucose metabolism. <i>Psychiatry Research - Neuroimaging</i> , 2009 , 173, 206-11	2.9	22
137	Genetic variant of prodynorphin gene is risk factor for methamphetamine dependence. <i>Neuroscience Letters</i> , 2006 , 400, 158-62	3.3	22
136	Current topics in neuroleptic-induced extrapyramidal symptoms in Japan. <i>Keio Journal of Medicine</i> , 1996 , 45, 95-9	1.6	22
135	The adenosine A2A receptor is associated with methamphetamine dependence/psychosis in the Japanese population. <i>Behavioral and Brain Functions</i> , 2010 , 6, 50	4.1	21
134	The glycine transporter 1 gene (GLYT1) is associated with methamphetamine-use disorder. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2008 , 147B, 54-8	3.5	21
133	Identification of Rare, Single-Nucleotide Mutations in NDE1 and Their Contributions to Schizophrenia Susceptibility. <i>Schizophrenia Bulletin</i> , 2015 , 41, 744-53	1.3	20

132	Prescription profiles for pharmacological treatment of Japanese inpatients with schizophrenia: comparison between 2007 and 2009. <i>Human Psychopharmacology</i> , 2012 , 27, 70-5	2.3	20
131	The Frizzled 3 gene is associated with methamphetamine psychosis in the Japanese population. <i>Behavioral and Brain Functions</i> , 2008 , 4, 37	4.1	20
130	Short allele of 5-HTTLPR as a risk factor for the development of psychosis in Japanese methamphetamine abusers. <i>Annals of the New York Academy of Sciences</i> , 2008 , 1139, 49-56	6.5	20
129	Current topics in tardive dyskinesia in Japan. <i>Psychiatry and Clinical Neurosciences</i> , 1995 , 49, 239-44	6.2	20
128	No association between polymorphisms of neuronal oxide synthase 1 gene (NOS1) and schizophrenia in a Japanese population. <i>NeuroMolecular Medicine</i> , 2009 , 11, 123-7	4.6	19
127	Association between neuropeptide Y gene and its receptor Y1 gene and methamphetamine dependence. <i>Psychiatry and Clinical Neurosciences</i> , 2009 , 63, 417-22	6.2	19
126	G72 gene is associated with susceptibility to methamphetamine psychosis. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2009 , 33, 1046-9	5.5	19
125	Replication study and meta-analysis of the genetic association of GRM3 gene polymorphisms with schizophrenia in a large Japanese case-control population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2008 , 147, 392-6	3.5	19
124	Association of SOX10 with schizophrenia in the Japanese population. <i>Psychiatric Genetics</i> , 2007 , 17, 227-31	3.9	19
123	Functional polymorphism of the NQO2 gene is associated with methamphetamine psychosis. <i>Addiction Biology</i> , 2005 , 10, 145-8	4.6	19
122	First episodes of behavioral symptoms in Alzheimer's disease patients at age 90 and over, and early-onset Alzheimer's disease: comparison with senile dementia of Alzheimer's type. <i>Psychiatry and Clinical Neurosciences</i> , 2005 , 59, 730-5	6.2	19
121	Genetic variants on 3q21 and in the Sp8 transcription factor gene (SP8) as susceptibility loci for psychotic disorders: a genetic association study. <i>PLoS ONE</i> , 2013 , 8, e70964	3.7	17
120	Supportive evidence for reduced expression of GNB1L in schizophrenia. <i>Schizophrenia Bulletin</i> , 2010 , 36, 756-65	1.3	17
119	A two-stage case-control association study of the dihydropyrimidinase-like 2 gene (DPYSL2) with schizophrenia in Japanese subjects. <i>Journal of Human Genetics</i> , 2010 , 55, 469-72	4.3	17
118	A functional polymorphism in estrogen receptor alpha gene is associated with Japanese methamphetamine induced psychosis. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2009 , 33, 895-8	5.5	17
117	An association study of monoamine oxidase A (MAOA) gene polymorphism in methamphetamine psychosis. <i>Neuroscience Letters</i> , 2009 , 455, 120-3	3.3	17
116	Possible association between ubiquitin-specific peptidase 46 gene and major depressive disorders in the Japanese population. <i>Journal of Affective Disorders</i> , 2011 , 133, 150-7	6.6	16
115	Reduced CYP2D6 activity is a negative risk factor for methamphetamine dependence. <i>Neuroscience Letters</i> , 2008 , 434, 88-92	3.3	16

114	Glutamate cysteine ligase modifier (GCLM) subunit gene is not associated with methamphetamine-use disorder or schizophrenia in the Japanese population. <i>Annals of the New York Academy of Sciences</i> , 2008 , 1139, 63-9	6.5	16
113	Association analysis of delta-opioid receptor gene polymorphisms in methamphetamine dependence/psychosis. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2006 , 141B, 482-6	3.5	16
112	Association between chromogranin A gene polymorphism and schizophrenia in the Japanese population. <i>Schizophrenia Research</i> , 2006 , 83, 179-83	3.6	16
111	Stable factor structure of the Edinburgh Postnatal Depression Scale during the whole peripartum period: Results from a Japanese prospective cohort study. <i>Scientific Reports</i> , 2018 , 8, 17659	4.9	16
110	Evaluation of the individual safe correction of antipsychotic agent polypharmacy in Japanese patients with chronic schizophrenia: validation of safe corrections for antipsychotic polypharmacy and the high-dose method. <i>International Journal of Neuropsychopharmacology</i> , 2014 , 18,	5.8	15
109	Identification of a rare variant in CHD8 that contributes to schizophrenia and autism spectrum disorder susceptibility. <i>Schizophrenia Research</i> , 2016 , 178, 104-106	3.6	15
108	Analysis of the VAV3 as candidate gene for schizophrenia: evidences from voxel-based morphometry and mutation screening. <i>Schizophrenia Bulletin</i> , 2013 , 39, 720-8	1.3	15
107	Criminal offenses among discharged mentally ill individuals. Determinants of the duration from discharge and absence of diagnostic specificity. <i>International Journal of Law and Psychiatry</i> , 1998 , 21, 197-207	2.6	15
106	Genetic analysis of the gene coding for DARPP-32 (PPP1R1B) in Japanese patients with schizophrenia or bipolar disorder. <i>Schizophrenia Research</i> , 2008 , 100, 334-41	3.6	15
105	Association study between polymorphisms in glutathione-related genes and methamphetamine use disorder in a Japanese population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2008 , 147B, 1040-6	3.5	15
104	Failure to confirm the association between the FEZ1 gene and schizophrenia in a Japanese population. <i>Neuroscience Letters</i> , 2007 , 417, 326-9	3.3	15
103	The 2',3'-cyclic nucleotide 3'-phosphodiesterase and oligodendrocyte lineage transcription factor 2 genes do not appear to be associated with schizophrenia in the Japanese population. <i>Schizophrenia Research</i> , 2006 , 88, 245-50	3.6	15
102	Positive association of phencyclidine-responsive genes, PDE4A and PLAT, with schizophrenia. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2011 , 156B, 850-8	3.5	14
101	PROKR2 is associated with methamphetamine dependence in the Japanese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2010 , 34, 1033-6	5.5	14
100	No association between tagging SNPs of SNARE complex genes (STX1A, VAMP2 and SNAP25) and schizophrenia in a Japanese population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2008 , 147B, 1327-31	3.5	14
99	Alpha4 and beta2 subunits of neuronal nicotinic acetylcholine receptor genes are not associated with methamphetamine-use disorder in the Japanese population. <i>Annals of the New York Academy of Sciences</i> , 2008 , 1139, 70-82	6.5	14
98	Study protocol: safety correction of high dose antipsychotic polypharmacy in Japan. <i>BMC Psychiatry</i> , 2014 , 14, 103	4.2	13
97	Associations between the orexin (hypocretin) receptor 2 gene polymorphism Val308Ile and nicotine dependence in genome-wide and subsequent association studies. <i>Molecular Brain</i> , 2015 , 8, 50	4.5	13

96	An evaluation of polymorphisms in casein kinase 1 delta and epsilon genes in major psychiatric disorders. <i>Neuroscience Letters</i> , 2012 , 529, 66-9	3.3	13
95	Association study of ubiquitin-specific peptidase 46 (USP46) with bipolar disorder and schizophrenia in a Japanese population. <i>Journal of Human Genetics</i> , 2010 , 55, 133-6	4.3	13
94	Genetic association study of KREMEN1 and DKK1 and schizophrenia in a Japanese population. <i>Schizophrenia Research</i> , 2010 , 118, 113-7	3.6	13
93	Serotonin 6 receptor gene and mood disorders: case-control study and meta-analysis. <i>Neuroscience Research</i> , 2010 , 67, 250-5	2.9	13
92	Association study between vesicle-associated membrane protein 2 gene polymorphisms and fluvoxamine response in Japanese major depressive patients. <i>Neuropsychobiology</i> , 2006 , 54, 226-30	4	13
91	Influence of HTR2A polymorphisms and parental rearing on personality traits in healthy Japanese subjects. <i>Journal of Human Genetics</i> , 2010 , 55, 838-41	4.3	12
90	No association between monoamine oxidase A promoter polymorphism and personality traits in Japanese females. <i>Neuroscience Letters</i> , 2005 , 389, 121-3	3.3	12
89	The neural correlates of mindful awareness: a possible buffering effect on anxiety-related reduction in subgenual anterior cingulate cortex activity. <i>PLoS ONE</i> , 2013 , 8, e75526	3.7	11
88	Serotonin 6 receptor gene is associated with methamphetamine-induced psychosis in a Japanese population. <i>Drug and Alcohol Dependence</i> , 2011 , 113, 1-7	4.9	11
87	Association analysis of the GDNF gene with methamphetamine use disorder in a Japanese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2011 , 35, 1268-72	5.5	11
86	Translin-associated factor X gene (TSNAX) may be associated with female major depressive disorder in the Japanese population. <i>NeuroMolecular Medicine</i> , 2010 , 12, 78-85	4.6	11
85	Replication study for associations between polymorphisms in the CLDN5 and DGCR2 genes in the 22q11 deletion syndrome region and schizophrenia. <i>Psychiatric Genetics</i> , 2008 , 18, 255-6	2.9	11
84	The X-box binding protein 1 (XBP1) gene is not associated with methamphetamine dependence. <i>Neuroscience Letters</i> , 2005 , 383, 194-8	3.3	11
83	Genetic association analysis of tagging SNPs in alpha4 and beta2 subunits of neuronal nicotinic acetylcholine receptor genes (CHRNA4 and CHRN2) with schizophrenia in the Japanese population. <i>Journal of Neural Transmission</i> , 2008 , 115, 1457-61	4.3	10
82	Efficacy of diazepam as an anti-anxiety agent: meta-analysis of double-blind, randomized controlled trials carried out in Japan. <i>Human Psychopharmacology</i> , 2003 , 18, 483-7	2.3	10
81	Association between catechol-O-methyltransferase (COMT) polymorphism and severe alcoholic withdrawal symptoms in male Japanese alcoholics. <i>Addiction Biology</i> , 2001 , 6, 233-238	4.6	10
80	Replication study of association between ADCYAP1 gene polymorphisms and schizophrenia. <i>Psychiatric Genetics</i> , 2010 , 20, 123-5	2.9	10
79	Novel rare variants in F-box protein 45 (FBXO45) in schizophrenia. <i>Schizophrenia Research</i> , 2014 , 157, 149-56	3.6	9

78	Replication in a Japanese population that a MIR30E gene variation is associated with schizophrenia. <i>Schizophrenia Research</i> , 2013 , 150, 596-7	3.6	9
77	Two-stage case-control association study of polymorphisms in rheumatoid arthritis susceptibility genes with schizophrenia. <i>Journal of Human Genetics</i> , 2009 , 54, 62-5	4.3	9
76	Association analysis of the adenosine A1 receptor gene polymorphisms in patients with methamphetamine dependence/psychosis. <i>Current Neuropharmacology</i> , 2011 , 9, 137-42	7.6	9
75	Genetic association analysis of NRG1 with methamphetamine-induced psychosis in a Japanese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2009 , 33, 903-5	5.5	9
74	A genetic association study of the FXFD domain containing ion transport regulator 6 (FXFD6) gene, encoding phosphohippolin, in susceptibility to schizophrenia in a Japanese population. <i>Neuroscience Letters</i> , 2008 , 438, 70-5	3.3	9
73	No association between the glutamate decarboxylase 67 gene (GAD1) and schizophrenia in the Japanese population. <i>Schizophrenia Research</i> , 2007 , 91, 22-6	3.6	9
72	An exact test for the association between the disease and alleles at highly polymorphic loci with particular interest in the haplotype analysis. <i>Biometrics</i> , 2001 , 57, 769-78	1.8	9
71	Application of eye trackers for understanding mental disorders: Cases for schizophrenia and autism spectrum disorder. <i>Neuropsychopharmacology Reports</i> , 2019 , 39, 72-77	2.2	9
70	Genetic association study between the detected risk variants based upon type II diabetes GWAS and psychotic disorders in the Japanese population. <i>Journal of Human Genetics</i> , 2014 , 59, 54-6	4.3	8
69	GTP cyclohydrolase 1 gene haplotypes as predictors of SSRI response in Japanese patients with major depressive disorder. <i>Journal of Affective Disorders</i> , 2012 , 142, 315-22	6.6	8
68	Brief PANSS to assess and monitor the overall severity of schizophrenia. <i>Psychiatry and Clinical Neurosciences</i> , 2010 , 64, 262-7	6.2	8
67	Gap junction coding genes and schizophrenia: a genetic association study. <i>Journal of Human Genetics</i> , 2007 , 52, 498-501	4.3	8
66	Mentally disordered criminal offenders: five years' data from the Tokyo district public prosecutor's office. <i>International Journal of Law and Psychiatry</i> , 1995 , 18, 221-30	2.6	8
65	An association analysis of the cardiomyopathy-associated 5 (CMYA5) gene with schizophrenia in a Japanese population. <i>Psychiatric Genetics</i> , 2013 , 23, 179-80	2.9	7
64	A case control association study and cognitive function analysis of neuropilin and tolloid-like 1 gene and schizophrenia in the Japanese population. <i>PLoS ONE</i> , 2011 , 6, e28929	3.7	7
63	Association analysis of functional polymorphism in estrogen receptor alpha gene with schizophrenia and mood disorders in the Japanese population. <i>Psychiatric Genetics</i> , 2009 , 19, 217-8	2.9	7
62	Methamphetamine psychosis in which tardive dystonia was successfully treated with clonazepam. <i>Psychiatry and Clinical Neurosciences</i> , 2007 , 61, 691-4	6.2	7
61	Association study between Apolipoprotein L and schizophrenia by exhaustive and rule-based combination analysis for identification of multilocus interactions. <i>Journal of Bioscience and Bioengineering</i> , 2007 , 103, 303-10	3.3	7

60	Assessment of pharmacological toxicity using serum anticholinergic activity in a patient with dementia. <i>Psychiatry and Clinical Neurosciences</i> , 2005 , 59, 508-10	6.2	7
59	Aberrant functional connectivity between the thalamus and visual cortex is related to attentional impairment in schizophrenia. <i>Psychiatry Research - Neuroimaging</i> , 2018 , 278, 35-41	2.9	7
58	Cannabis use correlates with aggressive behavior and long-acting injectable antipsychotic treatment in Asian patients with schizophrenia. <i>Nordic Journal of Psychiatry</i> , 2019 , 73, 323-330	2.3	6
57	Serotonin 6 receptor gene and schizophrenia: case-control study and meta-analysis. <i>Human Psychopharmacology</i> , 2012 , 27, 63-9	2.3	6
56	No significant association between SIRT1 gene and methamphetamine-induced psychosis in the Japanese population. <i>Human Psychopharmacology</i> , 2011 , 26, 445-50	2.3	6
55	Association analysis of the tryptophan hydroxylase 2 gene polymorphisms in patients with methamphetamine dependence/psychosis. <i>Current Neuropharmacology</i> , 2011 , 9, 176-82	7.6	6
54	Are the cochrane group registers comprehensive? A case study of Japanese psychiatry trials. <i>BMC Medical Research Methodology</i> , 2002 , 2, 6	4.7	6
53	Protective effect of histidine on MPP ⁺ -induced hydroxyl radical generation in rat striatum. <i>Brain Research</i> , 1999 , 817, 206-8	3.7	6
52	The Risk Factors Predicting Suicidal Ideation Among Perinatal Women in Japan. <i>Frontiers in Psychiatry</i> , 2020 , 11, 441	5	5
51	Genome-wide association study of schizophrenia using microsatellite markers in the Japanese population. <i>Psychiatric Genetics</i> , 2013 , 23, 117-23	2.9	5
50	Failure to find an association between myosin heavy chain 9, non-muscle (MYH9) and schizophrenia: a three-stage case-control association study. <i>Schizophrenia Research</i> , 2010 , 118, 106-12	3.6	5
49	Large-scale case-control study of a functional polymorphism in the glutamate receptor, metabotropic 3 gene in patients with schizophrenia. <i>Psychiatry and Clinical Neurosciences</i> , 2008 , 62, 239-40	6.2	5
48	Prostate apoptosis response 4 gene is not associated with methamphetamine-use disorder in the Japanese population. <i>Annals of the New York Academy of Sciences</i> , 2008 , 1139, 83-8	6.5	5
47	. <i>Psychiatric Genetics</i> , 2003 , 13, 111-113	2.9	5
46	Support vector machine-based classification of schizophrenia patients and healthy controls using structural magnetic resonance imaging from two independent sites. <i>PLoS ONE</i> , 2020 , 15, e0239615	3.7	5
45	Drug-induced Extrapyrimalidal Symptoms Scale (DIEPSS) Serbian Language version: Inter-rater and Test-retest Reliability. <i>Scientific Reports</i> , 2017 , 7, 8105	4.9	4
44	A two-stage case-control association study between the tryptophan hydroxylase 2 (TPH2) gene and schizophrenia in a Japanese population. <i>Schizophrenia Research</i> , 2012 , 137, 264-6	3.6	4
43	Association study of bromodomain-containing 1 gene with schizophrenia in Japanese population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2010 , 153B, 786-91	3.5	4

42	Association Analysis of Nuclear Receptor Rev-erb Alpha Gene (NR1D1) and Japanese Methamphetamine Dependence. <i>Current Neuropharmacology</i> , 2011 , 9, 129-32	7.6	4
41	An association study of tachykinin receptor 3 gene with schizophrenia in the Japanese population. <i>NeuroReport</i> , 2008 , 19, 471-3	1.7	4
40	Association study between the transferrin gene and schizophrenia in the Japanese population. <i>NeuroReport</i> , 2007 , 18, 517-20	1.7	4
39	Association study of BCL9 gene polymorphism rs583583 with schizophrenia and negative symptoms in Japanese population. <i>Scientific Reports</i> , 2015 , 5, 15705	4.9	3
38	Lack of association between translin-associated factor X gene (TSNAX) and methamphetamine dependence in the Japanese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2011 , 35, 1618-22	5.5	3
37	A two-stage case-control association study of PADI2 with schizophrenia. <i>Journal of Human Genetics</i> , 2009 , 54, 430-2	4.3	3
36	Association study between the PIK4CA gene and methamphetamine use disorder in a Japanese population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2009 , 150B, 233-8	3.5	3
35	Bofu-tsusho-san effectively attenuates the weight gain observed after receiving olanzapine. <i>Psychiatry and Clinical Neurosciences</i> , 2008 , 62, 747	6.2	3
34	No association between the oligodendrocyte-related gene PLP1 and schizophrenia in the Japanese population. <i>Journal of Human Genetics</i> , 2008 , 53, 863-866	4.3	3
33	No association between the protein tyrosine phosphatase, receptor-type, Z Polypeptide 1 (PTPRZ1) gene and schizophrenia in the Japanese population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2008 , 147B, 1013-8	3.5	3
32	The life expectancy of schizophrenic patients with tardive dyskinesia. <i>Human Psychopharmacology</i> , 1992 , 7, 249-254	2.3	3
31	Genome wide study of tardive dyskinesia in schizophrenia. <i>Translational Psychiatry</i> , 2021 , 11, 351	8.6	3
30	Drug-Induced Extrapyrarnidal Symptoms Scale of the Norwegian version: inter-rater and test-retest reliability. <i>Nordic Journal of Psychiatry</i> , 2019 , 73, 546-550	2.3	2
29	Common Variants in BCL9 Gene and Schizophrenia in a Japanese Population: Association Study, Meta-Analysis and Cognitive Function Analysis / UOBIAJENE VARIJANTE BCL9 GENA I IZOFRENIJA U JAPANSKOJ POPULACIJI: IJUDIJA POVEZANOSTI, METAANALIZA I ANALIZA KOGNITIVNIH FUNKCIJA. <i>Journal of Medical Biochemistry</i> , 2015 , 32, 361-367	1.9	2
28	Lack of association between prokineticin 2 gene and Japanese methamphetamine dependence. <i>Current Neuropharmacology</i> , 2011 , 9, 133-6	7.6	2
27	Genetic Association Analysis of NOS1 and Methamphetamine-Induced Psychosis Among Japanese. <i>Current Neuropharmacology</i> , 2011 , 9, 155-9	7.6	2
26	Lack of association between MAGEL2 and schizophrenia and mood disorders in the Japanese population. <i>NeuroMolecular Medicine</i> , 2010 , 12, 285-91	4.6	2
25	A Single Medical Marker for Diagnosis of Methamphetamine Addiction - DNA Methylation of SHATI/NAT8L Promoter Sites from Patient Blood. <i>Current Pharmaceutical Design</i> , 2020 , 26, 260-264	3.3	2

24	Peripheral biomarkers of attention-deficit hyperactivity disorder: Current status and future perspective. <i>Journal of Psychiatric Research</i> , 2021 , 137, 465-470	5.2	2
23	Severe and long-lasting neuropsychiatric symptoms after mild respiratory symptoms caused by COVID-19: A case report. <i>Neuropsychopharmacology Reports</i> , 2021 ,	2.2	2
22	Case of elderly patient who developed hallucinations upon administration of oseltamivir. <i>Psychiatry and Clinical Neurosciences</i> , 2015 , 69, 305-6	6.2	1
21	An association study between the dymeclin gene and schizophrenia in the Japanese population. <i>Journal of Human Genetics</i> , 2010 , 55, 631-4	4.3	1
20	No Association Between GRM3 and Japanese Methamphetamine-Induced Psychosis. <i>Current Neuropharmacology</i> , 2011 , 9, 160-2	7.6	1
19	Visual hallucinations in CBS: reply from the author. <i>Acta Neuropsychiatrica</i> , 2004 , 16, 183-4	3.9	1
18	'Awakenings' in demented patients. <i>Psychiatry and Clinical Neurosciences</i> , 2003 , 57, 237	6.2	1
17	Factors affecting hallucinations in patients with delirium. <i>Scientific Reports</i> , 2021 , 11, 13005	4.9	1
16	Dyskinesia is most centrally situated in an estimated network of extrapyramidal syndrome in Asian patients with schizophrenia: findings from research on Asian psychotropic prescription patterns for antipsychotics. <i>Nordic Journal of Psychiatry</i> , 2021 , 75, 9-17	2.3	1
15	Significant decrease in delirium referrals after changing hypnotic from benzodiazepine to suvorexant. <i>Psychogeriatrics</i> , 2021 , 21, 324-332	1.8	1
14	Assessment of a glyoxalase I frameshift variant, p.P122fs, in Japanese patients with schizophrenia. <i>Psychiatric Genetics</i> , 2018 , 28, 90-93	2.9	1
13	Autistic traits as predictors of persistent depression. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2021 , 1	5.1	1
12	Risk factors for inducing violence in patients with delirium. <i>Brain and Behavior</i> , 2021 , 11, e2276	3.4	1
11	Lack of association of EGR2 variants with bipolar disorder in Japanese population. <i>Gene</i> , 2013 , 526, 246-508	5.8	0
10	Pacing Rhythms of Wanderers with Dementia. <i>Psychogeriatrics</i> , 2001 , 1, 76-81	1.8	0
9	A Call for a Rational Polypharmacy Policy: International Insights From Psychiatrists. <i>Psychiatry Investigation</i> , 2021 , 18, 1058-1067	3.1	0
8	Validation and factor structure of the Japanese version of the inventory to diagnose depression, lifetime version for pregnant women. <i>PLoS ONE</i> , 2020 , 15, e0234240	3.7	
7	Response to Dr. Okumura. <i>International Journal of Neuropsychopharmacology</i> , 2017 , 20, 548-549	5.8	

- 6 The inter-rater reliability of the Japanese version of the Inventory of Depressive Symptomatology, clinician version. *Human Psychopharmacology*, **2011**, 26, 267-9 2.3
- 5 Three cases of schizophrenia for which olanzapine was effective after early acute phase. *Psychiatry and Clinical Neurosciences*, **2008**, 62, 93-7 6.2
- 4 Effects of a single injection of methamphetamine on central dopaminergic systems in rats with hippocampal lesions. *Neuroscience Research Communications*, **1996**, 19, 1-7
- 3 Extraction of Potential Difficult-to-Manage Psychiatric Cases among Criminal Offenders: 5-Year Data from the Tokyo District Public Prosecutors Office. *Psychiatry and Clinical Neurosciences*, **1994**, 48, 729-735 6.2
- 2 A case of schizophrenia with twilight state due to nonconvulsive status epilepticus (NCSE). *International Clinical Psychopharmacology*, **2001**, 16, 304 2.2
- 1 Factors associated with the severity of delirium. *Human Psychopharmacology*, **2021**, 36, e2787 2.3