

# Toshiya Inada

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/6298874/toshiya-inada-publications-by-year.pdf>

**Version:** 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

203  
papers

4,779  
citations

34  
h-index

58  
g-index

211  
ext. papers

5,339  
ext. citations

4.1  
avg. IF

4.87  
L-index

#	Paper	IF	Citations
203	A Call for a Rational Polypharmacy Policy: International Insights From Psychiatrists. <i>Psychiatry Investigation</i> , <b>2021</b> , 18, 1058-1067	3.1	0
202	Peripheral biomarkers of attention-deficit hyperactivity disorder: Current status and future perspective. <i>Journal of Psychiatric Research</i> , <b>2021</b> , 137, 465-470	5.2	2
201	Factors affecting hallucinations in patients with delirium. <i>Scientific Reports</i> , <b>2021</b> , 11, 13005	4.9	1
200	Genome wide study of tardive dyskinesia in schizophrenia. <i>Translational Psychiatry</i> , <b>2021</b> , 11, 351	8.6	3
199	Dyskinesia is most centrally situated in an estimated network of extrapyramidal syndrome in Asian patients with schizophrenia: findings from research on Asian psychotropic prescription patterns for antipsychotics. <i>Nordic Journal of Psychiatry</i> , <b>2021</b> , 75, 9-17	2.3	1
198	Significant decrease in delirium referrals after changing hypnotic from benzodiazepine to suvorexant. <i>Psychogeriatrics</i> , <b>2021</b> , 21, 324-332	1.8	1
197	Factors associated with the severity of delirium. <i>Human Psychopharmacology</i> , <b>2021</b> , 36, e2787	2.3	
196	Autistic traits as predictors of persistent depression. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , <b>2021</b> , 1	5.1	1
195	Risk factors for inducing violence in patients with delirium. <i>Brain and Behavior</i> , <b>2021</b> , 11, e2276	3.4	1
194	Severe and long-lasting neuropsychiatric symptoms after mild respiratory symptoms caused by COVID-19: A case report. <i>Neuropsychopharmacology Reports</i> , <b>2021</b> ,	2.2	2
193	Validation and factor structure of the Japanese version of the inventory to diagnose depression, lifetime version for pregnant women. <i>PLoS ONE</i> , <b>2020</b> , 15, e0234240	3.7	
192	The Risk Factors Predicting Suicidal Ideation Among Perinatal Women in Japan. <i>Frontiers in Psychiatry</i> , <b>2020</b> , 11, 441	5	5
191	Support vector machine-based classification of schizophrenia patients and healthy controls using structural magnetic resonance imaging from two independent sites. <i>PLoS ONE</i> , <b>2020</b> , 15, e0239615	3.7	5
190	A Single Medical Marker for Diagnosis of Methamphetamine Addiction - DNA Methylation of SHATI/NAT8L Promoter Sites from Patient Blood. <i>Current Pharmaceutical Design</i> , <b>2020</b> , 26, 260-264	3.3	2
189	Drug-Induced Extrapyramidal Symptoms Scale of the Norwegian version: inter-rater and test-retest reliability. <i>Nordic Journal of Psychiatry</i> , <b>2019</b> , 73, 546-550	2.3	2
188	Cannabis use correlates with aggressive behavior and long-acting injectable antipsychotic treatment in Asian patients with schizophrenia. <i>Nordic Journal of Psychiatry</i> , <b>2019</b> , 73, 323-330	2.3	6
187	Application of eye trackers for understanding mental disorders: Cases for schizophrenia and autism spectrum disorder. <i>Neuropsychopharmacology Reports</i> , <b>2019</b> , 39, 72-77	2.2	9

186	Assessment of a glyoxalase I frameshift variant, p.P122fs, in Japanese patients with schizophrenia. <i>Psychiatric Genetics</i> , <b>2018</b> , 28, 90-93	2.9	1
185	Stable factor structure of the Edinburgh Postnatal Depression Scale during the whole peripartum period: Results from a Japanese prospective cohort study. <i>Scientific Reports</i> , <b>2018</b> , 8, 17659	4.9	16
184	Comparative Analyses of Copy-Number Variation in Autism Spectrum Disorder and Schizophrenia Reveal Etiological Overlap and Biological Insights. <i>Cell Reports</i> , <b>2018</b> , 24, 2838-2856	10.6	110
183	Aberrant functional connectivity between the thalamus and visual cortex is related to attentional impairment in schizophrenia. <i>Psychiatry Research - Neuroimaging</i> , <b>2018</b> , 278, 35-41	2.9	7
182	Response to Dr. Okumura. <i>International Journal of Neuropsychopharmacology</i> , <b>2017</b> , 20, 548-549	5.8	
181	Drug-induced Extrapyramidal Symptoms Scale (DIEPSS) Serbian Language version: Inter-rater and Test-retest Reliability. <i>Scientific Reports</i> , <b>2017</b> , 7, 8105	4.9	4
180	Identification of a rare variant in CHD8 that contributes to schizophrenia and autism spectrum disorder susceptibility. <i>Schizophrenia Research</i> , <b>2016</b> , 178, 104-106	3.6	15
179	Identification of Rare, Single-Nucleotide Mutations in NDE1 and Their Contributions to Schizophrenia Susceptibility. <i>Schizophrenia Bulletin</i> , <b>2015</b> , 41, 744-53	1.3	20
178	Case of elderly patient who developed hallucinations upon administration of oseltamivir. <i>Psychiatry and Clinical Neurosciences</i> , <b>2015</b> , 69, 305-6	6.2	1
177	Association study of BCL9 gene polymorphism rs583583 with schizophrenia and negative symptoms in Japanese population. <i>Scientific Reports</i> , <b>2015</b> , 5, 15705	4.9	3
176	Associations between the orexin (hypocretin) receptor 2 gene polymorphism Val308Ile and nicotine dependence in genome-wide and subsequent association studies. <i>Molecular Brain</i> , <b>2015</b> , 8, 50	4.5	13
175	Psychotropic dose equivalence in Japan. <i>Psychiatry and Clinical Neurosciences</i> , <b>2015</b> , 69, 440-7	6.2	251
174	Evaluation of the individual safe correction of antipsychotic agent polypharmacy in Japanese patients with chronic schizophrenia: validation of safe corrections for antipsychotic polypharmacy and the high-dose method. <i>International Journal of Neuropsychopharmacology</i> , <b>2014</b> , 18,	5.8	15
173	Replication and cross-phenotype study based upon schizophrenia GWASs data in the Japanese population: support for association of MHC region with psychosis. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2014</b> , 165B, 421-7	3.5	23
172	Study protocol: safety correction of high dose antipsychotic polypharmacy in Japan. <i>BMC Psychiatry</i> , <b>2014</b> , 14, 103	4.2	13
171	Novel rare variants in F-box protein 45 (FBXO45) in schizophrenia. <i>Schizophrenia Research</i> , <b>2014</b> , 157, 149-56	3.6	9
170	Genetic association study between the detected risk variants based upon type II diabetes GWAS and psychotic disorders in the Japanese population. <i>Journal of Human Genetics</i> , <b>2014</b> , 59, 54-6	4.3	8
169	Replication in a Japanese population that a MIR30E gene variation is associated with schizophrenia. <i>Schizophrenia Research</i> , <b>2013</b> , 150, 596-7	3.6	9

168	Lack of association of EGR2 variants with bipolar disorder in Japanese population. <i>Gene</i> , <b>2013</b> , 526, 246-508	0	0
167	A population-specific uncommon variant in GRIN3A associated with schizophrenia. <i>Biological Psychiatry</i> , <b>2013</b> , 73, 532-9	7.9	33
166	Analysis of the VAV3 as candidate gene for schizophrenia: evidences from voxel-based morphometry and mutation screening. <i>Schizophrenia Bulletin</i> , <b>2013</b> , 39, 720-8	1.3	15
165	An association analysis of the cardiomyopathy-associated 5 (CMYA5) gene with schizophrenia in a Japanese population. <i>Psychiatric Genetics</i> , <b>2013</b> , 23, 179-80	2.9	7
164	Genome-wide association study of schizophrenia using microsatellite markers in the Japanese population. <i>Psychiatric Genetics</i> , <b>2013</b> , 23, 117-23	2.9	5
163	Common Variants in BCL9 Gene and Schizophrenia in a Japanese Population: Association Study, Meta-Analysis and Cognitive Function Analysis / UOBIAJENE VARIJANTE BCL9 GENA I IZOFRENIJA U JAPANSKOJ POPULACIJI: IJUDIJA POVEZANOSTI, METAANALIZA I ANALIZA KOGNITIVNIH FUNKCIJA. <i>Journal of Medical Genetics</i> , <b>2013</b> , 33, 311-317	1.9	2
162	Genetic variants on 3q21 and in the Sp8 transcription factor gene (SP8) as susceptibility loci for psychotic disorders: a genetic association study. <i>PLoS ONE</i> , <b>2013</b> , 8, e70964	3.7	17
161	The neural correlates of mindful awareness: a possible buffering effect on anxiety-related reduction in subgenual anterior cingulate cortex activity. <i>PLoS ONE</i> , <b>2013</b> , 8, e75526	3.7	11
160	Serotonin 6 receptor gene and schizophrenia: case-control study and meta-analysis. <i>Human Psychopharmacology</i> , <b>2012</b> , 27, 63-9	2.3	6
159	Prescription profiles for pharmacological treatment of Japanese inpatients with schizophrenia: comparison between 2007 and 2009. <i>Human Psychopharmacology</i> , <b>2012</b> , 27, 70-5	2.3	20
158	Association of SNPs linked to increased expression of SLC1A1 with schizophrenia. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2012</b> , 159B, 30-7	3.5	22
157	An evaluation of polymorphisms in casein kinase 1 delta and epsilon genes in major psychiatric disorders. <i>Neuroscience Letters</i> , <b>2012</b> , 529, 66-9	3.3	13
156	GTP cyclohydrolase 1 gene haplotypes as predictors of SSRI response in Japanese patients with major depressive disorder. <i>Journal of Affective Disorders</i> , <b>2012</b> , 142, 315-22	6.6	8
155	A two-stage case-control association study between the tryptophan hydroxylase 2 (TPH2) gene and schizophrenia in a Japanese population. <i>Schizophrenia Research</i> , <b>2012</b> , 137, 264-6	3.6	4
154	Haplotypes in the expression quantitative trait locus of interleukin-1 $\beta$ gene are associated with schizophrenia. <i>Schizophrenia Research</i> , <b>2012</b> , 140, 185-91	3.6	25
153	Common variants in MAGI2 gene are associated with increased risk for cognitive impairment in schizophrenic patients. <i>PLoS ONE</i> , <b>2012</b> , 7, e36836	3.7	27
152	Resequencing and association analysis of the KALRN and EPHB1 genes and their contribution to schizophrenia susceptibility. <i>Schizophrenia Bulletin</i> , <b>2012</b> , 38, 552-60	1.3	59
151	Genome-wide association study of schizophrenia in a Japanese population. <i>Biological Psychiatry</i> , <b>2011</b> , 69, 472-8	7.9	145

150	The CLOCK gene and mood disorders: a case-control study and meta-analysis. <i>Chronobiology International</i> , <b>2011</b> , 28, 825-33	3.6	33
149	Serotonin 6 receptor gene is associated with methamphetamine-induced psychosis in a Japanese population. <i>Drug and Alcohol Dependence</i> , <b>2011</b> , 113, 1-7	4.9	11
148	Association analysis of the GDNF gene with methamphetamine use disorder in a Japanese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , <b>2011</b> , 35, 1268-72	5.5	11
147	Lack of association between translin-associated factor X gene (TSNAX) and methamphetamine dependence in the Japanese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , <b>2011</b> , 35, 1618-22	5.5	3
146	Serotonin 1A receptor gene, schizophrenia and bipolar disorder: an association study and meta-analysis. <i>Psychiatry Research</i> , <b>2011</b> , 185, 20-6	9.9	39
145	A case control association study and cognitive function analysis of neuropilin and tolloid-like 1 gene and schizophrenia in the Japanese population. <i>PLoS ONE</i> , <b>2011</b> , 6, e28929	3.7	7
144	Possible association between ubiquitin-specific peptidase 46 gene and major depressive disorders in the Japanese population. <i>Journal of Affective Disorders</i> , <b>2011</b> , 133, 150-7	6.6	16
143	Association of ANK3 with bipolar disorder confirmed in East Asia. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2011</b> , 156B, 312-5	3.5	28
142	Positive association of phencyclidine-responsive genes, PDE4A and PLAT, with schizophrenia. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2011</b> , 156B, 850-8	3.5	14
141	The inter-rater reliability of the Japanese version of the Inventory of Depressive Symptomatology, clinician version. <i>Human Psychopharmacology</i> , <b>2011</b> , 26, 267-9	2.3	
140	No significant association between SIRT1 gene and methamphetamine-induced psychosis in the Japanese population. <i>Human Psychopharmacology</i> , <b>2011</b> , 26, 445-50	2.3	6
139	Association Analysis of Nuclear Receptor Rev-erb Alpha Gene (NR1D1) and Japanese Methamphetamine Dependence. <i>Current Neuropharmacology</i> , <b>2011</b> , 9, 129-32	7.6	4
138	No Association Between GRM3 and Japanese Methamphetamine-Induced Psychosis. <i>Current Neuropharmacology</i> , <b>2011</b> , 9, 160-2	7.6	1
137	Association analysis of the tryptophan hydroxylase 2 gene polymorphisms in patients with methamphetamine dependence/psychosis. <i>Current Neuropharmacology</i> , <b>2011</b> , 9, 176-82	7.6	6
136	Association analysis of the adenosine A1 receptor gene polymorphisms in patients with methamphetamine dependence/psychosis. <i>Current Neuropharmacology</i> , <b>2011</b> , 9, 137-42	7.6	9
135	Lack of association between prokineticin 2 gene and Japanese methamphetamine dependence. <i>Current Neuropharmacology</i> , <b>2011</b> , 9, 133-6	7.6	2
134	Genetic Association Analysis of NOS1 and Methamphetamine-Induced Psychosis Among Japanese. <i>Current Neuropharmacology</i> , <b>2011</b> , 9, 155-9	7.6	2
133	Brief PANSS to assess and monitor the overall severity of schizophrenia. <i>Psychiatry and Clinical Neurosciences</i> , <b>2010</b> , 64, 262-7	6.2	8

132	Genetic association analysis of functional polymorphisms in neuronal nitric oxide synthase 1 gene (NOS1) and mood disorders and fluvoxamine response in major depressive disorder in the Japanese population. <i>Neuropsychobiology</i> , <b>2010</b> , 61, 57-63	4	23
131	Association of the HSPG2 gene with neuroleptic-induced tardive dyskinesia. <i>Neuropsychopharmacology</i> , <b>2010</b> , 35, 1155-64	8.7	49
130	Influence of HTR2A polymorphisms and parental rearing on personality traits in healthy Japanese subjects. <i>Journal of Human Genetics</i> , <b>2010</b> , 55, 838-41	4.3	12
129	Supportive evidence for reduced expression of GNB1L in schizophrenia. <i>Schizophrenia Bulletin</i> , <b>2010</b> , 36, 756-65	1.3	17
128	Association study of ubiquitin-specific peptidase 46 (USP46) with bipolar disorder and schizophrenia in a Japanese population. <i>Journal of Human Genetics</i> , <b>2010</b> , 55, 133-6	4.3	13
127	A two-stage case-control association study of the dihydropyrimidinase-like 2 gene (DPYSL2) with schizophrenia in Japanese subjects. <i>Journal of Human Genetics</i> , <b>2010</b> , 55, 469-72	4.3	17
126	An association study between the dymeclin gene and schizophrenia in the Japanese population. <i>Journal of Human Genetics</i> , <b>2010</b> , 55, 631-4	4.3	1
125	Failure to find an association between myosin heavy chain 9, non-muscle (MYH9) and schizophrenia: a three-stage case-control association study. <i>Schizophrenia Research</i> , <b>2010</b> , 118, 106-12	3.6	5
124	Brain cannabinoid CB2 receptor in schizophrenia. <i>Biological Psychiatry</i> , <b>2010</b> , 67, 974-82	7.9	139
123	The dopamine D3 receptor (DRD3) gene and risk of schizophrenia: case-control studies and an updated meta-analysis. <i>Schizophrenia Research</i> , <b>2010</b> , 116, 61-7	3.6	38
122	Genetic association study of KREMEN1 and DKK1 and schizophrenia in a Japanese population. <i>Schizophrenia Research</i> , <b>2010</b> , 118, 113-7	3.6	13
121	Gene-wide association study between the methylenetetrahydrofolate reductase gene (MTHFR) and schizophrenia in the Japanese population, with an updated meta-analysis on currently available data. <i>Schizophrenia Research</i> , <b>2010</b> , 124, 216-22	3.6	26
120	Serotonin 6 receptor gene and mood disorders: case-control study and meta-analysis. <i>Neuroscience Research</i> , <b>2010</b> , 67, 250-5	2.9	13
119	Association analysis of GRM2 and HTR2A with methamphetamine-induced psychosis and schizophrenia in the Japanese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , <b>2010</b> , 34, 639-44	5.5	23
118	PROKR2 is associated with methamphetamine dependence in the Japanese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , <b>2010</b> , 34, 1033-6	5.5	14
117	Serotonin 1A receptor gene is associated with Japanese methamphetamine-induced psychosis patients. <i>Neuropharmacology</i> , <b>2010</b> , 58, 452-6	5.5	29
116	Association study of bromodomain-containing 1 gene with schizophrenia in Japanese population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2010</b> , 153B, 786-91	3.5	4
115	Translin-associated factor X gene (TSNAX) may be associated with female major depressive disorder in the Japanese population. <i>NeuroMolecular Medicine</i> , <b>2010</b> , 12, 78-85	4.6	11

114	Lack of association between MAGEL2 and schizophrenia and mood disorders in the Japanese population. <i>NeuroMolecular Medicine</i> , <b>2010</b> , 12, 285-91	4.6	2
113	SIRT1 gene is associated with major depressive disorder in the Japanese population. <i>Journal of Affective Disorders</i> , <b>2010</b> , 126, 167-73	6.6	87
112	Association analyses between brain-expressed fatty-acid binding protein (FABP) genes and schizophrenia and bipolar disorder. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2010</b> , 153B, 484-493	3.5	25
111	The adenosine A2A receptor is associated with methamphetamine dependence/psychosis in the Japanese population. <i>Behavioral and Brain Functions</i> , <b>2010</b> , 6, 50	4.1	21
110	Replication study of association between ADCYAP1 gene polymorphisms and schizophrenia. <i>Psychiatric Genetics</i> , <b>2010</b> , 20, 123-5	2.9	10
109	A two-stage case-control association study of PADI2 with schizophrenia. <i>Journal of Human Genetics</i> , <b>2009</b> , 54, 430-2	4.3	3
108	Two-stage case-control association study of polymorphisms in rheumatoid arthritis susceptibility genes with schizophrenia. <i>Journal of Human Genetics</i> , <b>2009</b> , 54, 62-5	4.3	9
107	Serotonin 1A receptor gene and major depressive disorder: an association study and meta-analysis. <i>Journal of Human Genetics</i> , <b>2009</b> , 54, 629-33	4.3	53
106	Gender difference in relationship between anxiety-related personality traits and cerebral brain glucose metabolism. <i>Psychiatry Research - Neuroimaging</i> , <b>2009</b> , 173, 206-11	2.9	22
105	Association study between the PIK4CA gene and methamphetamine use disorder in a Japanese population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2009</b> , 150B, 233-8	3.5	3
104	Preliminary genome-wide association study of bipolar disorder in the Japanese population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2009</b> , 150B, 1110-7	3.5	61
103	Association study of clock gene (CLOCK) and schizophrenia and mood disorders in the Japanese population. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , <b>2009</b> , 259, 293-7	5.1	71
102	No association between polymorphisms of neuronal oxide synthase 1 gene (NOS1) and schizophrenia in a Japanese population. <i>NeuroMolecular Medicine</i> , <b>2009</b> , 11, 123-7	4.6	19
101	Association between neuropeptide Y gene and its receptor Y1 gene and methamphetamine dependence. <i>Psychiatry and Clinical Neurosciences</i> , <b>2009</b> , 63, 417-22	6.2	19
100	Genetic variants of D2 but not D3 or D4 dopamine receptor gene are associated with rapid onset and poor prognosis of methamphetamine psychosis. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , <b>2009</b> , 33, 625-9	5.5	29
99	Association analysis of group II metabotropic glutamate receptor genes (GRM2 and GRM3) with mood disorders and fluvoxamine response in a Japanese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , <b>2009</b> , 33, 875-9	5.5	29
98	A functional polymorphism in estrogen receptor alpha gene is associated with Japanese methamphetamine induced psychosis. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , <b>2009</b> , 33, 895-8	5.5	17
97	Genetic association analysis of NRG1 with methamphetamine-induced psychosis in a Japanese population. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , <b>2009</b> , 33, 903-5	5.5	9

96	G72 gene is associated with susceptibility to methamphetamine psychosis. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , <b>2009</b> , 33, 1046-9	5.5	19
95	An association study of monoamine oxidase A (MAOA) gene polymorphism in methamphetamine psychosis. <i>Neuroscience Letters</i> , <b>2009</b> , 455, 120-3	3.3	17
94	Genetic association analysis of serotonin 2A receptor gene (HTR2A) with bipolar disorder and major depressive disorder in the Japanese population. <i>Neuroscience Research</i> , <b>2009</b> , 64, 231-4	2.9	30
93	Meta-analysis of association between genetic variants in COMT and schizophrenia: an update. <i>Schizophrenia Research</i> , <b>2009</b> , 110, 140-8	3.6	103
92	BDNF is not associated with schizophrenia: data from a Japanese population study and meta-analysis. <i>Schizophrenia Research</i> , <b>2009</b> , 112, 72-9	3.6	45
91	Involvement of SMARCA2/BRM in the SWI/SNF chromatin-remodeling complex in schizophrenia. <i>Human Molecular Genetics</i> , <b>2009</b> , 18, 2483-94	5.6	89
90	Association analysis of functional polymorphism in estrogen receptor alpha gene with schizophrenia and mood disorders in the Japanese population. <i>Psychiatric Genetics</i> , <b>2009</b> , 19, 217-8	2.9	7
89	Three cases of schizophrenia for which olanzapine was effective after early acute phase. <i>Psychiatry and Clinical Neurosciences</i> , <b>2008</b> , 62, 93-7	6.2	
88	Large-scale case-control study of a functional polymorphism in the glutamate receptor, metabotropic 3 gene in patients with schizophrenia. <i>Psychiatry and Clinical Neurosciences</i> , <b>2008</b> , 62, 239-40	6.2	5
87	Bofu-tsusho-san effectively attenuates the weight gain observed after receiving olanzapine. <i>Psychiatry and Clinical Neurosciences</i> , <b>2008</b> , 62, 747	6.2	3
86	Genetic analysis of the gene coding for DARPP-32 (PPP1R1B) in Japanese patients with schizophrenia or bipolar disorder. <i>Schizophrenia Research</i> , <b>2008</b> , 100, 334-41	3.6	15
85	Failure to replicate the association between NRG1 and schizophrenia using Japanese large sample. <i>Schizophrenia Research</i> , <b>2008</b> , 101, 1-8	3.6	30
84	A polymorphism of the metabotropic glutamate receptor mGluR7 (GRM7) gene is associated with schizophrenia. <i>Schizophrenia Research</i> , <b>2008</b> , 101, 9-16	3.6	49
83	The Frizzled 3 gene is associated with methamphetamine psychosis in the Japanese population. <i>Behavioral and Brain Functions</i> , <b>2008</b> , 4, 37	4.1	20
82	The dysbindin gene (DTNBP1) is associated with methamphetamine psychosis. <i>Biological Psychiatry</i> , <b>2008</b> , 63, 191-6	7.9	53
81	Reduced CYP2D6 activity is a negative risk factor for methamphetamine dependence. <i>Neuroscience Letters</i> , <b>2008</b> , 434, 88-92	3.3	16
80	Relationship between three serotonin receptor subtypes (HTR3A, HTR2A and HTR4) and treatment-resistant schizophrenia in the Japanese population. <i>Neuroscience Letters</i> , <b>2008</b> , 435, 95-8	3.3	32
79	A genetic association study of the FXFD domain containing ion transport regulator 6 (FXFD6) gene, encoding phosphohippolin, in susceptibility to schizophrenia in a Japanese population. <i>Neuroscience Letters</i> , <b>2008</b> , 438, 70-5	3.3	9



78	Identification of YWHAE, a gene encoding 14-3-3epsilon, as a possible susceptibility gene for schizophrenia. <i>Human Molecular Genetics</i> , <b>2008</b> , 17, 3212-22	5.6	88
77	Genome-wide association for methamphetamine dependence: convergent results from 2 samples. <i>Archives of General Psychiatry</i> , <b>2008</b> , 65, 345-55		118
76	Replication study for associations between polymorphisms in the CLDN5 and DGCR2 genes in the 22q11 deletion syndrome region and schizophrenia. <i>Psychiatric Genetics</i> , <b>2008</b> , 18, 255-6	2.9	11
75	Pathway-based association analysis of genome-wide screening data suggest that genes associated with the gamma-aminobutyric acid receptor signaling pathway are involved in neuroleptic-induced, treatment-resistant tardive dyskinesia. <i>Pharmacogenetics and Genomics</i> , <b>2008</b> , 18, 317-23	1.9	82
74	An association study of tachykinin receptor 3 gene with schizophrenia in the Japanese population. <i>NeuroReport</i> , <b>2008</b> , 19, 471-3	1.7	4
73	No association between the oligodendrocyte-related gene PLP1 and schizophrenia in the Japanese population. <i>Journal of Human Genetics</i> , <b>2008</b> , 53, 863-866	4.3	3
72	Genetic association analysis of tagging SNPs in alpha4 and beta2 subunits of neuronal nicotinic acetylcholine receptor genes (CHRNA4 and CHRN2) with schizophrenia in the Japanese population. <i>Journal of Neural Transmission</i> , <b>2008</b> , 115, 1457-61	4.3	10
71	The glycine transporter 1 gene (GLYT1) is associated with methamphetamine-use disorder. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2008</b> , 147B, 54-8	3.5	21
70	Replication study and meta-analysis of the genetic association of GRM3 gene polymorphisms with schizophrenia in a large Japanese case-control population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2008</b> , 147, 392-6	3.5	19
69	No association between the protein tyrosine phosphatase, receptor-type, Z Polypeptide 1 (PTPRZ1) gene and schizophrenia in the Japanese population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2008</b> , 147B, 1013-8	3.5	3
68	Association study between polymorphisms in glutathione-related genes and methamphetamine use disorder in a Japanese population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2008</b> , 147B, 1040-6	3.5	15
67	No association between tagging SNPs of SNARE complex genes (STX1A, VAMP2 and SNAP25) and schizophrenia in a Japanese population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2008</b> , 147B, 1327-31	3.5	14
66	Short allele of 5-HTTLPR as a risk factor for the development of psychosis in Japanese methamphetamine abusers. <i>Annals of the New York Academy of Sciences</i> , <b>2008</b> , 1139, 49-56	6.5	20
65	Glutamate cysteine ligase modifier (GCLM) subunit gene is not associated with methamphetamine-use disorder or schizophrenia in the Japanese population. <i>Annals of the New York Academy of Sciences</i> , <b>2008</b> , 1139, 63-9	6.5	16
64	Alpha4 and beta2 subunits of neuronal nicotinic acetylcholine receptor genes are not associated with methamphetamine-use disorder in the Japanese population. <i>Annals of the New York Academy of Sciences</i> , <b>2008</b> , 1139, 70-82	6.5	14
63	Prostate apoptosis response 4 gene is not associated with methamphetamine-use disorder in the Japanese population. <i>Annals of the New York Academy of Sciences</i> , <b>2008</b> , 1139, 83-8	6.5	5
62	Functional expression of brain neuronal CB2 cannabinoid receptors are involved in the effects of drugs of abuse and in depression. <i>Annals of the New York Academy of Sciences</i> , <b>2008</b> , 1139, 434-49	6.5	139
61	Brain neuronal CB2 cannabinoid receptors in drug abuse and depression: from mice to human subjects. <i>PLoS ONE</i> , <b>2008</b> , 3, e1640	3.7	194

60	No association between the glutamate decarboxylase 67 gene (GAD1) and schizophrenia in the Japanese population. <i>Schizophrenia Research</i> , <b>2007</b> , 91, 22-6	3.6	9
59	Methamphetamine psychosis in which tardive dystonia was successfully treated with clonazepam. <i>Psychiatry and Clinical Neurosciences</i> , <b>2007</b> , 61, 691-4	6.2	7
58	Association study between Apolipoprotein L and schizophrenia by exhaustive and rule-based combination analysis for identification of multilocus interactions. <i>Journal of Bioscience and Bioengineering</i> , <b>2007</b> , 103, 303-10	3.3	7
57	Gap junction coding genes and schizophrenia: a genetic association study. <i>Journal of Human Genetics</i> , <b>2007</b> , 52, 498-501	4.3	8
56	A promoter haplotype of the inositol monophosphatase 2 gene (IMPA2) at 18p11.2 confers a possible risk for bipolar disorder by enhancing transcription. <i>Neuropsychopharmacology</i> , <b>2007</b> , 32, 1727-37	8.7	29
55	Identification of functional polymorphisms in the promoter region of the human PICK1 gene and their association with methamphetamine psychosis. <i>American Journal of Psychiatry</i> , <b>2007</b> , 164, 1105-14	11.9	29
54	Association of SOX10 with schizophrenia in the Japanese population. <i>Psychiatric Genetics</i> , <b>2007</b> , 17, 227-31	3.1	19
53	Association study between the transferrin gene and schizophrenia in the Japanese population. <i>NeuroReport</i> , <b>2007</b> , 18, 517-20	1.7	4
52	Gene-gene interaction analysis of personality traits in a Japanese population using an electrochemical DNA array chip analysis. <i>Neuroscience Letters</i> , <b>2007</b> , 414, 209-12	3.3	24
51	Failure to confirm the association between the FEZ1 gene and schizophrenia in a Japanese population. <i>Neuroscience Letters</i> , <b>2007</b> , 417, 326-9	3.3	15
50	Association analysis of delta-opioid receptor gene polymorphisms in methamphetamine dependence/psychosis. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2006</b> , 141B, 482-6	3.5	16
49	Association study between vesicle-associated membrane protein 2 gene polymorphisms and fluvoxamine response in Japanese major depressive patients. <i>Neuropsychobiology</i> , <b>2006</b> , 54, 226-30	4	13
48	Association between chromogranin A gene polymorphism and schizophrenia in the Japanese population. <i>Schizophrenia Research</i> , <b>2006</b> , 83, 179-83	3.6	16
47	The 2',3'-cyclic nucleotide 3'-phosphodiesterase and oligodendrocyte lineage transcription factor 2 genes do not appear to be associated with schizophrenia in the Japanese population. <i>Schizophrenia Research</i> , <b>2006</b> , 88, 245-50	3.6	15
46	Regional brain cerebral glucose metabolism and temperament: a positron emission tomography study. <i>Neuroscience Letters</i> , <b>2006</b> , 396, 33-7	3.3	24
45	Genetic variant of prodynorphin gene is risk factor for methamphetamine dependence. <i>Neuroscience Letters</i> , <b>2006</b> , 400, 158-62	3.3	22
44	Positive association of AKT1 haplotype to Japanese methamphetamine use disorder. <i>International Journal of Neuropsychopharmacology</i> , <b>2006</b> , 9, 77-81	5.8	34
43	An association study between catechol-O-methyl transferase gene polymorphism and methamphetamine psychotic disorder. <i>Psychiatric Genetics</i> , <b>2006</b> , 16, 133-8	2.9	28

42	Association between gene polymorphisms of SLC22A3 and methamphetamine use disorder. <i>Alcoholism: Clinical and Experimental Research</i> , <b>2006</b> , 30, 1644-9	3.7	42
41	Association analysis of SOD2 variants with methamphetamine psychosis in Japanese and Taiwanese populations. <i>Human Genetics</i> , <b>2006</b> , 120, 243-52	6.3	23
40	Genomewide high-density SNP linkage analysis of 236 Japanese families supports the existence of schizophrenia susceptibility loci on chromosomes 1p, 14q, and 20p. <i>American Journal of Human Genetics</i> , <b>2005</b> , 77, 937-44	11	83
39	No association was found between a functional SNP in ZDHHC8 and schizophrenia in a Japanese case-control population. <i>Neuroscience Letters</i> , <b>2005</b> , 374, 21-4	3.3	34
38	A nonsynonymous polymorphism in the human fatty acid amide hydrolase gene did not associate with either methamphetamine dependence or schizophrenia. <i>Neuroscience Letters</i> , <b>2005</b> , 376, 182-7	3.3	50
37	No association between monoamine oxidase A promoter polymorphism and personality traits in Japanese females. <i>Neuroscience Letters</i> , <b>2005</b> , 389, 121-3	3.3	12
36	The X-box binding protein 1 (XBP1) gene is not associated with methamphetamine dependence. <i>Neuroscience Letters</i> , <b>2005</b> , 383, 194-8	3.3	11
35	Association analysis of chromosome 5 GABAA receptor cluster in Japanese schizophrenia patients. <i>Biological Psychiatry</i> , <b>2005</b> , 58, 440-5	7.9	26
34	No association of haplotype-tagging SNPs in TRAR4 with schizophrenia in Japanese patients. <i>Schizophrenia Research</i> , <b>2005</b> , 78, 127-30	3.6	30
33	Functional polymorphism of the NQO2 gene is associated with methamphetamine psychosis. <i>Addiction Biology</i> , <b>2005</b> , 10, 145-8	4.6	19
32	Assessment of pharmacological toxicity using serum anticholinergic activity in a patient with dementia. <i>Psychiatry and Clinical Neurosciences</i> , <b>2005</b> , 59, 508-10	6.2	7
31	First episodes of behavioral symptoms in Alzheimer's disease patients at age 90 and over, and early-onset Alzheimer's disease: comparison with senile dementia of Alzheimer's type. <i>Psychiatry and Clinical Neurosciences</i> , <b>2005</b> , 59, 730-5	6.2	19
30	Association study between brain-derived neurotrophic factor gene polymorphisms and methamphetamine abusers in Japan. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2005</b> , 132B, 70-3	3.5	48
29	A functional glutathione S-transferase P1 gene polymorphism is associated with methamphetamine-induced psychosis in Japanese population. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , <b>2005</b> , 135B, 5-9	3.5	38
28	Visual hallucinations in CBS: reply from the author. <i>Acta Neuropsychiatrica</i> , <b>2004</b> , 16, 183-4	3.9	1
27	Mitochondrial DNA 3644T-->C mutation associated with bipolar disorder. <i>Genomics</i> , <b>2004</b> , 84, 1041-50	4.3	86
26	Extrapyramidal symptom profiles assessed with the Drug-Induced Extrapyramidal Symptom Scale: comparison with Western scales in the clinical double-blind studies of schizophrenic patients treated with either olanzapine or haloperidol. <i>International Clinical Psychopharmacology</i> , <b>2003</b> , 18, 39-48	2.2	43
25	Cytochrome P450 II D6 gene polymorphisms and the neuroleptic-induced extrapyramidal symptoms in Japanese schizophrenic patients. <i>Psychiatric Genetics</i> , <b>2003</b> , 13, 163-8	2.9	32

24	. <i>Psychiatric Genetics</i> , <b>2003</b> , 13, 111-113	2.9	5
23	Cytochrome P450 2D6 polymorphism and character traits. <i>Psychiatric Genetics</i> , <b>2003</b> , 13, 111-3	2.9	29
22	Efficacy of diazepam as an anti-anxiety agent: meta-analysis of double-blind, randomized controlled trials carried out in Japan. <i>Human Psychopharmacology</i> , <b>2003</b> , 18, 483-7	2.3	10
21	Relationship between catechol-O-methyltransferase polymorphism and treatment-resistant schizophrenia. <i>American Journal of Medical Genetics Part A</i> , <b>2003</b> , 120B, 35-9		59
20	'Awakenings' in demented patients. <i>Psychiatry and Clinical Neurosciences</i> , <b>2003</b> , 57, 237	6.2	1
19	Are the cochrane group registers comprehensive? A case study of Japanese psychiatry trials. <i>BMC Medical Research Methodology</i> , <b>2002</b> , 2, 6	4.7	6
18	Extrapyramidal symptom profiles in Japanese patients with schizophrenia treated with olanzapine or haloperidol. <i>Schizophrenia Research</i> , <b>2002</b> , 57, 227-38	3.6	56
17	Olanzapine versus haloperidol in the treatment of patients with chronic schizophrenia: results of the Japan multicenter, double-blind olanzapine trial. <i>Psychiatry and Clinical Neurosciences</i> , <b>2001</b> , 55, 403-14	6.2	48
16	Association between catechol-O-methyltransferase (COMT) polymorphism and severe alcoholic withdrawal symptoms in male Japanese alcoholics. <i>Addiction Biology</i> , <b>2001</b> , 6, 233-238	4.6	10
15	An exact test for the association between the disease and alleles at highly polymorphic loci with particular interest in the haplotype analysis. <i>Biometrics</i> , <b>2001</b> , 57, 769-78	1.8	9
14	Pacing Rhythms of Wanderers with Dementia. <i>Psychogeriatrics</i> , <b>2001</b> , 1, 76-81	1.8	0
13	A case of schizophrenia with twilight state due to nonconvulsive status epilepticus (NCSE). <i>International Clinical Psychopharmacology</i> , <b>2001</b> , 16, 304	2.2	
12	Protective effect of histidine on MPP+-induced hydroxyl radical generation in rat striatum. <i>Brain Research</i> , <b>1999</b> , 817, 206-8	3.7	6
11	Association between a polymorphism in the promoter region of the dopamine D2 receptor gene and schizophrenia in Japanese subjects: replication and evaluation for antipsychotic-related features. <i>International Journal of Neuropsychopharmacology</i> , <b>1999</b> , 2, 181-186	5.8	29
10	Criminal offenses among discharged mentally ill individuals. Determinants of the duration from discharge and absence of diagnostic specificity. <i>International Journal of Law and Psychiatry</i> , <b>1998</b> , 21, 197-207	2.6	15
9	Search for a Susceptibility Locus to Tardive Dyskinesia. <i>Human Psychopharmacology</i> , <b>1997</b> , 12, 35-39	2.3	26
8	Dopamine transporter gene polymorphism and psychiatric symptoms seen in schizophrenic patients at their first episode. <i>American Journal of Medical Genetics Part A</i> , <b>1996</b> , 67, 406-8		30
7	Effects of a single injection of methamphetamine on central dopaminergic systems in rats with hippocampal lesions. <i>Neuroscience Research Communications</i> , <b>1996</b> , 19, 1-7		

6	Current topics in neuroleptic-induced extrapyramidal symptoms in Japan. <i>Keio Journal of Medicine</i> , <b>1996</b> , 45, 95-9	1.6	22
5	Mentally disordered criminal offenders: five years' data from the Tokyo district public prosecutor's office. <i>International Journal of Law and Psychiatry</i> , <b>1995</b> , 18, 221-30	2.6	8
4	Rolipram, a selective c-AMP phosphodiesterase inhibitor suppresses oro-facial dyskinesic movements in rats. <i>Life Sciences</i> , <b>1995</b> , 56, PL443-7	6.8	30
3	Current topics in tardive dyskinesia in Japan. <i>Psychiatry and Clinical Neurosciences</i> , <b>1995</b> , 49, 239-44	6.2	20
2	Extraction of Potential Difficult-to-Manage Psychiatric Cases among Criminal Offenders: 5-Year Data from the Tokyo District Public Prosecutors Office. <i>Psychiatry and Clinical Neurosciences</i> , <b>1994</b> , 48, 729-735	6.2	
1	The life expectancy of schizophrenic patients with tardive dyskinesia. <i>Human Psychopharmacology</i> , <b>1992</b> , 7, 249-254	2.3	3