

Takuya Imatoh

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6300609/publications.pdf>

Version: 2024-02-01

26
papers

300
citations

1039880

9
h-index

887953

17
g-index

26
all docs

26
docs citations

26
times ranked

544
citing authors

#	ARTICLE	IF	CITATIONS
1	Adiponectin Levels Associated with the Development of Hypertension: A Prospective Study. Hypertension Research, 2008, 31, 229-233.	1.5	57
2	Coffee consumption but not green tea consumption is associated with adiponectin levels in Japanese males. European Journal of Nutrition, 2011, 50, 279-284.	1.8	52
3	Evaluation of the fatty liver index as a predictor for the development of diabetes among insurance beneficiaries with prediabetes. Journal of Diabetes Investigation, 2015, 6, 309-316.	1.1	23
4	Coffee but not green tea consumption is associated with prevalence and severity of hepatic steatosis: the impact on leptin level. European Journal of Clinical Nutrition, 2015, 69, 1023-1027.	1.3	21
5	Is heat shock protein 60 associated with type 2 diabetes mellitus?. Diabetes Research and Clinical Practice, 2009, 85, 208-212.	1.1	15
6	A possible role for HLA-DRB1*04:06 in statin-related myopathy in Japanese patients. Drug Metabolism and Pharmacokinetics, 2016, 31, 467-470.	1.1	15
7	Influence of Japanese Regulatory Action on Denosumab-Related Hypocalcemia Using Japanese Adverse Drug Event Report Database. Biological and Pharmaceutical Bulletin, 2017, 40, 1447-1453.	0.6	13
8	Interaction of low serum adiponectin levels and smoking on coronary stenosis in Japanese men. International Journal of Cardiology, 2006, 110, 251-255.	0.8	12
9	Pharmacogenomic information in the Warning section of drug labels: A comparison between labels in the United States and those in five other countries/regions. Journal of Clinical Pharmacy and Therapeutics, 2018, 43, 493-499.	0.7	11
10	Association between dipeptidyl peptidase-4 inhibitors and urinary tract infection in elderly patients: A retrospective cohort study. Pharmacoepidemiology and Drug Safety, 2018, 27, 931-939.	0.9	10
11	Association between C-reactive protein and risk of overall and 18 site-specific cancers in a Japanese case-cohort. British Journal of Cancer, 2022, 126, 1481-1489.	2.9	9
12	Association between infection and severe drug adverse reactions: an analysis using data from the Japanese Adverse Drug Event Report database. European Journal of Clinical Pharmacology, 2017, 73, 1643-1653.	0.8	8
13	Identification of risk factors and development of detection algorithm for denosumab-induced hypocalcaemia. Journal of Clinical Pharmacy and Therapeutics, 2019, 44, 62-68.	0.7	8
14	Development of a novel algorithm for detecting glucocorticoid-induced diabetes mellitus using a medical information database. Journal of Clinical Pharmacy and Therapeutics, 2017, 42, 215-220.	0.7	7
15	<i>ALDH2</i> Polymorphism rs671, but Not <i>ADH1B</i> Polymorphism rs1229984, Increases Risk for Hypo-€HDL-€Cholesterolemia in a/a Carriers Compared to the G/G Carriers. Lipids, 2018, 53, 797-807.	0.7	7
16	Moderate Oxidative Stress and High Antioxidative Activity Are Associated with Steatosis in Japanese Males. Clinical and Translational Science, 2013, 6, 45-49.	1.5	6
17	Population/regional differences in efficacy of 3 drug categories (antidiabetic, respiratory and) Tj ETQq1 1 0.784314 rgBT /Overlock 10 British Journal of Clinical Pharmacology, 2019, 85, 1270-1282.	1.1	5
18	Does Elevated High-Sensitivity Serum C-Reactive Protein Associate with Hypertension in Non-Obese Japanese Males?. Clinical and Experimental Hypertension, 2007, 29, 395-401.	0.5	4

#	ARTICLE	IF	CITATIONS
19	Associations Between Stevensâ€“Johnson Syndrome and Infection: Overview of Pharmacoepidemiological Studies. <i>Frontiers in Medicine</i> , 2021, 8, 644871.	1.2	4
20	Does antihypertensive treatment with renin-angiotensin system inhibitors prevent the development of diabetic kidney disease?. <i>BMC Pharmacology & Toxicology</i> , 2015, 16, 22.	1.0	3
21	Association between coffee consumption and risk of prostate cancer in Japanese men: a population-based cohort study in Japan. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2021, , cebp.0484.2021.	1.1	3
22	Hyperleptinemia is associated with hypertension in Japanese males. <i>Acta Medica Okayama</i> , 2008, 62, 169-74.	0.1	3
23	Evaluating the impact of regulatory action on denosumabâ€“induced hypocalcaemia in Japan. <i>Journal of Clinical Pharmacy and Therapeutics</i> , 2019, 44, 788-795.	0.7	2
24	Measurement of high-molecular-weight adiponectin is not useful in assessing coronary stenosis. <i>Clinical Chemistry and Laboratory Medicine</i> , 2012, 50, 741-5.	1.4	1
25	The Association Between Concurrence of Infection and the Onset of Severe Eruption or Liver Injury in Patients Using Antipyretic Analgesics: A Matched, Nested Caseâ€“Control Study. <i>Journal of Clinical Pharmacology</i> , 2020, 60, 1177-1184.	1.0	1
26	No causal impact of serum vascular endothelial growth factor level on temporal changes in body mass index in Japanese male workers: a five-year longitudinal study. <i>Endocrine</i> , 2017, 55, 831-838.	1.1	0