

Jeong-Sik Yu

List of Publications by Citations

Source: <https://exaly.com/author-pdf/6305871/jeong-sik-yu-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

129
papers

2,638
citations

31
h-index

46
g-index

138
ext. papers

2,981
ext. citations

5
avg, IF

4.68
L-index

| # | Paper | IF | Citations |
|-----|--|------|-----------|
| 129 | Urachal remnant diseases: spectrum of CT and US findings. <i>Radiographics</i> , 2001 , 21, 451-61 | 5.4 | 218 |
| 128 | Nontumorous hepatic arterial-portal venous shunts: MR imaging findings. <i>Radiology</i> , 2000 , 217, 750-6 | 20.5 | 104 |
| 127 | Restaging of Rectal Cancer with MR Imaging after Concurrent Chemotherapy and Radiation Therapy. <i>Radiographics</i> , 2010 , 30, 503-16 | 5.4 | 92 |
| 126 | Greater and lesser omenta: normal anatomy and pathologic processes. <i>Radiographics</i> , 2007 , 27, 707-20 | 5.4 | 92 |
| 125 | Intraabdominal complications secondary to ventriculoperitoneal shunts: CT findings and review of the literature. <i>American Journal of Roentgenology</i> , 2009 , 193, 1311-7 | 5.4 | 86 |
| 124 | Capecitabine combined with gemcitabine (CapGem) as first-line treatment in patients with advanced/metastatic biliary tract carcinoma. <i>Cancer</i> , 2005 , 104, 2753-8 | 6.4 | 86 |
| 123 | Abdominal cocoon: preoperative diagnostic clues from radiologic imaging with pathologic correlation. <i>American Journal of Roentgenology</i> , 2004 , 182, 639-41 | 5.4 | 66 |
| 122 | Patient selection guidelines in MR-guided focused ultrasound surgery of uterine fibroids: a pictorial guide to relevant findings in screening pelvic MRI. <i>European Radiology</i> , 2008 , 18, 2997-3006 | 8 | 64 |
| 121 | Hepatic cavernous hemangioma: temporal peritumoral enhancement during multiphase dynamic MR imaging. <i>Radiology</i> , 2000 , 216, 692-7 | 20.5 | 61 |
| 120 | Predisposing factors of bile duct injury after transcatheter arterial chemoembolization (TACE) for hepatic malignancy. <i>CardioVascular and Interventional Radiology</i> , 2002 , 25, 270-4 | 2.7 | 59 |
| 119 | Nontumorous arterioportal shunts in the liver: CT and MRI findings considering mechanisms and fate. <i>European Radiology</i> , 2010 , 20, 385-94 | 8 | 57 |
| 118 | Added value of diffusion-weighted imaging in the MRI assessment of perilesional tumor recurrence after chemoembolization of hepatocellular carcinomas. <i>Journal of Magnetic Resonance Imaging</i> , 2009 , 30, 153-60 | 5.6 | 53 |
| 117 | Diagnostic accuracy of multidetector row computed tomography in T- and N staging of gastric cancer with histopathologic correlation. <i>Journal of Computer Assisted Tomography</i> , 2006 , 30, 372-7 | 2.2 | 52 |
| 116 | Recurrent pyogenic cholangitis: comparison between MR cholangiography and direct cholangiography. <i>Radiology</i> , 2001 , 220, 677-82 | 20.5 | 50 |
| 115 | Transcatheter arterial chemoembolization for hepatocellular carcinoma: anatomic and hemodynamic considerations in the hepatic artery and portal vein. <i>Radiographics</i> , 2002 , 22, 1077-91 | 5.4 | 49 |
| 114 | Nodal staging of rectal cancer: high-resolution pelvic MRI versus 18 F-FDGPET/CT. <i>Journal of Computer Assisted Tomography</i> , 2011 , 35, 531-4 | 2.2 | 46 |
| 113 | Acute cholecystitis at T2-weighted and manganese-enhanced T1-weighted MR cholangiography: preliminary study. <i>Radiology</i> , 2003 , 227, 580-4 | 20.5 | 45 |

| | | | |
|-----|--|------|----|
| 112 | Biliary ductal involvement of hilar cholangiocarcinoma: multidetector computed tomography versus magnetic resonance cholangiography. <i>Journal of Computer Assisted Tomography</i> , 2007 , 31, 72-8 | 2.2 | 41 |
| 111 | Dynamic subtraction MR imaging of the liver: advantages and pitfalls. <i>American Journal of Roentgenology</i> , 2003 , 180, 1351-7 | 5.4 | 41 |
| 110 | Dynamic subtraction magnetic resonance imaging of cirrhotic liver: assessment of high signal intensity lesions on nonenhanced T1-weighted images. <i>Journal of Computer Assisted Tomography</i> , 2005 , 29, 51-8 | 2.2 | 40 |
| 109 | Early biliary complications of laparoscopic cholecystectomy: evaluation on T2-weighted MR cholangiography in conjunction with mangafodipir trisodium-enhanced 3D T1-weighted MR cholangiography. <i>American Journal of Roentgenology</i> , 2004 , 183, 1559-66 | 5.4 | 38 |
| 108 | CT angiography of the renal arteries: comparison of lower-tube-voltage CTA with moderate-concentration iodinated contrast material and conventional CTA. <i>American Journal of Roentgenology</i> , 2012 , 199, 96-102 | 5.4 | 37 |
| 107 | Fat-containing nodules in the cirrhotic liver: chemical shift MRI features and clinical implications. <i>American Journal of Roentgenology</i> , 2007 , 188, 1009-16 | 5.4 | 35 |
| 106 | Hepatocellular carcinoma variants: radiologic-pathologic correlation. <i>American Journal of Roentgenology</i> , 2009 , 193, W7-13 | 5.4 | 34 |
| 105 | Radiologic findings of gastrointestinal tract involvement in hepatocellular carcinoma. <i>Journal of Computer Assisted Tomography</i> , 2002 , 26, 95-101 | 2.2 | 34 |
| 104 | Detection of small intrahepatic metastases of hepatocellular carcinomas using diffusion-weighted imaging: comparison with conventional dynamic MRI. <i>Magnetic Resonance Imaging</i> , 2011 , 29, 985-92 | 3.3 | 33 |
| 103 | Small hypervascular enhancing lesions on arterial phase images of multiphase dynamic computed tomography in cirrhotic liver: fate and implications. <i>Journal of Computer Assisted Tomography</i> , 2008 , 32, 39-45 | 2.2 | 33 |
| 102 | Radiological and clinical features of sarcomatoid hepatocellular carcinoma in 11 cases. <i>Journal of Computer Assisted Tomography</i> , 2008 , 32, 745-9 | 2.2 | 33 |
| 101 | Bile duct injuries leading to portal vein obliteration after transcatheter arterial chemoembolization in the liver: CT findings and initial observations. <i>Radiology</i> , 2001 , 221, 429-36 | 20.5 | 33 |
| 100 | Comparison of multishot turbo spin echo and HASTE sequences for T2-weighted MRI of liver lesions. <i>Journal of Magnetic Resonance Imaging</i> , 1998 , 8, 1079-84 | 5.6 | 31 |
| 99 | Radiographic findings of primary B-cell lymphoma of the stomach: low-grade versus high-grade malignancy in relation to the mucosa-associated lymphoid tissue concept. <i>American Journal of Roentgenology</i> , 2002 , 179, 1297-304 | 5.4 | 31 |
| 98 | Mucinous versus nonmucinous gastric carcinoma: differentiation with helical CT. <i>Radiology</i> , 2002 , 223, 540-6 | 20.5 | 30 |
| 97 | Small hypervascular hepatocellular carcinoma: limited value of portal and delayed phases on dynamic magnetic resonance imaging. <i>Acta Radiologica</i> , 2008 , 49, 735-43 | 2 | 28 |
| 96 | Preoperative staging accuracy of multidetector row computed tomography for extrahepatic bile duct carcinoma. <i>Journal of Computer Assisted Tomography</i> , 2006 , 30, 362-7 | 2.2 | 26 |
| 95 | CT venography for deep vein thrombosis using a low tube voltage (100 kVp) setting could increase venous enhancement and reduce the amount of administered iodine. <i>Korean Journal of Radiology</i> , 2013 , 14, 183-93 | 6.9 | 22 |

| | | | |
|----|---|-----|----|
| 94 | Lymphoepithelial cysts in the pancreas: MRI of two cases with emphasis of diffusion-weighted imaging characteristics. <i>Journal of Magnetic Resonance Imaging</i> , 2010 , 32, 692-6 | 5.6 | 22 |
| 93 | Hepatic metastases: perilesional enhancement on dynamic MRI. <i>American Journal of Roentgenology</i> , 2006 , 186, 1051-8 | 5.4 | 22 |
| 92 | Advanced gastric carcinoma with signet ring cell carcinoma versus non-signet ring cell carcinoma: differentiation with multidetector CT. <i>Journal of Computer Assisted Tomography</i> , 2006 , 30, 880-4 | 2.2 | 22 |
| 91 | Diffusion-Weighted MRI of Malignant versus Benign Portal Vein Thrombosis. <i>Korean Journal of Radiology</i> , 2016 , 17, 533-40 | 6.9 | 22 |
| 90 | Hypervascular hepatocellular carcinoma in the cirrhotic liver: diffusion-weighted imaging versus superparamagnetic iron oxide-enhanced MRI. <i>Magnetic Resonance Imaging</i> , 2011 , 29, 1235-43 | 3.3 | 21 |
| 89 | Choledochal cyst and anomalous pancreaticobiliary ductal union in adults: radiological spectrum and complications. <i>Journal of Computer Assisted Tomography</i> , 2008 , 32, 17-22 | 2.2 | 21 |
| 88 | Value of nonvisualized primary lesions of gastric cancer on preoperative MDCT. <i>American Journal of Roentgenology</i> , 2007 , 189, W315-9 | 5.4 | 21 |
| 87 | Pure epidural cavernous hemangioma of the cervical spine that presented with an acute sensory deficit caused by hemorrhage. <i>Yonsei Medical Journal</i> , 2006 , 47, 877-80 | 3 | 21 |
| 86 | Adrenocortical oncocytoma displaying intense activity on 18F-FDG-PET: a case report and a literature review. <i>Annals of Nuclear Medicine</i> , 2008 , 22, 821-4 | 2.5 | 20 |
| 85 | Transient peritumoral enhancement during dynamic MRI of the liver: cavernous hemangioma versus hepatocellular carcinoma. <i>Journal of Computer Assisted Tomography</i> , 2002 , 26, 411-7 | 2.2 | 19 |
| 84 | Unusual cystic neoplasms in the pancreas: radiologic-pathologic correlation. <i>Journal of Computer Assisted Tomography</i> , 2005 , 29, 610-6 | 2.2 | 18 |
| 83 | Magnetic resonance imaging of arterioportal shunts in the liver. <i>Topics in Magnetic Resonance Imaging</i> , 2002 , 13, 165-76 | 2.3 | 18 |
| 82 | Double contrast-enhanced MRI of viral hepatitis-induced cirrhosis: correlation of gross morphological signs with hepatic fibrosis. <i>British Journal of Radiology</i> , 2010 , 83, 212-7 | 3.4 | 16 |
| 81 | Newly developed hepatocellular carcinoma (HCC) in chronic liver disease: MR imaging findings before the diagnosis of HCC. <i>Journal of Computer Assisted Tomography</i> , 2006 , 30, 765-71 | 2.2 | 16 |
| 80 | Eosinophilic hepatic necrosis: magnetic resonance imaging and computed tomography comparison. <i>Journal of Computer Assisted Tomography</i> , 2005 , 29, 765-71 | 2.2 | 16 |
| 79 | Evaluation of lateral pelvic nodes in patients with advanced rectal cancer. <i>American Journal of Roentgenology</i> , 2014 , 202, 1245-55 | 5.4 | 15 |
| 78 | Sclerotic changes of cavernous hemangioma in the cirrhotic liver: long-term follow-up using dynamic contrast-enhanced computed tomography. <i>Radiologia Medica</i> , 2020 , 125, 1225-1232 | 6.5 | 14 |
| 77 | A case of mass-forming splenic tuberculosis: MRI findings with emphasis of diffusion-weighted imaging characteristics. <i>Journal of Korean Medical Science</i> , 2011 , 26, 457-60 | 4.7 | 14 |

| | | | |
|----|---|------|----|
| 76 | Prostate cancer: added value of subtraction dynamic imaging in 3T magnetic resonance imaging with a phased-array body coil. <i>Yonsei Medical Journal</i> , 2008 , 49, 765-74 | 3 | 14 |
| 75 | A Phase II study of capecitabine combined with gemcitabine in patients with advanced gallbladder carcinoma. <i>Yonsei Medical Journal</i> , 2005 , 46, 526-31 | 3 | 14 |
| 74 | Percutaneous placement of self-expandable metallic biliary stents in malignant extrahepatic strictures: indications of transpapillary and suprapapillary methods. <i>Korean Journal of Radiology</i> , 2000 , 1, 65-72 | 6.9 | 14 |
| 73 | Small hypervascular hepatocellular carcinomas: value of "washout" on gadolinium-enhanced dynamic MR imaging compared to superparamagnetic iron oxide-enhanced imaging. <i>European Radiology</i> , 2009 , 19, 2614-22 | 8 | 13 |
| 72 | Hyperintense nodules on non-enhanced T1-weighted gradient-echo magnetic resonance imaging of cirrhotic liver: fate and clinical implications. <i>Journal of Magnetic Resonance Imaging</i> , 2006 , 24, 630-6 | 5.6 | 13 |
| 71 | Abdominal sparganosis presenting as an abscess with fistulous communication to the bowel. <i>American Journal of Roentgenology</i> , 2005 , 185, 1084-5 | 5.4 | 13 |
| 70 | Focal eosinophilic necrosis versus metastasis in the liver: the usefulness of two-phase dynamic CT. <i>American Journal of Roentgenology</i> , 2005 , 184, 1085-90 | 5.4 | 12 |
| 69 | Prediction of pancreatic fistula after pancreatoduodenectomy by preoperative dynamic CT and fecal elastase-1 levels. <i>PLoS ONE</i> , 2017 , 12, e0177052 | 3.7 | 12 |
| 68 | Optimization of Split-Bolus CT Urography: Effect of Differences in Allocation of Contrast Medium and Prolongation of Imaging Delay. <i>American Journal of Roentgenology</i> , 2017 , 209, W10-W17 | 5.4 | 11 |
| 67 | Segmental difference of the hepatic fibrosis from chronic viral hepatitis due to hepatitis B versus C virus infection: comparison using dual contrast material-enhanced MRI. <i>Korean Journal of Radiology</i> , 2011 , 12, 431-8 | 6.9 | 11 |
| 66 | Hepatic cavernous hemangioma in cirrhotic liver: imaging findings. <i>Korean Journal of Radiology</i> , 2000 , 1, 185-90 | 6.9 | 11 |
| 65 | High-flow haemangiomas versus hypervascular hepatocellular carcinoma showing "pseudo-washout" on gadoteric acid-enhanced hepatic MRI: value of diffusion-weighted imaging in the differential diagnosis of small lesions. <i>Clinical Radiology</i> , 2017 , 72, 247-254 | 2.9 | 10 |
| 64 | Percutaneous biliary approach as a successful rescue procedure after failed endoscopic therapy for drainage in advanced hilar tumors. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2017 , 32, 932-938 | 4.38 | 10 |
| 63 | Transient hepatic attenuation difference (THAD) following transcatheter arterial chemoembolization for hepatic malignancy: changes on serial CT examinations. <i>European Radiology</i> , 2008 , 18, 1596-603 | 8 | 10 |
| 62 | Distinguishing hepatic metastases from hemangiomas: qualitative and quantitative diagnostic performance through dual echo respiratory-triggered fast spin echo magnetic resonance imaging. <i>Journal of Computer Assisted Tomography</i> , 2005 , 29, 571-9 | 2.2 | 10 |
| 61 | Hepatic cavernous hemangiomas: relationship between speed of intratumoral enhancement during dynamic MRI and apparent diffusion coefficient on diffusion-weighted imaging. <i>Korean Journal of Radiology</i> , 2012 , 13, 728-35 | 6.9 | 9 |
| 60 | Focal eosinophilic necrosis on superparamagnetic iron oxide-enhanced MRI. <i>American Journal of Roentgenology</i> , 2010 , 194, 1296-302 | 5.4 | 9 |
| 59 | Gadobenate dimeglumine-enhanced MRI of intraductal papillary mucinous tumor of the bile ducts. <i>Journal of Magnetic Resonance Imaging</i> , 2007 , 25, 625-7 | 5.6 | 9 |

| | | | |
|----|---|------|---|
| 58 | MR cholangiography of accessory bile duct connected to the stomach. <i>American Journal of Roentgenology</i> , 2007 , 189, W344-7 | 5.4 | 9 |
| 57 | Hepatic MRI using the double-echo chemical shift phase-selective gradient-echo technique. <i>American Journal of Roentgenology</i> , 2007 , 188, W49-56 | 5.4 | 9 |
| 56 | Retroperitoneal paragonimiasis: a case of ectopic paragonimiasis presenting as periureteral masses. <i>Journal of Computer Assisted Tomography</i> , 1999 , 23, 696-8 | 2.2 | 9 |
| 55 | Computed tomographic venography for varicose veins of the lower extremities: prospective comparison of 80-kVp and conventional 120-kVp protocols. <i>Journal of Computer Assisted Tomography</i> , 2012 , 36, 583-90 | 2.2 | 8 |
| 54 | Large (>or=2 cm) non-hypervascular nodules depicted on MRI in the cirrhotic liver: fate and implications. <i>Clinical Radiology</i> , 2008 , 63, 1121-30 | 2.9 | 8 |
| 53 | Inflammatory myofibroblastic tumors in the liver: MRI of two immunohistochemically-verified cases. <i>Journal of Magnetic Resonance Imaging</i> , 2007 , 26, 418-21 | 5.6 | 8 |
| 52 | Apparent diffusion coefficient of hepatocellular carcinoma on diffusion-weighted imaging: Histopathologic tumor grade versus arterial vascularity during dynamic magnetic resonance imaging. <i>PLoS ONE</i> , 2018 , 13, e0197070 | 3.7 | 7 |
| 51 | Usefulness of a lead shielding device for reducing the radiation dose to tissues outside the primary beams during CT. <i>Radiologia Medica</i> , 2014 , 119, 951-957 | 6.5 | 7 |
| 50 | Feasibility of 5-minute delayed transition phase imaging with 30° flip angle in gadoteric acid-enhanced 3D gradient-echo MRI of liver, compared with 20-minute delayed hepatocyte phase MRI with standard 10° flip angle. <i>American Journal of Roentgenology</i> , 2015 , 204, 69-75 | 5.4 | 7 |
| 49 | Preoperative evaluation of lymphovascular invasion using high-resolution pelvic magnetic resonance in patients with rectal cancer: a 2-year follow-up study. <i>Journal of Computer Assisted Tomography</i> , 2013 , 37, 583-8 | 2.2 | 7 |
| 48 | Value of manganese-enhanced T1- and T2-weighted MR cholangiography for differentiating cystic parenchymal lesions from cystic abnormalities which communicate with bile ducts. <i>Yonsei Medical Journal</i> , 2007 , 48, 1072-4 | 3 | 7 |
| 47 | Stent-graft for TIPS in a hepatocellular carcinoma patient with main portal vein invasion. <i>American Journal of Roentgenology</i> , 2004 , 182, 1301-4 | 5.4 | 7 |
| 46 | Efficacy of a paclitaxel-eluting biliary metal stent with sodium caprate in malignant biliary obstruction: a prospective randomized comparative study. <i>Endoscopy</i> , 2019 , 51, 843-851 | 3.4 | 7 |
| 45 | Comparison of abdominal ultrasonographic findings with endoscopic ultrasonographic findings of solid pseudopapillary neoplasms of the pancreas. <i>Ultrasound Quarterly</i> , 2014 , 30, 173-8 | 1.4 | 6 |
| 44 | Diagnosing small hepatic cysts on multidetector CT: an additional merit of thinner coronal reformations. <i>Korean Journal of Radiology</i> , 2011 , 12, 341-50 | 6.9 | 6 |
| 43 | Hypervascular focus in the nonhypervascular nodule ("nodule-in-nodule") on dynamic computed tomography: imaging evidence of aggressive progression in hepatocellular carcinoma. <i>Journal of Computer Assisted Tomography</i> , 2009 , 33, 131-5 | 2.2 | 6 |
| 42 | Intratumoral Blood Flow in Cavernous Hemangioma of the Liver: Radiologic-Pathologic Correlation. <i>Radiology</i> , 1998 , 208, 549-549 | 20.5 | 6 |
| 41 | Prognostic value of gallbladder wall thickening in patients with acute hepatitis A. <i>Ultrasonography</i> , 2015 , 34, 139-43 | 4.3 | 6 |

| | | | |
|----|--|------|---|
| 40 | Validation of 10-Minute Delayed Hepatocyte Phase Imaging with 30° Flip Angle in Gadoteric Acid-Enhanced MRI for the Detection of Liver Metastasis. <i>PLoS ONE</i> , 2015 , 10, e0139863 | 3.7 | 5 |
| 39 | Limited value of diffusion-weighted MR imaging for differentiating bland from malignant portal venous thrombi. <i>Radiology</i> , 2010 , 256, 673-4; author reply 674 | 20.5 | 5 |
| 38 | Diagnostic accuracy of multidetector-row computed tomography for common bile duct calculi: is it necessary to add non-contrast-enhanced images to contrast-enhanced images?. <i>Journal of Computer Assisted Tomography</i> , 2007 , 31, 508-12 | 2.2 | 5 |
| 37 | Isolated adrenal hemorrhage after minor blunt trauma. <i>American Journal of Emergency Medicine</i> , 2007 , 25, 984.e5-6 | 2.9 | 5 |
| 36 | ULTRASONOGRAPHY: a new beginning with international open access. <i>Ultrasonography</i> , 2014 , 33, 1 | 4.3 | 5 |
| 35 | Feasibility of 10-Minute Delayed Hepatocyte Phase Imaging Using a 30° Flip Angle in Gd-EOB-DTPA-Enhanced Liver MRI for the Detection of Hepatocellular Carcinoma in Patients with Chronic Hepatitis or Cirrhosis. <i>PLoS ONE</i> , 2016 , 11, e0167701 | 3.7 | 5 |
| 34 | Hepatic cavernous hemangiomas: long-term (>5 years) follow-up changes on contrast-enhanced dynamic computed tomography or magnetic resonance imaging and determinant factors of the size change. <i>Radiologia Medica</i> , 2018 , 123, 323-330 | 6.5 | 4 |
| 33 | Diffusion-weighted imaging versus superparamagnetic iron oxide (SPIO)-enhanced MRI: exclusive and combined values in the assessment of hepatic metastases. <i>Magnetic Resonance Imaging</i> , 2012 , 30, 554-61 | 3.3 | 4 |
| 32 | Pneumoperitoneum caused by transhepatic air leak after metallic biliary stent placement. <i>CardioVascular and Interventional Radiology</i> , 2000 , 23, 482-4 | 2.7 | 4 |
| 31 | ULTRASONOGRAPHY: on the way to becoming an internationally renowned journal. <i>Ultrasonography</i> , 2015 , 34, 1-2 | 4.3 | 4 |
| 30 | ULTRASONOGRAPHY: two years of globalization and the achievement of ESCI status. <i>Ultrasonography</i> , 2016 , 35, 1-2 | 4.3 | 4 |
| 29 | Texture Analysis of Hepatocellular Carcinoma on Magnetic Resonance Imaging: Assessment for Performance in Predicting Histopathologic Grade. <i>Journal of Computer Assisted Tomography</i> , 2020 , 44, 901-910 | 2.2 | 3 |
| 28 | Three cases of pancreatic pseudocysts associated with dorsal pancreatic agenesis. <i>Radiology Case Reports</i> , 2019 , 14, 79-82 | 1 | 3 |
| 27 | Subtraction Images From Portal Venous Phase Gadoteric Acid-Enhanced MRI for Observing Washout and Enhancing Capsule Features in LI-RADS Version 2018. <i>American Journal of Roentgenology</i> , 2020 , 214, 72-80 | 5.4 | 3 |
| 26 | Diffusion-weighted MR imaging before and after contrast enhancement with superparamagnetic iron oxide for assessment of hepatic metastasis. <i>Yonsei Medical Journal</i> , 2012 , 53, 825-33 | 3 | 2 |
| 25 | Islet Cell Tumors of the Pancreas: A Variety of Multiphase Dynamic Imaging Findings with Pathologic Correlations Focusing on Nonfunctioning Tumors and Insulinomas. <i>Journal of the Korean Society of Radiology</i> , 2010 , 63, 463 | 0.2 | 2 |
| 24 | Pseudoglandular Formation in Hepatocellular Carcinoma Determines Apparent Diffusion Coefficient in Diffusion-Weighted MRI 2018 , 22, 79 | | 2 |
| 23 | ULTRASONOGRAPHY: how to raise the impact factor. <i>Ultrasonography</i> , 2017 , 36, 1-2 | 4.3 | 2 |

| | | | |
|----|---|------|---|
| 22 | Effect of hepatic steatosis on native T1 mapping of 3T magnetic resonance imaging in the assessment of T1 values for patients with non-alcoholic fatty liver disease. <i>Magnetic Resonance Imaging</i> , 2021 , 80, 1-8 | 3.3 | 2 |
| 21 | Multidetector CT findings of histopathologically proven peritoneal tuberculous cold abscesses. <i>Radiologia Medica</i> , 2017 , 122, 248-256 | 6.5 | 1 |
| 20 | Factors affecting survival after concurrent chemoradiation therapy for advanced hepatocellular carcinoma: a retrospective study. <i>Radiation Oncology</i> , 2017 , 12, 133 | 4.2 | 1 |
| 19 | Serous Cystic Neoplasms of the Pancreas: Endoscopic Ultrasonographic Versus Computed Tomography and Magnetic Resonance Imaging Features of Surgically Removed Masses. <i>Ultrasound Quarterly</i> , 2018 , 34, 122-127 | 1.4 | 1 |
| 18 | Attenuation-Based Automatic Tube Potential Selection in Cerebral Computed Tomography Angiography: Effects on Radiation Exposure and Image Quality. <i>Journal of the Korean Society of Radiology</i> , 2018 , 78, 35 | 0.2 | 1 |
| 17 | Preoperative Evaluation of Lower Rectal Cancer by Pelvic MR with and without Gel Filling. <i>Journal of the Korean Society of Magnetic Resonance in Medicine</i> , 2014 , 18, 323 | | 1 |
| 16 | Corona enhancement can substitute enhancing capsule in the imaging diagnosis of small (≤ 1 cm) HCCs on gadoteric acid-enhanced MRI. <i>European Radiology</i> , 2021 , 31, 8628-8637 | 8 | 1 |
| 15 | IgG4-Related Hepatic Inflammatory Pseudotumor Complicated by Actinomycosis during Steroid Therapy. <i>Journal of the Korean Society of Radiology</i> , 2019 , 80, 351 | 0.2 | 0 |
| 14 | Iatrogenic Injury in the Lateral Segment of the Liver after Pancreatoduodenectomy: Early Follow-Up CT Features and Clinical Implications. <i>Journal of the Korean Society of Radiology</i> , 2012 , 66, 451 | 0.2 | 0 |
| 13 | Diffusion-Weighted MR Imaging of Upper Abdomen: Comparison of Breath-Hold, Free-Breathing, and Respiratory-Triggered Techniques. <i>Journal of the Korean Society of Radiology</i> , 2011 , 64, 465 | 0.2 | |
| 12 | Ferucarbotran-Enhanced Hepatic MRI at 3T Unit: Quantitative and Qualitative Comparison of Fast Breath-hold Imaging Sequences. <i>Journal of the Korean Society of Magnetic Resonance in Medicine</i> , 2010 , 14, 31 | | |
| 11 | Alpha-fetoprotein-producing Gastric Cancer with Metastasis to the Scrotum: Case Report. <i>Journal of the Korean Society of Radiology</i> , 2010 , 63, 471 | 0.2 | |
| 10 | Ferucarbotran-enhanced T2-weighted magnetic resonance imaging: differentiation of hepatic cavernous hemangiomas from malignant solid lesions. <i>Abdominal Imaging</i> , 2009 , 34, 494-501 | | |
| 9 | Letter to the editor re: Diffusion-weighted magnetic resonance imaging of pancreas tumours. <i>European Radiology</i> , 2010 , 20, 1768-9; author reply 1770-1 | 8 | |
| 8 | Hepatocellular carcinoma: progression in patients with hepatitis B versus C virus infection. <i>Radiology</i> , 2008 , 249, 1082; author reply 1082-3 | 20.5 | |
| 7 | Secondary achalasia caused by peripheral cholangiocarcinoma in left lobe of the liver. <i>European Journal of Radiology Extra</i> , 2004 , 50, 89-92 | | |
| 6 | Retrospective Comparison of Renal Ultrasonographic and Clinical Findings in Patients with Rhabdomyolysis. <i>Journal of Medical Ultrasound</i> , 2020 , 28, 151-155 | 0.8 | |
| 5 | Segmental Difference of the Hepatic Fibrosis from Chronic Viral Hepatitis due to Hepatitis B versus C Virus Infection: Comparison Using Dual Contrast Material-Enhanced MRI. <i>Korean Journal of Radiology</i> , 2011 , 12, 440 | 6.9 | |

- | | | |
|---|---|-----|
| 4 | Ultrasonography : road to SCIE listing. <i>Science Editing</i> , 2020 , 7, 194-197 | 1.2 |
| 3 | Multidetector CT Evaluation of Food Stasis in Remnant Stomach and Body Fat Change after Subtotal Gastrectomy by Laparoscopic versus Open Abdominal Approach. <i>Journal of the Korean Society of Radiology</i> , 2013 , 68, 33 | 0.2 |
| 2 | Preoperative Multidetector CT Manifestations of Perigastric Lymph Nodes in Patients with Early Gastric Cancer and pN0. <i>Journal of the Korean Society of Radiology</i> , 2013 , 69, 391 | 0.2 |
| 1 | Advanced Gastric Cancer: Differentiation of Borrmann Type IV versus Borrmann Type III by Two-Phased Dynamic Multi-Detector Row CT with Use of the Water Filling Method. <i>Journal of the Korean Society of Radiology</i> , 2013 , 68, 117 | 0.2 |