Jeong-Sik Yu

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/6305871/jeong-sik-yu-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

 129
 2,638
 31
 46

 papers
 citations
 h-index
 g-index

 138
 2,981
 5
 4.68

 ext. papers
 ext. citations
 avg, IF
 L-index

#	Paper	IF	Citations
129	Corona enhancement can substitute enhancing capsule in the imaging diagnosis of small (IB cm) HCCs on gadoxetic acid-enhanced MRI. <i>European Radiology</i> , 2021 , 31, 8628-8637	8	1
128	Effect of hepatic steatosis on native T1 mapping of 3T magnetic resonance imaging in the assessment of T1 values for patients with non-alcoholic fatty liver disease. <i>Magnetic Resonance Imaging</i> , 2021 , 80, 1-8	3.3	2
127	Sclerotic changes of cavernous hemangioma in the cirrhotic liver: long-term follow-up using dynamic contrast-enhanced computed tomography. <i>Radiologia Medica</i> , 2020 , 125, 1225-1232	6.5	14
126	Retrospective Comparison of Renal Ultrasonographic and Clinical Findings in Patients with Rhabdomyolysis. <i>Journal of Medical Ultrasound</i> , 2020 , 28, 151-155	0.8	
125	Ultrasonography : road to SCIE listing. <i>Science Editing</i> , 2020 , 7, 194-197	1.2	
124	Texture Analysis of Hepatocellular Carcinoma on Magnetic Resonance Imaging: Assessment for Performance in Predicting Histopathologic Grade. <i>Journal of Computer Assisted Tomography</i> , 2020 , 44, 901-910	2.2	3
123	Subtraction Images From Portal Venous Phase Gadoxetic Acid-Enhanced MRI for Observing Washout and Enhancing Capsule Features in LI-RADS Version 2018. <i>American Journal of Roentgenology</i> , 2020 , 214, 72-80	5.4	3
122	IgG4-Related Hepatic Inflammatory Pseudotumor Complicated by Actinomycosis during Steroid Therapy. <i>Journal of the Korean Society of Radiology</i> , 2019 , 80, 351	0.2	О
121	Efficacy of a paclitaxel-eluting biliary metal stent with sodium caprate in malignant biliary obstruction: a prospective randomized comparative study. <i>Endoscopy</i> , 2019 , 51, 843-851	3.4	7
120	Three cases of pancreatic pseudocysts associated with dorsal pancreatic agenesis. <i>Radiology Case Reports</i> , 2019 , 14, 79-82	1	3
119	Serous Cystic Neoplasms of the Pancreas: Endoscopic Ultrasonographic Versus Computed Tomography and Magnetic Resonance Imaging Features of Surgically Removed Masses. <i>Ultrasound Quarterly</i> , 2018 , 34, 122-127	1.4	1
118	Hepatic cavernous hemangiomas: long-term (>ြ@lyears) follow-up changes on contrast-enhanced dynamic computed tomography or magnetic resonance imaging and determinant factors of the size change. <i>Radiologia Medica</i> , 2018 , 123, 323-330	6.5	4
117	Apparent diffusion coefficient of hepatocellular carcinoma on diffusion-weighted imaging: Histopathologic tumor grade versus arterial vascularity during dynamic magnetic resonance imaging. <i>PLoS ONE</i> , 2018 , 13, e0197070	3.7	7
116	Attenuation-Based Automatic Tube Potential Selection in Cerebral Computed Tomography Angiography: Effects on Radiation Exposure and Image Quality. <i>Journal of the Korean Society of Radiology</i> , 2018 , 78, 35	0.2	1
115	Pseudoglandular Formation in Hepatocellular Carcinoma Determines Apparent Diffusion Coefficient in Diffusion-Weighted MRI 2018 , 22, 79		2
114	Optimization of Split-Bolus CT Urography: Effect of Differences in Allocation of Contrast Medium and Prolongation of Imaging Delay. <i>American Journal of Roentgenology</i> , 2017 , 209, W10-W17	5.4	11
113	Multidetector CT findings of histopathologically proven peritoneal tuberculous cold abscesses. <i>Radiologia Medica</i> , 2017 , 122, 248-256	6.5	1

(2013-2017)

112	Factors affecting survival after concurrent chemoradiation therapy for advanced hepatocellular carcinoma: a retrospective study. <i>Radiation Oncology</i> , 2017 , 12, 133	4.2	1	
111	High-flow haemangiomas versus hypervascular hepatocellular carcinoma showing "pseudo-washout" on gadoxetic acid-enhanced hepatic MRI: value of diffusion-weighted imaging in the differential diagnosis of small lesions. <i>Clinical Radiology</i> , 2017 , 72, 247-254	2.9	10	
110	Percutaneous biliary approach as a successful rescue procedure after failed endoscopic therapy for drainage in advanced hilar tumors. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2017 , 32, 93	2-938	10	
109	Prediction of pancreatic fistula after pancreatoduodenectomy by preoperative dynamic CT and fecal elastase-1 levels. <i>PLoS ONE</i> , 2017 , 12, e0177052	3.7	12	
108	ULTRASONOGRAPHY: how to raise the impact factor. <i>Ultrasonography</i> , 2017 , 36, 1-2	4.3	2	
107	ULTRASONOGRAPHY: two years of globalization and the achievement of ESCI status. <i>Ultrasonography</i> , 2016 , 35, 1-2	4.3	4	
106	Diffusion-Weighted MRI of Malignant versus Benign Portal Vein Thrombosis. <i>Korean Journal of Radiology</i> , 2016 , 17, 533-40	6.9	22	
105	Feasibility of 10-Minute Delayed Hepatocyte Phase Imaging Using a 30°IFlip Angle in Gd-EOB-DTPA-Enhanced Liver MRI for the Detection of Hepatocellular Carcinoma in Patients with Chronic Hepatitis or Cirrhosis. <i>PLoS ONE</i> , 2016 , 11, e0167701	3.7	5	
104	Validation of 10-Minute Delayed Hepatocyte Phase Imaging with 30 th Flip Angle in Gadoxetic Acid-Enhanced MRI for the Detection of Liver Metastasis. <i>PLoS ONE</i> , 2015 , 10, e0139863	3.7	5	
103	Feasibility of 5-minute delayed transition phase imaging with 30º flip angle in gadoxetic acid-enhanced 3D gradient-echo MRI of liver, compared with 20-minute delayed hepatocyte phase MRI with standard 10º flip angle. <i>American Journal of Roentgenology</i> , 2015 , 204, 69-75	5.4	7	
102	Prognostic value of gallbladder wall thickening in patients with acute hepatitis A. <i>Ultrasonography</i> , 2015 , 34, 139-43	4.3	6	
101	ULTRASONOGRAPHY: on the way to becoming an internationally renowned journal. <i>Ultrasonography</i> , 2015 , 34, 1-2	4.3	4	
100	Evaluation of lateral pelvic nodes in patients with advanced rectal cancer. <i>American Journal of Roentgenology</i> , 2014 , 202, 1245-55	5.4	15	
99	Usefulness of a lead shielding device for reducing the radiation dose to tissues outside the primary beams during CT. <i>Radiologia Medica</i> , 2014 , 119, 951-957	6.5	7	
98	Comparison of abdominal ultrasonographic findings with endoscopic ultrasonographic findings of solid pseudopapillary neoplasms of the pancreas. <i>Ultrasound Quarterly</i> , 2014 , 30, 173-8	1.4	6	
97	ULTRASONOGRAPHY: a new beginning with international open access. <i>Ultrasonography</i> , 2014 , 33, 1	4.3	5	
96	Preoperative Evaluation of Lower Rectal Cancer by Pelvic MR with and without Gel Filling. <i>Journal of the Korean Society of Magnetic Resonance in Medicine</i> , 2014 , 18, 323		1	
95	Preoperative evaluation of lymphovascular invasion using high-resolution pelvic magnetic resonance in patients with rectal cancer: a 2-year follow-up study. <i>Journal of Computer Assisted Tomography</i> , 2013 , 37, 583-8	2.2	7	

94	CT venography for deep vein thrombosis using a low tube voltage (100 kVp) setting could increase venous enhancement and reduce the amount of administered iodine. <i>Korean Journal of Radiology</i> , 2013 , 14, 183-93	6.9	22
93	Multidetector CT Evaluation of Food Stasis in Remnant Stomach and Body Fat Change after Subtotal Gastrectomy by Laparoscopic versus Open Abdominal Approach. <i>Journal of the Korean Society of Radiology</i> , 2013 , 68, 33	0.2	
92	Preoperative Multidetector CT Manifestations of Perigastric Lymph Nodes in Patients with Early Gastric Cancer and pN0. <i>Journal of the Korean Society of Radiology</i> , 2013 , 69, 391	0.2	
91	Advanced Gastric Cancer: Differentiation of Borrmann Type IV versus Borrmann Type III by Two-Phased Dynamic Multi-Detector Row CT with Use of the Water Filling Method. <i>Journal of the Korean Society of Radiology</i> , 2013 , 68, 117	0.2	
90	Diffusion-weighted imaging versus superparamagnetic iron oxide (SPIO)-enhanced MRI: exclusive and combined values in the assessment of hepatic metastases. <i>Magnetic Resonance Imaging</i> , 2012 , 30, 554-61	3.3	4
89	Hepatic cavernous hemangiomas: relationship between speed of intratumoral enhancement during dynamic MRI and apparent diffusion coefficient on diffusion-weighted imaging. <i>Korean Journal of Radiology</i> , 2012 , 13, 728-35	6.9	9
88	Diffusion-weighted MR imaging before and after contrast enhancement with superparamagnetic iron oxide for assessment of hepatic metastasis. <i>Yonsei Medical Journal</i> , 2012 , 53, 825-33	3	2
87	CT angiography of the renal arteries: comparison of lower-tube-voltage CTA with moderate-concentration iodinated contrast material and conventional CTA. <i>American Journal of Roentgenology</i> , 2012 , 199, 96-102	5.4	37
86	Computed tomographic venography for varicose veins of the lower extremities: prospective comparison of 80-kVp and conventional 120-kVp protocols. <i>Journal of Computer Assisted Tomography</i> , 2012 , 36, 583-90	2.2	8
85	Iatrogenic Injury in the Lateral Segment of the Liver after Pancreatoduodenectomy: Early Follow-Up CT Features and Clinical Implications. <i>Journal of the Korean Society of Radiology</i> , 2012 , 66, 451	0.2	O
84	Diagnosing small hepatic cysts on multidetector CT: an additional merit of thinner coronal reformations. <i>Korean Journal of Radiology</i> , 2011 , 12, 341-50	6.9	6
83	Diffusion-Weighted MR Imaging of Upper Abdomen: Comparison of Breath-Hold, Free-Breathing, and Respiratory-Triggered Techniques. <i>Journal of the Korean Society of Radiology</i> , 2011 , 64, 465	0.2	
82	A case of mass-forming splenic tuberculosis: MRI findings with emphasis of diffusion-weighted imaging characteristics. <i>Journal of Korean Medical Science</i> , 2011 , 26, 457-60	4.7	14
81	Segmental difference of the hepatic fibrosis from chronic viral hepatitis due to hepatitis B versus C virus infection: comparison using dual contrast material-enhanced MRI. <i>Korean Journal of Radiology</i> , 2011 , 12, 431-8	6.9	11
80	Nodal staging of rectal cancer: high-resolution pelvic MRI versus IE-FDGPET/CT. <i>Journal of Computer Assisted Tomography</i> , 2011 , 35, 531-4	2.2	46
79	Detection of small intrahepatic metastases of hepatocellular carcinomas using diffusion-weighted imaging: comparison with conventional dynamic MRI. <i>Magnetic Resonance Imaging</i> , 2011 , 29, 985-92	3.3	33
78	Hypervascular hepatocellular carcinoma in the cirrhotic liver: diffusion-weighted imaging versus superparamagnetic iron oxide-enhanced MRI. <i>Magnetic Resonance Imaging</i> , 2011 , 29, 1235-43	3.3	21
77	Segmental Difference of the Hepatic Fibrosis from Chronic Viral Hepatitis due to Hepatitis B versus C Virus Infection: Comparison Using Dual Contrast Material-Enhanced MRI. <i>Korean Journal of Radiology</i> 2011 12 440	6.9	

(2008-2010)

Breath-hold Imaging Sequences. Journal of the Korean Society of Magnetic Resonance in Medicine, 76 2010, 14, 31 Alpha-fetoprotein-producing Gastric Cancer with Metastasis to the Scrotum: Case Report. Journal 0.2 75 of the Korean Society of Radiology, 2010, 63, 471 Restaging of Rectal Cancer with MR Imaging after Concurrent Chemotherapy and Radiation 92 74 5.4 Therapy. Radiographics, 2010, 30, 503-16 Double contrast-enhanced MRI of viral hepatitis-induced cirrhosis: correlation of gross 16 73 3.4 morphological signs with hepatic fibrosis. British Journal of Radiology, 2010, 83, 212-7 Limited value of diffusion-weighted MR imaging for differentiating bland from malignant portal 72 20.5 5 venous thrombi. Radiology, 2010, 256, 673-4; author reply 674 Focal eosinophilic necrosis on superparamagnetic iron oxide-enhanced MRI. American Journal of 71 5.4 9 Roentgenology, **2010**, 194, 1296-302 Nontumorous arterioportal shunts in the liver: CT and MRI findings considering mechanisms and 8 70 57 fate. European Radiology, **2010**, 20, 385-94 Letter to the editor re: Diffusion-weighted magnetic resonance imaging of pancreas tumours. 8 69 European Radiology, 2010, 20, 1768-9; author reply 1770-1 Lymphoepithelial cysts in the pancreas: MRI of two cases with emphasis of diffusion-weighted 68 5.6 22 imaging characteristics. Journal of Magnetic Resonance Imaging, 2010, 32, 692-6 Islet Cell Tumors of the Pancreas: A Variety of MultiphaseDynamic Imaging Findings with Pathologic Correlations Focusing on Nonfunctioning Tumors and Insulinomas. Journal of the Korean 67 0.2 Society of Radiology, **2010**, 63, 463 Intraabdominal complications secondary to ventriculoperitoneal shunts: CT findings and review of 66 86 5.4 the literature. American Journal of Roentgenology, 2009, 193, 1311-7 Hepatocellular carcinoma variants: radiologic-pathologic correlation. American Journal of 65 5.4 34 Roentgenology, **2009**, 193, W7-13 Added value of diffusion-weighted imaging in the MRI assessment of perilesional tumor recurrence 64 after chemoembolization of hepatocellular carcinomas. Journal of Magnetic Resonance Imaging, 5.6 53 2009, 30, 153-60 Ferucarbotran-enhanced T2-weighted magnetic resonance imaging: differentiation of hepatic 63 cavernous hemangiomas from malignant solid lesions. Abdominal Imaging, 2009, 34, 494-501 Small hypervascular hepatocellular carcinomas: value of "washout" on gadolinium-enhanced 62 dynamic MR imaging compared to superparamagnetic iron oxide-enhanced imaging. European 8 13 Radiology, 2009, 19, 2614-22 Hypervascular focus in the nonhypervascular nodule ("nodule-in-nodule") on dynamic computed 61 tomography: imaging evidence of aggressive progression in hepatocellular carcinoma. Journal of 6 2.2 Computer Assisted Tomography, 2009, 33, 131-5 Large (>or=2 cm) non-hypervascular nodules depicted on MRI in the cirrhotic liver: fate and 60 8 2.9 implications. Clinical Radiology, 2008, 63, 1121-30 Small hypervascular hepatocellular carcinoma: limited value of portal and delayed phases on 28 59 dynamic magnetic resonance imaging. Acta Radiologica, 2008, 49, 735-43

Ferucarbotran-Enhanced Hepatic MRI at 3T Unit: Quantitative and Qualitative Comparison of Fast

58	Hepatocellular carcinoma: progression in patients with hepatitis B versus C virus infection. <i>Radiology</i> , 2008 , 249, 1082; author reply 1082-3	20.5	
57	Small hypervascular enhancing lesions on arterial phase images of multiphase dynamic computed tomography in cirrhotic liver: fate and implications. <i>Journal of Computer Assisted Tomography</i> , 2008 , 32, 39-45	2.2	33
56	Choledochal cyst and anomalous pancreaticobiliary ductal union in adults: radiological spectrum and complications. <i>Journal of Computer Assisted Tomography</i> , 2008 , 32, 17-22	2.2	21
55	Radiological and clinical features of sarcomatoid hepatocellular carcinoma in 11 cases. <i>Journal of Computer Assisted Tomography</i> , 2008 , 32, 745-9	2.2	33
54	Prostate cancer: added value of subtraction dynamic imaging in 3T magnetic resonance imaging with a phased-array body coil. <i>Yonsei Medical Journal</i> , 2008 , 49, 765-74	3	14
53	Adrenocortical oncocytoma displaying intense activity on 18F-FDG-PET: a case report and a literature review. <i>Annals of Nuclear Medicine</i> , 2008 , 22, 821-4	2.5	20
52	Transient hepatic attenuation difference (THAD) following transcatheter arterial chemoembolization for hepatic malignancy: changes on serial CT examinations. <i>European Radiology</i> , 2008 , 18, 1596-603	8	10
51	Patient selection guidelines in MR-guided focused ultrasound surgery of uterine fibroids: a pictorial guide to relevant findings in screening pelvic MRI. <i>European Radiology</i> , 2008 , 18, 2997-3006	8	64
50	Value of manganese-enhanced T1- and T2-weighted MR cholangiography for differentiating cystic parenchymal lesions from cystic abnormalities which communicate with bile ducts. <i>Yonsei Medical Journal</i> , 2007 , 48, 1072-4	3	7
49	Gadobenate dimeglumine-enhanced MRI of intraductal papillary mucinous tumor of the bile ducts. Journal of Magnetic Resonance Imaging, 2007, 25, 625-7	5.6	9
48	Inflammatory myofibroblastic tumors in the liver: MRI of two immunohistochemically-verified cases. <i>Journal of Magnetic Resonance Imaging</i> , 2007 , 26, 418-21	5.6	8
47	Greater and lesser omenta: normal anatomy and pathologic processes. <i>Radiographics</i> , 2007 , 27, 707-20	5.4	92
46	Value of nonvisualized primary lesions of gastric cancer on preoperative MDCT. <i>American Journal of Roentgenology</i> , 2007 , 189, W315-9	5.4	21
45	MR cholangiography of accessory bile duct connected to the stomach. <i>American Journal of Roentgenology</i> , 2007 , 189, W344-7	5.4	9
44	Hepatic MRI using the double-echo chemical shift phase-selective gradient-echo technique. <i>American Journal of Roentgenology</i> , 2007 , 188, W49-56	5.4	9
43	Fat-containing nodules in the cirrhotic liver: chemical shift MRI features and clinical implications. <i>American Journal of Roentgenology</i> , 2007 , 188, 1009-16	5.4	35
42	Diagnostic accuracy of multidetector-row computed tomography for common bile duct calculi: is it necessary to add non-contrast-enhanced images to contrast-enhanced images?. <i>Journal of Computer Assisted Tomography</i> , 2007 , 31, 508-12	2.2	5
41	Biliary ductal involvement of hilar cholangiocarcinoma: multidetector computed tomography versus magnetic resonance cholangiography. <i>Journal of Computer Assisted Tomography</i> , 2007 , 31, 72-8	2.2	41

(2004-2007)

40	Isolated adrenal hemorrhage after minor blunt trauma. <i>American Journal of Emergency Medicine</i> , 2007 , 25, 984.e5-6	2.9	5
39	Hyperintense nodules on non-enhanced T1-weighted gradient-echo magnetic resonance imaging of cirrhotic liver: fate and clinical implications. <i>Journal of Magnetic Resonance Imaging</i> , 2006 , 24, 630-6	5.6	13
38	Hepatic metastases: perilesional enhancement on dynamic MRI. <i>American Journal of Roentgenology</i> , 2006 , 186, 1051-8	5.4	22
37	Pure epidural cavernous hemangioma of the cervical spine that presented with an acute sensory deficit caused by hemorrhage. <i>Yonsei Medical Journal</i> , 2006 , 47, 877-80	3	21
36	Preoperative staging accuracy of multidetector row computed tomography for extrahepatic bile duct carcinoma. <i>Journal of Computer Assisted Tomography</i> , 2006 , 30, 362-7	2.2	26
35	Diagnostic accuracy of multidetector row computed tomography in T- and N staging of gastric cancer with histopathologic correlation. <i>Journal of Computer Assisted Tomography</i> , 2006 , 30, 372-7	2.2	52
34	Advanced gastric carcinoma with signet ring cell carcinoma versus non-signet ring cell carcinoma: differentiation with multidetector CT. <i>Journal of Computer Assisted Tomography</i> , 2006 , 30, 880-4	2.2	22
33	Newly developed hepatocellular carcinoma (HCC) in chronic liver disease: MR imaging findings before the diagnosis of HCC. <i>Journal of Computer Assisted Tomography</i> , 2006 , 30, 765-71	2.2	16
32	Focal eosinophilic necrosis versus metastasis in the liver: the usefulness of two-phase dynamic CT. <i>American Journal of Roentgenology</i> , 2005 , 184, 1085-90	5.4	12
31	Eosinophilic hepatic necrosis: magnetic resonance imaging and computed tomography comparison. <i>Journal of Computer Assisted Tomography</i> , 2005 , 29, 765-71	2.2	16
30	Distinguishing hepatic metastases from hemangiomas: qualitative and quantitative diagnostic performance through dual echo respiratory-triggered fast spin echo magnetic resonance imaging. <i>Journal of Computer Assisted Tomography</i> , 2005 , 29, 571-9	2.2	10
29	Dynamic subtraction magnetic resonance imaging of cirrhotic liver: assessment of high signal intensity lesions on nonenhanced T1-weighted images. <i>Journal of Computer Assisted Tomography</i> , 2005 , 29, 51-8	2.2	40
28	Unusual cystic neoplasms in the pancreas: radiologic-pathologic correlation. <i>Journal of Computer Assisted Tomography</i> , 2005 , 29, 610-6	2.2	18
27	Capecitabine combined with gemcitabine (CapGem) as first-line treatment in patients with advanced/metastatic biliary tract carcinoma. <i>Cancer</i> , 2005 , 104, 2753-8	6.4	86
26	A Phase II study of capecitabine combined with gemcitabine in patients with advanced gallbladder carcinoma. <i>Yonsei Medical Journal</i> , 2005 , 46, 526-31	3	14
25	Abdominal sparganosis presenting as an abscess with fistulous communication to the bowel. <i>American Journal of Roentgenology</i> , 2005 , 185, 1084-5	5.4	13
24	Abdominal cocoon: preoperative diagnostic clues from radiologic imaging with pathologic correlation. <i>American Journal of Roentgenology</i> , 2004 , 182, 639-41	5.4	66
23	Early biliary complications of laparoscopic cholecystectomy: evaluation on T2-weighted MR cholangiography in conjunction with mangafodipir trisodium-enhanced 3D T1-weighted MR cholangiography. <i>American Journal of Roentgenology</i> , 2004 , 183, 1559-66	5.4	38

22	Stent-graft for TIPS in a hepatocellular carcinoma patient with main portal vein invasion. <i>American Journal of Roentgenology</i> , 2004 , 182, 1301-4	5.4	7
21	Secondary achalasia caused by peripheral cholangiocarcinoma in left lobe of the liver. <i>European Journal of Radiology Extra</i> , 2004 , 50, 89-92		
20	Acute cholecystitis at T2-weighted and manganese-enhanced T1-weighted MR cholangiography: preliminary study. <i>Radiology</i> , 2003 , 227, 580-4	20.5	45
19	Dynamic subtraction MR imaging of the liver: advantages and pitfalls. <i>American Journal of Roentgenology</i> , 2003 , 180, 1351-7	5.4	41
18	Predisposing factors of bile duct injury after transcatheter arterial chemoembolization (TACE) for hepatic malignancy. <i>CardioVascular and Interventional Radiology</i> , 2002 , 25, 270-4	2.7	59
17	Transcatheter arterial chemoembolization for hepatocellular carcinoma: anatomic and hemodynamic considerations in the hepatic artery and portal vein. <i>Radiographics</i> , 2002 , 22, 1077-91	5.4	49
16	Mucinous versus nonmucinous gastric carcinoma: differentiation with helical CT. <i>Radiology</i> , 2002 , 223, 540-6	20.5	30
15	Radiographic findings of primary B-cell lymphoma of the stomach: low-grade versus high-grade malignancy in relation to the mucosa-associated lymphoid tissue concept. <i>American Journal of Roentgenology</i> , 2002 , 179, 1297-304	5.4	31
14	Transient peritumoral enhancement during dynamic MRI of the liver: cavernous hemangioma versus hepatocellular carcinoma. <i>Journal of Computer Assisted Tomography</i> , 2002 , 26, 411-7	2.2	19
13	Radiologic findings of gastrointestinal tract involvement in hepatocellular carcinoma. <i>Journal of Computer Assisted Tomography</i> , 2002 , 26, 95-101	2.2	34
12	Magnetic resonance imaging of arterioportal shunts in the liver. <i>Topics in Magnetic Resonance Imaging</i> , 2002 , 13, 165-76	2.3	18
11	Urachal remnant diseases: spectrum of CT and US findings. <i>Radiographics</i> , 2001 , 21, 451-61	5.4	218
10	Recurrent pyogenic cholangitis: comparison between MR cholangiography and direct cholangiography. <i>Radiology</i> , 2001 , 220, 677-82	20.5	50
9	Bile duct injuries leading to portal vein obliteration after transcatheter arterial chemoembolization in the liver: CT findings and initial observations. <i>Radiology</i> , 2001 , 221, 429-36	20.5	33
8	Pneumoperitoneum caused by transhepatic air leak after metallic biliary stent placement. <i>CardioVascular and Interventional Radiology</i> , 2000 , 23, 482-4	2.7	4
7	Hepatic cavernous hemangioma in cirrhotic liver: imaging findings. <i>Korean Journal of Radiology</i> , 2000 , 1, 185-90	6.9	11
6	Percutaneous placement of self-expandable metallic biliary stents in malignant extrahepatic strictures: indications of transpapillary and suprapapillary methods. <i>Korean Journal of Radiology</i> , 2000 , 1, 65-72	6.9	14
5	Hepatic cavernous hemangioma: temporal peritumoral enhancement during multiphase dynamic MR imaging. <i>Radiology</i> , 2000 , 216, 692-7	20.5	61

LIST OF PUBLICATIONS

4	Nontumorous hepatic arterial-portal venous shunts: MR imaging findings. <i>Radiology</i> , 2000 , 217, 750-6	20.5	104
3	Retroperitoneal paragonimiasis: a case of ectopic paragonimiasis presenting as periureteral masses. <i>Journal of Computer Assisted Tomography</i> , 1999 , 23, 696-8	2.2	9
2	Comparison of multishot turbo spin echo and HASTE sequences for T2-weighted MRI of liver lesions. <i>Journal of Magnetic Resonance Imaging</i> , 1998 , 8, 1079-84	5.6	31
1	Intratumoral Blood Flow in Cavernous Hemangioma of the Liver: Radiologic-Pathologic Correlation. <i>Radiology</i> , 1998 , 208, 549-549	20.5	6