

Frans Pouwer

List of Publications by Citations

Source: <https://exaly.com/author-pdf/6306528/frans-pouwer-publications-by-citations.pdf>

Version: 2024-04-29

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

199
papers

7,229
citations

43
h-index

80
g-index

215
ext. papers

8,568
ext. citations

6
avg, IF

5.94
L-index

#	Paper	IF	Citations
199	Depression as a risk factor for the onset of type 2 diabetes mellitus. A meta-analysis. <i>Diabetologia</i> , 2006 , 49, 837-45	10.3	653
198	Type 2 diabetes mellitus as a risk factor for the onset of depression: a systematic review and meta-analysis. <i>Diabetologia</i> , 2010 , 53, 2480-6	10.3	470
197	Diabetes Attitudes, Wishes and Needs second study (DAWN2) cross-national benchmarking of diabetes-related psychosocial outcomes for people with diabetes. <i>Diabetic Medicine</i> , 2013 , 30, 767-77	3.5	405
196	Diabetes-related emotional distress in Dutch and U.S. diabetic patients: cross-cultural validity of the problem areas in diabetes scale. <i>Diabetes Care</i> , 2000 , 23, 1305-9	14.6	316
195	Validation of the WHO-5 Well-Being Index in adolescents with type 1 diabetes. <i>Diabetes Care</i> , 2007 , 30, 2003-6	14.6	172
194	Diabetes Attitudes, Wishes and Needs second study (DAWN2) cross-national benchmarking indicators for family members living with people with diabetes. <i>Diabetic Medicine</i> , 2013 , 30, 778-88	3.5	168
193	Psychological distress two years after diagnosis of breast cancer: frequency and prediction. <i>Patient Education and Counseling</i> , 2000 , 40, 209-17	3.1	166
192	Diabetes Attitudes Wishes and Needs 2 (DAWN2): a multinational, multi-stakeholder study of psychosocial issues in diabetes and person-centred diabetes care. <i>Diabetes Research and Clinical Practice</i> , 2013 , 99, 174-84	7.4	159
191	Prevalence of depression in individuals with impaired glucose metabolism or undiagnosed diabetes: a systematic review and meta-analysis of the European Depression in Diabetes (EDID) Research Consortium. <i>Diabetes Care</i> , 2011 , 34, 752-62	14.6	157
190	Web-based depression treatment for type 1 and type 2 diabetic patients: a randomized, controlled trial. <i>Diabetes Care</i> , 2011 , 34, 320-5	14.6	156
189	Rates and risks for co-morbid depression in patients with Type 2 diabetes mellitus: results from a community-based study. <i>Diabetologia</i> , 2003 , 46, 892-8	10.3	150
188	Association of Microvascular Dysfunction With Late-Life Depression: A Systematic Review and Meta-analysis. <i>JAMA Psychiatry</i> , 2017 , 74, 729-739	14.5	138
187	Does emotional stress cause type 2 diabetes mellitus? A review from the European Depression in Diabetes (EDID) Research Consortium. <i>Discovery Medicine</i> , 2010 , 9, 112-8	2.5	125
186	The confidence in diabetes self-care scale: psychometric properties of a new measure of diabetes-specific self-efficacy in Dutch and US patients with type 1 diabetes. <i>Diabetes Care</i> , 2003 , 26, 713-8	14.6	124
185	Psychometric and screening properties of the WHO-5 well-being index in adult outpatients with Type 1 or Type 2 diabetes mellitus. <i>Diabetic Medicine</i> , 2013 , 30, e63-9	3.5	119
184	Identification of barriers to insulin therapy and approaches to overcoming them. <i>Diabetes, Obesity and Metabolism</i> , 2018 , 20, 488-496	6.7	110
183	Prevalence of comorbid depression is high in out-patients with Type 1 or Type 2 diabetes mellitus. Results from three out-patient clinics in the Netherlands. <i>Diabetic Medicine</i> , 2010 , 27, 217-24	3.5	110

182	Diabetes-specific emotional distress mediates the association between depressive symptoms and glycaemic control in Type 1 and Type 2 diabetes. <i>Diabetic Medicine</i> , 2010 , 27, 798-803	3.5	108
181	The effects of a mindfulness-based intervention on emotional distress, quality of life, and HbA(1c) in outpatients with diabetes (DiaMind): a randomized controlled trial. <i>Diabetes Care</i> , 2013 , 36, 823-30	14.6	105
180	Monitoring of psychological well-being in outpatients with diabetes: effects on mood, HbA(1c), and the patient's evaluation of the quality of diabetes care: a randomized controlled trial. <i>Diabetes Care</i> , 2001 , 24, 1929-35	14.6	97
179	Screening tools used for measuring depression among people with Type 1 and Type 2 diabetes: a systematic review. <i>Diabetic Medicine</i> , 2012 , 29, 164-75	3.5	92
178	Effects of oxidative stress on fatty acid- and one-carbon-metabolism in psychiatric and cardiovascular disease comorbidity. <i>Acta Psychiatrica Scandinavica</i> , 2014 , 130, 163-80	6.5	83
177	Development and validation of the insulin treatment appraisal scale (ITAS) in patients with type 2 diabetes. <i>Health and Quality of Life Outcomes</i> , 2007 , 5, 69	3	83
176	The impact of comorbidity on Health-Related Quality of Life among cancer survivors: analyses of data from the PROFILES registry. <i>Journal of Cancer Survivorship</i> , 2013 , 7, 602-13	5.1	79
175	Associations of low grade inflammation and endothelial dysfunction with depression - The Maastricht Study. <i>Brain, Behavior, and Immunity</i> , 2016 , 56, 390-6	16.6	71
174	A comparison of the standard and the computerized versions of the Well-being Questionnaire (WBQ) and the Diabetes Treatment Satisfaction Questionnaire (DTSQ). <i>Quality of Life Research</i> , 1998 , 7, 33-8	3.7	70
173	Longitudinal associations between depression and diabetes complications: a systematic review and meta-analysis. <i>Diabetic Medicine</i> , 2019 , 36, 1562-1572	3.5	66
172	The 12-item well-being questionnaire. An evaluation of its validity and reliability in Dutch people with diabetes. <i>Diabetes Care</i> , 1999 , 22, 2004-10	14.6	66
171	Diabetes Fear of Injecting and Self-Testing Questionnaire: a psychometric evaluation. <i>Diabetes Care</i> , 2000 , 23, 765-9	14.6	65
170	Quality of life of children with type 1 diabetes: a systematic review. <i>Current Diabetes Reviews</i> , 2012 , 8, 434-43	2.7	62
169	The course of depressive symptoms in primary care patients with type 2 diabetes: results from the Diabetes, Depression, Type D Personality Zuidoost-Brabant (DiaDDZoB) Study. <i>Diabetologia</i> , 2012 , 55, 608-16	10.3	60
168	Effects of exercise training on quality of life, symptoms of depression, symptoms of anxiety and emotional well-being in type 2 diabetes mellitus: a systematic review. <i>Diabetologia</i> , 2013 , 56, 1210-25	10.3	60
167	Differential associations between depressive symptoms and glycaemic control in outpatients with diabetes. <i>Diabetic Medicine</i> , 2013 , 30, e115-22	3.5	56
166	Symptoms of depression and diabetes-specific emotional distress are associated with a negative appraisal of insulin therapy in insulin-naïve patients with Type 2 diabetes mellitus. A study from the European Depression in Diabetes [EDID] Research Consortium. <i>Diabetic Medicine</i> , 2009 , 26, 28-33	3.5	56
165	If it does not significantly change HbA1c levels why should we waste time on it? A plea for the prioritization of psychological well-being in people with diabetes. <i>Diabetic Medicine</i> , 2015 , 32, 155-63	3.5	52

164	Monitoring health related quality of life in adolescents with diabetes: a review of measures. <i>Archives of Disease in Childhood</i> , 2007 , 92, 434-9	2.2	52
163	The association between diabetes and an episode of depressive symptoms in the 2002 World Health Survey: an analysis of 231,797 individuals from 47 countries. <i>Diabetic Medicine</i> , 2013 , 30, e208-14 ^{3.5}	3.5	50
162	Limited effect of screening for depression with written feedback in outpatients with diabetes mellitus: a randomised controlled trial. <i>Diabetologia</i> , 2011 , 54, 741-8	10.3	49
161	Endothelial dysfunction is associated with a greater depressive symptom score in a general elderly population: the Hoorn Study. <i>Psychological Medicine</i> , 2014 , 44, 1403-16	6.9	48
160	Diabetes MILES--Australia (management and impact for long-term empowerment and success): methods and sample characteristics of a national survey of the psychological aspects of living with type 1 or type 2 diabetes in Australian adults. <i>BMC Public Health</i> , 2012 , 12, 120	4.1	46
159	Depressive symptoms are associated with physical inactivity in patients with type 2 diabetes. The DIAZOB Primary Care Diabetes study. <i>Family Practice</i> , 2009 , 26, 171-3	1.9	45
158	Associations between depressive symptoms and insulin resistance: the Hoorn Study. <i>Diabetologia</i> , 2006 , 49, 2874-7	10.3	45
157	Depression, anxiety and self-care behaviours of young adults with Type 2 diabetes: results from the International Diabetes Management and Impact for Long-term Empowerment and Success (MILES) Study. <i>Diabetic Medicine</i> , 2015 , 32, 133-40	3.5	44
156	The well-being questionnaire: evidence for a three-factor structure with 12 items (W-BQ12). <i>Psychological Medicine</i> , 2000 , 30, 455-62	6.9	43
155	Eicosapentaenoic acid as an add-on to antidepressant medication for co-morbid major depression in patients with diabetes mellitus: a randomized, double-blind placebo-controlled study. <i>Journal of Affective Disorders</i> , 2010 , 126, 282-6	6.6	42
154	Symptoms of depression in people with impaired glucose metabolism or Type 2 diabetes mellitus: The Hoorn Study. <i>Diabetic Medicine</i> , 2008 , 25, 843-9	3.5	40
153	Association between symptoms of depression and glycaemic control may be unstable across gender. <i>Diabetic Medicine</i> , 2001 , 18, 595-8	3.5	40
152	Diabetes-specific emotional distress in people with Type 2 diabetes: a comparison between primary and secondary care. <i>Diabetic Medicine</i> , 2014 , 31, 1252-9	3.5	39
151	Serious diabetes-specific emotional problems in patients with type 2 diabetes who have different levels of comorbid depression: a Polish study from the European Depression in Diabetes (EDID) Research Consortium. <i>European Psychiatry</i> , 2009 , 24, 425-30	6	39
150	Diabetes-related symptom distress in association with glucose metabolism and comorbidity: the Hoorn Study. <i>Diabetes Care</i> , 2008 , 31, 2268-70	14.6	38
149	Diabetes distress is more strongly associated with HbA1c than depressive symptoms in adolescents with type 1 diabetes: Results from Diabetes MILES Youth-Australia. <i>Pediatric Diabetes</i> , 2018 , 19, 840-847 ^{3.6}	3.6	37
148	Suboptimal glycemic control in type 2 diabetes: a key role for anhedonia?. <i>Journal of Psychiatric Research</i> , 2012 , 46, 549-54	5.2	37
147	Association of coexisting diabetes and depression with mortality after myocardial infarction. <i>Diabetes Care</i> , 2012 , 35, 503-9	14.6	35

146	Psychological and personality factors in type 2 diabetes mellitus, presenting the rationale and exploratory results from The Maastricht Study, a population-based cohort study. <i>BMC Psychiatry</i> , 2016 , 16, 17	4.2	34
145	Effectiveness of a stepped care intervention for anxiety and depression in people with diabetes, asthma or COPD in primary care: A randomized controlled trial. <i>Journal of Affective Disorders</i> , 2015 , 184, 269-76	6.6	33
144	Type D personality, suboptimal health behaviors and emotional distress in adults with diabetes: results from Diabetes MILES-The Netherlands. <i>Diabetes Research and Clinical Practice</i> , 2015 , 108, 94-105	7.4	33
143	The association between glucose-lowering drug use and mortality among breast cancer patients with type 2 diabetes. <i>Breast Cancer Research and Treatment</i> , 2015 , 150, 427-37	4.4	33
142	Web-based cognitive behavioural therapy (W-CBT) for diabetes patients with co-morbid depression: design of a randomised controlled trial. <i>BMC Psychiatry</i> , 2008 , 8, 9	4.2	33
141	Association of Type D personality with increased vulnerability to depression: Is there a role for inflammation or endothelial dysfunction? - The Maastricht Study. <i>Journal of Affective Disorders</i> , 2016 , 189, 118-25	6.6	32
140	Gender differences in disordered eating behaviors and body dissatisfaction among adolescents with type 1 diabetes: Results from diabetes MILES youth-Australia. <i>International Journal of Eating Disorders</i> , 2017 , 50, 1183-1193	6.3	31
139	Toward defining a cutoff score for elevated fear of hypoglycemia on the hypoglycemia fear survey worry subscale in patients with type 2 diabetes. <i>Diabetes Care</i> , 2014 , 37, 102-8	14.6	31
138	Living with type 1 diabetes is challenging for Zambian adolescents: qualitative data on stress, coping with stress and quality of care and life. <i>BMC Endocrine Disorders</i> , 2015 , 15, 20	3.3	30
137	What is the best measure for assessing diabetes distress? A comparison of the Problem Areas in Diabetes and Diabetes Distress Scale: results from Diabetes MILES-Australia. <i>Journal of Health Psychology</i> , 2018 , 23, 667-680	3.1	29
136	Managing hypoglycemia in diabetes may be more fear management than glucose management: a practical guide for diabetes care providers. <i>Current Diabetes Reviews</i> , 2014 , 10, 364-70	2.7	29
135	Continuous intraperitoneal insulin infusion in patients with brittle diabetes: favourable effects on glycaemic control and hospital stay. <i>Diabetic Medicine</i> , 2002 , 19, 496-501	3.5	28
134	Negative appraisals of insulin therapy are common among adults with Type 2 diabetes using insulin: Results from Diabetes MILES - Australia cross-sectional survey. <i>Diabetic Medicine</i> , 2015 , 32, 1297-303	3.5	27
133	Initiation of insulin glargine in patients with Type 2 diabetes in suboptimal glycaemic control positively impacts health-related quality of life. A prospective cohort study in primary care. <i>Diabetic Medicine</i> , 2011 , 28, 1096-102	3.5	26
132	The Use of Mobile Applications Among Adults with Type 1 and Type 2 Diabetes: Results from the Second MILES-Australia (MILES-2) Study. <i>Diabetes Technology and Therapeutics</i> , 2017 , 19, 730-738	8.1	25
131	Severely obese people with diabetes experience impaired emotional well-being associated with socioeconomic disadvantage: results from diabetes MILES - Australia. <i>Diabetes Research and Clinical Practice</i> , 2013 , 101, 131-40	7.4	24
130	Fear of hypoglycaemia in adults with Type 1 diabetes: results from Diabetes MILES - The Netherlands. <i>Diabetic Medicine</i> , 2015 , 32, 1289-96	3.5	24
129	Associations between vascular co-morbidities and depression in insulin-naïve diabetes patients: the DIAZOB Primary Care Diabetes study. <i>Diabetologia</i> , 2009 , 52, 2056-63	10.3	23

128	PatientsPEvaluation of the Quality of Diabetes Care (PEQD): development and validation of a new instrument. <i>Quality and Safety in Health Care</i> , 2002 , 11, 131-6		23
127	The impact of having both cancer and diabetes on patient-reported outcomes: a systematic review and directions for future research. <i>Journal of Cancer Survivorship</i> , 2016 , 10, 406-15	5.1	22
126	The relationship between parenting stress and parent-child interaction with health outcomes in the youngest patients with type 1 diabetes (0-7 years). <i>European Journal of Pediatrics</i> , 2016 , 175, 329-38	4.1	21
125	The longitudinal association between glycaemic control and health-related quality of life following insulin therapy optimisation in type 2 diabetes patients. A prospective observational study in secondary care. <i>Quality of Life Research</i> , 2012 , 21, 1359-65	3.7	21
124	Advanced Glycation End Product (AGE) Accumulation in the Skin is Associated with Depression: The Maastricht Study. <i>Depression and Anxiety</i> , 2017 , 34, 59-67	8.4	19
123	Stressful life events and incident metabolic syndrome: the Hoorn study. <i>Stress</i> , 2015 , 18, 507-13	3	19
122	Diabetes MILES Youth-Australia: methods and sample characteristics of a national survey of the psychological aspects of living with type 1 diabetes in Australian youth and their parents. <i>BMC Psychology</i> , 2016 , 4, 42	2.8	19
121	Fat food for a bad mood. Could we treat and prevent depression in Type 2 diabetes by means of omega-3 polyunsaturated fatty acids? A review of the evidence. <i>Diabetic Medicine</i> , 2005 , 22, 1465-75	3.5	19
120	Ceiling effect reduces the validity of the Diabetes Treatment Satisfaction Questionnaire. <i>Diabetes Care</i> , 1998 , 21, 2039	14.6	19
119	Psychometric properties of the Exercise Self-efficacy Scale in Dutch Primary care patients with type 2 diabetes mellitus. <i>International Journal of Behavioral Medicine</i> , 2014 , 21, 394-401	2.6	18
118	What Do Adults with Type 2 Diabetes Want from the "Perfect" App? Results from the Second Diabetes MILES: Australia (MILES-2) Study. <i>Diabetes Technology and Therapeutics</i> , 2019 , 21, 393-399	8.1	17
117	Compromised quality of life in patients with both Type 1 diabetes mellitus and coeliac disease. <i>Diabetic Medicine</i> , 2013 , 30, 835-9	3.5	17
116	How 25 years of psychosocial research has contributed to a better understanding of the links between depression and diabetes. <i>Diabetic Medicine</i> , 2020 , 37, 383-392	3.5	17
115	Cohort profiles of the cross-sectional and prospective participant groups in the second Diabetes MILES-Australia (MILES-2) study. <i>BMJ Open</i> , 2017 , 7, e012926	3	16
114	Modeling Interactions Between Latent Variables in Research on Type D Personality: A Monte Carlo Simulation and Clinical Study of Depression and Anxiety. <i>Multivariate Behavioral Research</i> , 2019 , 54, 637-685	2.3	16
113	Cut Points for Identifying Clinically Significant Diabetes Distress in Adolescents With Type 1 Diabetes Using the PAID-T: Results From Diabetes MILES Youth-Australia. <i>Diabetes Care</i> , 2017 , 40, 1462-1468	14.6	16
112	Back to the future: 25 years of R Guidelines for encouraging psychological well-beingAmong people affected by diabetes. <i>Diabetic Medicine</i> , 2020 , 37, 1225-1229	3.5	16
111	Suicidal ideation reported by adults with Type 1 or Type 2 diabetes: results from Diabetes MILES-Australia. <i>Diabetic Medicine</i> , 2016 , 33, 1582-1589	3.5	16

110	Strengths, Risk Factors, and Resilient Outcomes in Adolescents With Type 1 Diabetes: Results From Diabetes MILES Youth-Australia. <i>Diabetes Care</i> , 2017 , 40, 849-855	14.6	15
109	Severe hypoglycemia, impaired awareness of hypoglycemia, and self-monitoring in adults with type 1 diabetes: Results from Diabetes MILES-Australia. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 577-582	3.2	15
108	Diabetes stigma is associated with negative treatment appraisals among adults with insulin-treated Type 2 diabetes: results from the second Diabetes MILES - Australia (MILES-2) survey. <i>Diabetic Medicine</i> , 2018 , 35, 658-662	3.5	15
107	Do you wish my health professionals understood that it's not just all about your HbA1c? Qualitative responses from the second Diabetes MILES - Australia (MILES-2) study. <i>Diabetic Medicine</i> , 2020 , 37, 971-981	3.5	15
106	Paediatric parenting stress in fathers and mothers of young children with Type 1 diabetes: a longitudinal study. <i>Diabetic Medicine</i> , 2017 , 34, 821-827	3.5	14
105	Further investigation of the psychometric properties of the Insulin Treatment Appraisal Scale among insulin-using and non-insulin-using adults with type 2 diabetes: results from Diabetes MILES-Australia. <i>Health and Quality of Life Outcomes</i> , 2014 , 12, 87	3	14
104	The impact of diabetes on neuropathic symptoms and receipt of chemotherapy among colorectal cancer patients: results from the PROFILES registry. <i>Journal of Cancer Survivorship</i> , 2015 , 9, 523-31	5.1	13
103	Explaining psychological insulin resistance in adults with non-insulin-treated type 2 diabetes: The roles of diabetes distress and current medication concerns. Results from Diabetes MILES-Australia. <i>Primary Care Diabetes</i> , 2016 , 10, 75-82	2.4	13
102	The associations between diabetes distress and self-efficacy, medication adherence, self-care activities and disease control depend on the way diabetes distress is measured: Comparing the DDS-17, DDS-2 and the PAID-5. <i>Diabetes Research and Clinical Practice</i> , 2018 , 142, 74-84	7.4	13
101	Assessing the perceived impact of diabetes on quality of life: Psychometric validation of the DAWN2 Impact of Diabetes Profile in the second Diabetes MILES - Australia (MILES-2) survey. <i>Diabetes Research and Clinical Practice</i> , 2019 , 150, 253-263	7.4	12
100	Anxiety, depression and timing of insulin treatment among people with type 2 diabetes: Nine-year follow-up of the Nord-Trøndelag Health Study, Norway. <i>Journal of Psychosomatic Research</i> , 2015 , 79, 309-15	4.1	12
99	Reducing the burden of hypoglycaemia in people with diabetes through increased understanding: design of the Hypoglycaemia REdefining SOLUTIONS for better liVEs (Hypo-RESOLVE) project. <i>Diabetic Medicine</i> , 2020 , 37, 1066-1073	3.5	12
98	Mindfulness and fear of hypoglycaemia in parents of children with Type 1 diabetes: results from Diabetes MILES Youth - The Netherlands. <i>Diabetic Medicine</i> , 2018 , 35, 650-657	3.5	12
97	Prospectively measured lifestyle factors and BMI explain differences in health-related quality of life between colorectal cancer patients with and without comorbid diabetes. <i>Supportive Care in Cancer</i> , 2016 , 24, 2591-601	3.9	12
96	The individual and combined effect of colorectal cancer and diabetes on health-related quality of life and sexual functioning: results from the PROFILES registry. <i>Supportive Care in Cancer</i> , 2014 , 22, 3071-3079	3.9	12
95	Severe obesity and diabetes self-care attitudes, behaviours and burden: implications for weight management from a matched case-controlled study. Results from Diabetes MILES-Australia. <i>Diabetic Medicine</i> , 2014 , 31, 232-40	3.5	12
94	Associations between economic hardship and markers of self-management in adults with type 2 diabetes: results from Diabetes MILES - Australia. <i>Australian and New Zealand Journal of Public Health</i> , 2014 , 38, 466-72	2.3	12
93	Psychosocial research in the diabetic foot: Are we making progress?. <i>Diabetes/Metabolism Research and Reviews</i> , 2020 , 36 Suppl 1, e3257	7.5	12

92	The Use of Mobile Applications Among Adolescents with Type 1 Diabetes: Results from Diabetes MILES Youth-Australia. <i>Diabetes Technology and Therapeutics</i> , 2016 , 18, 813-819	8.1	11
91	Does the shortage of diabetes specialists in regional and rural Australia matter? Results from Diabetes MILES--Australia. <i>Diabetes Research and Clinical Practice</i> , 2013 , 100, 222-9	7.4	11
90	Ethnic aspects of emotional distress in patients with diabetes--the Amsterdam Health Monitor Study. <i>Diabetic Medicine</i> , 2013 , 30, e25-31	3.5	11
89	No evidence for increased self-reported cognitive failure in Type 1 and Type 2 diabetes: a cross-sectional study. <i>Diabetic Medicine</i> , 2007 , 24, 735-40	3.5	11
88	Prevalence of Insomnia (Symptoms) in T2D and Association With Metabolic Parameters and Glycemic Control: Meta-Analysis. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020 , 105,	5.6	11
87	Diabetes Complications and Depressive Symptoms: Prospective Results From the Montreal Diabetes Health and Well-Being Study. <i>Psychosomatic Medicine</i> , 2017 , 79, 603-612	3.7	10
86	The Association of Mindful Parenting with Glycemic Control and Quality of Life in Adolescents with Type 1 Diabetes: Results from Diabetes MILES-The Netherlands. <i>Mindfulness</i> , 2016 , 7, 1227-1237	2.9	10
85	Is Self-Compassion Related to Behavioural, Clinical and Emotional Outcomes in Adults with Diabetes? Results from the Second Diabetes MILES Australia (MILES-2) Study. <i>Mindfulness</i> , 2019 , 10, 1222-1231	2.9	10
84	Factor structure of the Disability and Impact Profile in patients with multiple sclerosis. <i>Quality of Life Research</i> , 1999 , 8, 141-50	3.7	9
83	Subjective Wellbeing Among Adults with Diabetes: Results from Diabetes MILES Australia. <i>Journal of Happiness Studies</i> , 2016 , 17, 1205-1217	3.7	8
82	The Epidemiology of Depression and Diabetes 2010 , 1-27		8
81	Anxiety and Depressive Symptoms as Predictors of All-Cause Mortality among People with Insulin-Dependent Type 2 Diabetes: 17-Year Follow-Up of the Second Nord-Trøndelag Health Survey (HUNT2), Norway. <i>PLoS ONE</i> , 2016 , 11, e0160861	3.7	8
80	Assessing Psychological Insulin Resistance in Type 2 Diabetes: a Critical Comparison of Measures. <i>Current Diabetes Reports</i> , 2017 , 17, 46	5.6	7
79	Prevalence and correlates of diabetes distress, perceived stress and depressive symptoms among adults with early-onset Type 2 diabetes: cross-sectional survey results from the Danish DD2 study. <i>Diabetic Medicine</i> , 2020 , 37, 1679-1687	3.5	7
78	Depressive symptoms, insulin sensitivity and insulin secretion in the RISC cohort study. <i>Diabetes and Metabolism</i> , 2013 , 39, 42-9	5.4	7
77	Quality of the parent-child interaction in young children with type 1 diabetes mellitus: study protocol. <i>BMC Pediatrics</i> , 2011 , 11, 28	2.6	7
76	Psychosocial Moderators of the Impact of Diabetes Stigma: Results From the Second Diabetes MILES - Australia (MILES-2) Study. <i>Diabetes Care</i> , 2020 , 43, 2651-2659	14.6	7
75	Sustainable improvement of HbA and satisfaction with diabetes care after adding telemedicine in patients on adaptable insulin regimens: Results of the TeleDiabetes randomized controlled trial. <i>Health Informatics Journal</i> , 2020 , 26, 628-641	3	7

74	Intraperitoneal versus subcutaneous insulin therapy in the treatment of type 1 diabetes mellitus. <i>Netherlands Journal of Medicine</i> , 2015 , 73, 399-409	0.5	7
73	Diabetes mellitus and comorbid depression in Zambia. <i>Diabetic Medicine</i> , 2015 , 32, 814-8	3.5	6
72	Effect of vitamin D supplementation on health status in non-vitamin D deficient people with type 2 diabetes mellitus. <i>Endocrine Connections</i> , 2016 , 5, 61-69	3.5	6
71	Does self-efficacy mediate the cross-sectional relationship between perceived quality of health care and self-management of diabetes? Results from Diabetes MILES - Australia. <i>Psychology and Health</i> , 2016 , 31, 592-604	2.9	6
70	Emotional Well-Being Factors Associated with Sexual Dysfunction in Adults with Type 1 or Type 2 Diabetes: Results from Diabetes MILES Australia. <i>International Journal of Sexual Health</i> , 2018 , 30, 237-249	2.9	6
69	Type D personality and social relations in adults with diabetes: results from diabetes MILES - The Netherlands. <i>Psychology and Health</i> , 2018 , 33, 1456-1471	2.9	6
68	Psychological correlates of disordered eating in youth with type 1 diabetes: Results from diabetes MILES Youth-Australia. <i>Pediatric Diabetes</i> , 2020 , 21, 664-672	3.6	5
67	Review of research grant allocation to psychosocial studies in diabetes research. <i>Diabetic Medicine</i> , 2016 , 33, 1673-1676	3.5	5
66	Vitamin D status and health-related quality of life in patients with Type 2 diabetes. <i>Diabetic Medicine</i> , 2016 , 33, 300-6	3.5	5
65	Social support and self-care outcomes in adults with diabetes: The mediating effects of self-efficacy and diabetes distress. Results of the second diabetes MILES - Australia (MILES-2) study. <i>Diabetes Research and Clinical Practice</i> , 2020 , 166, 108314	7.4	5
64	The role of parental support for emerging adults with type 1 diabetes: A scoping review. <i>Pediatric Diabetes</i> , 2020 , 21, 995-1030	3.6	5
63	Associations Between Physical Activity and Depressive Symptoms by Weight Status Among Adults With Type 2 Diabetes: Results From Diabetes MILES-Australia. <i>Journal of Physical Activity and Health</i> , 2017 , 14, 195-202	2.5	4
62	Qualitative observation instrument to measure the quality of parent-child interactions in young children with type 1 diabetes mellitus. <i>BMC Pediatrics</i> , 2014 , 14, 145	2.6	4
61	Low vitamin D levels are not a contributing factor to higher prevalence of depressive symptoms in people with Type 2 diabetes mellitus: the Hoorn study. <i>Diabetic Medicine</i> , 2017 , 34, 577-581	3.5	4
60	Suicidal ideation or non-suicidal self-harm? A mismatch between the DSM-IV criterion and PHQ-9 item nine. <i>Diabetes Research and Clinical Practice</i> , 2015 , 108, e5-6	7.4	4
59	Prevalence and course of mood and anxiety disorders, and correlates of symptom severity in adolescents with type 1 diabetes: Results from diabetes LEAP. <i>Pediatric Diabetes</i> , 2021 , 22, 638-648	3.6	4
58	The impact of hypoglycaemia on quality of life outcomes among adults with type 1 diabetes: A systematic review. <i>Diabetes Research and Clinical Practice</i> , 2021 , 174, 108752	7.4	4
57	The suitability of patient-reported outcome measures used to assess the impact of hypoglycaemia on quality of life in people with diabetes: a systematic review using COSMIN methods. <i>Diabetologia</i> , 2021 , 64, 1213-1225	10.3	4

56	The bidirectional longitudinal association between depressive symptoms and HbA : A systematic review and meta-analysis. <i>Diabetic Medicine</i> , 2021 , e14671	3.5	4
55	Psychiatric disorders as risk factors for the development of type 2 diabetes mellitus: an umbrella review protocol. <i>BMJ Open</i> , 2019 , 9, e024981	3	3
54	Impact of cancer on use of glucose-lowering drug treatment in individuals with diabetes: potential mechanisms. <i>Diabetologia</i> , 2015 , 58, 1378-9	10.3	3
53	Depressive Symptoms Are Negatively Associated with Glucose Testing and Eating Meals on Time among Individuals with Diabetes in Zambia. <i>Diabetes and Metabolism Journal</i> , 2017 , 41, 440-448	5	3
52	Comment on Young-Hyman et al. Psychosocial Care for People With Diabetes: A Position Statement of the American Diabetes Association. <i>Diabetes Care</i> 2016;39:2126-2140. <i>Diabetes Care</i> , 2018 , 41, e31-e32	14.6	3
51	Anxiety is common and costly in T2DM - why psychology matters. <i>Nature Reviews Endocrinology</i> , 2019 , 15, 567-568	15.2	3
50	Reply to comment on: Knol MJ, Twisk JWR, Beekman ATF, Heine RJ, Snoek FJ, Pouwer F (2006) Depression as a risk factor for the onset of type 2 diabetes mellitus. A meta-analysis. <i>Diabetologia</i> 49:837-845. <i>Diabetologia</i> , 2006 , 49, 2799-2800	10.3	3
49	Depressive and anxiety symptoms and following of the Dutch Dietary Guidelines 2015 in adults with diabetes: Results from Diabetes MILES-The Netherlands. <i>Journal of Psychosomatic Research</i> , 2020 , 135, 110160	4.1	3
48	Positive facilitators of diabetes management in emerging adults with type 1 diabetes-A qualitative analysis of blogs. <i>Endocrinology, Diabetes and Metabolism</i> , 2020 , 3, e00161	2.7	3
47	Prospective memory slips are associated with forgetting to take glucose-lowering therapies among adults with diabetes: results from the second Diabetes MILES - Australia (MILES-2) survey. <i>Diabetic Medicine</i> , 2019 , 36, 569-577	3.5	3
46	Measuring and Assessing Depression in People with Diabetes: Implications for Clinical Practice 2013 , 199-209		2
45	Online cognitieve gedragstherapie voor diabetespatiënten met depressieve klachten: een gerandomiseerd gecontroleerd onderzoek. <i>Psychologie and Gezondheid</i> , 2009 , 37, 222-228		2
44	Depressive symptoms, insulin resistance, and risk of diabetes in women at midlife: response to Everson-Rose et al. <i>Diabetes Care</i> , 2005 , 28, 1265-6; author reply 1266	14.6	2
43	Depression Is Associated With Progression of Diabetic Nephropathy in Type 1 Diabetes. <i>Diabetes Care</i> , 2021 , 44, 174-180	14.6	2
42	The division and transfer of care responsibilities in paediatric type 1 diabetes: A qualitative study on parental perspectives. <i>Journal of Advanced Nursing</i> , 2021 , 77, 1968-1979	3.1	2
41	Prevalence of and risk factors for sexual dysfunctions in adults with type 1 or type 2 diabetes: Results from Diabetes MILES - Flanders. <i>Diabetic Medicine</i> , 2021 , e14676	3.5	2
40	Fatigue in type 1 diabetes: an understudied problem. <i>Lancet Diabetes and Endocrinology</i> , 2017 , 5, 409-410	18.1	1
39	The prevalence of type 2 diabetes in people with psychiatric disorders: an umbrella review protocol. <i>Systematic Reviews</i> , 2020 , 9, 101	3	1

38	Correlates of fear of hypoglycemia among patients with type 1 and 2 diabetes mellitus in outpatient hospitals in Zambia. <i>International Journal of Diabetes in Developing Countries</i> , 2020 , 40, 619-626	6.8	1
37	Diabetes, Depression, and Cardiovascular Risk 2016 , 831-847		1
36	More work is needed to better understand diabetes distress as a predictor of all-cause mortality in type 2 diabetes. <i>Diabetologia</i> , 2018 , 61, 2247-2248	10.3	1
35	Study protocol of Diabetes LEAP: a longitudinal study examining emotional problems in adolescents with type 1 diabetes and their parents/caregivers. <i>BMC Pediatrics</i> , 2019 , 19, 377	2.6	1
34	Comment on Young-Hyman et al. Psychosocial Care for People With Diabetes: A Position Statement of the American Diabetes Association. <i>Diabetes Care</i> 2016;39:2126-2140. <i>Diabetes Care</i> , 2017 , 40, e127-e128	14.6	1
33	Psychological and pharmacological interventions for depression in patients with diabetes mellitus: an abridged Cochrane review: some critical comments. <i>Diabetic Medicine</i> , 2015 , 32, 839	3.5	1
32	Comment on Schmitz et al. Recurrent subthreshold depression in type 2 diabetes: an important risk factor for poor health outcomes. <i>Diabetes care</i> 2014;37:970-978. <i>Diabetes Care</i> , 2014 , 37, e145	14.6	1
31	Personality characteristics in adolescence predict long-term changes in body fatness. <i>Diabetes Research and Clinical Practice</i> , 2008 , 79, e10-3	7.4	1
30	Omissions impact measurements of patient-reported outcomes. <i>Quality of Life Research</i> , 2008 , 17, 963-4; author reply 965-6	3.7	1
29	Prevalence of type 2 diabetes in psychiatric disorders: an umbrella review with meta-analysis of 245 observational studies from 32 systematic reviews. <i>Diabetologia</i> , 2021 , 1	10.3	1
28	The impact of hypoglycemia on quality of life and related outcomes in children and adolescents with type 1 diabetes: A systematic review. <i>PLoS ONE</i> , 2021 , 16, e0260896	3.7	1
27	Depression and anxiety in adolescents with type 1 diabetes and their parents. <i>Pediatric Research</i> , 2021 ,	3.2	1
26	Symptoms of depression and anxiety in adults with type 1 diabetes: Associations with self-care behaviour, glycaemia and incident complications over four years - Results from diabetes MILES-Australia. <i>Journal of Affective Disorders</i> , 2021 , 282, 803-811	6.6	1
25	Hypoglycaemia and diabetes-specific quality of life in adolescents with type 1 diabetes. <i>Diabetic Medicine</i> , 2021 , 38, e14565	3.5	1
24	Does nocturnal hypoglycaemia really improve quality of life?. <i>Diabetologia</i> , 2021 , 64, 1893-1894	10.3	1
23	Data on diabetes-specific distress are needed to improve the quality of diabetes care. <i>Lancet, The</i> , 2021 , 397, 2149	4.0	1
22	Psychiatric disorders as risk factors for type 2 diabetes: An umbrella review of systematic reviews with and without meta-analyses. <i>Diabetes Research and Clinical Practice</i> , 2021 , 176, 108855	7.4	1
21	Comment on the consensus report on the management of hyperglycaemia in Type 2 diabetes by the American Diabetes Association and the European Association for the Study of Diabetes. <i>Diabetic Medicine</i> , 2019 , 36, 911-912	3.5	1

20	The impact of hypoglycaemia on the quality of life of family members of adults with type 1 or type 2 diabetes: A qualitative systematic review. <i>Diabetic Medicine</i> , 2021 , 38, e14666	3.5	1
19	‘Never again will I be carefree’ a qualitative study of the impact of hypoglycemia on quality of life among adults with type 1 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2021 , 9,	4.5	1
18	Self-reported access to and quality of healthcare for diabetes: do the severely obese experience equal access?. <i>Australian Family Physician</i> , 2014 , 43, 552-6		1
17	Investigating the day-to-day impact of hypoglycaemia in adults with type 1 or type 2 diabetes: design and validation protocol of the Hypo-METRICS application.. <i>BMJ Open</i> , 2022 , 12, e051651	3	0
16	Changes in quality of life following hypoglycaemia in adults with type 2 diabetes: A systematic review of longitudinal studies. <i>Diabetic Medicine</i> , 2021 , e14706	3.5	0
15	How to identify clinically significant diabetes distress using the Problem Areas in Diabetes (PAID) scale in adults with diabetes treated in primary or secondary care? Evidence for new cut points based on latent class analyses.. <i>BMJ Open</i> , 2022 , 12, e056304	3	0
14	Overeating, binge eating, quality of life, emotional difficulties, and HbA in adolescents with type 1 diabetes: A Danish national survey. <i>Diabetes Research and Clinical Practice</i> , 2021 , 182, 109150	7.4	0
13	Diabetes in Sub-Saharan African Children: Risks, Care, and Challenges 2017 , 157-172		
12	Response to Kawada. Depressive symptoms and HbA1c in patients with Type 1 and Type 2 diabetes. <i>Diabetic Medicine</i> , 2014 , 31, 760-1	3.5	
11	PS12 - 7. The quality of parent-child interaction and disease-related parenting stress is associated with HbA1c and quality of life in young children with type 1 diabetes.. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2013 , 11, 180-180	0	
10	Screening for type 2 diabetes does not reduce mortality over 10 years. <i>Evidence-Based Medicine</i> , 2013 , 18, e44		
9	PS11 - 59. Development of a new observational rating scale to measure the quality of parent-child interactions in young children with type 1 diabetes mellitus. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2011 , 9, 131-131	0	
8	Psychosociale interventies bij mensen met diabetes. <i>Psychologie and Gezondheid</i> , 2008 , 36, 209-220		
7	--to: M. Koopmanschap: Coping with Type 2 diabetes: the patient’s perspective. <i>Diabetologia</i> 45:S18-S22. <i>Diabetologia</i> , 2003 , 46, 302-3	10.3	
6	Unmet support needs relating to hypoglycaemia among adults with type 1 diabetes: Results of a multi-country web-based qualitative study. <i>Diabetic Medicine</i> , 2021 , e14727	3.5	
5	Diabetes, Depression, and Cardiovascular Risk 2015 , 1-17		
4	Psychological Factors and Diabetes Mellitus 2016 , 819-839		
3	Comment on van Duinkerken et al. Biopsychosocial factors associated with a current depressive episode in diabetes: The ELSA-Brasil study. <i>Diabetic Medicine</i> , 2021 , 38, e14418	3.5	

- | | | |
|---|---|-----|
| 2 | Facilitating participation in cardiovascular preventive initiatives among people with diabetes: a qualitative study. <i>BMC Public Health</i> , 2021 , 21, 203 | 4.1 |
| 1 | Mindfulness in Relation to Diet Quality in Adults with Type 1 and Type 2 Diabetes: Results from Diabetes MILES-The Netherlands. <i>Mindfulness</i> ,1 | 2.9 |