Frans Pouwer

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/6306528/frans-pouwer-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

 199
 7,229
 43
 80

 papers
 h-index
 g-index

 215
 8,568
 6
 5.94

 ext. papers
 ext. citations
 avg, IF
 L-index

#	Paper	IF	Citations
199	Investigating the day-to-day impact of hypoglycaemia in adults with type 1 or type 2 diabetes: design and validation protocol of the Hypo-METRICS application <i>BMJ Open</i> , 2022 , 12, e051651	3	O
198	How to identify clinically significant diabetes distress using the Problem Areas in Diabetes (PAID) scale in adults with diabetes treated in primary or secondary care? Evidence for new cut points based on latent class analyses <i>BMJ Open</i> , 2022 , 12, e056304	3	О
197	Prevalence of type 2 diabetes in psychiatric disorders: an umbrella review with meta-analysis of 245 observational studies from 32 systematic reviews. <i>Diabetologia</i> , 2021 , 1	10.3	1
196	The impact of hypoglycemia on quality of life and related outcomes in children and adolescents with type 1 diabetes: A systematic review. <i>PLoS ONE</i> , 2021 , 16, e0260896	3.7	1
195	Unmet support needs relating to hypoglycaemia among adults with type 1 diabetes: Results of a multi-country web-based qualitative study. <i>Diabetic Medicine</i> , 2021 , e14727	3.5	
194	Depression and anxiety in adolescents with type 1 diabetes and their parents. <i>Pediatric Research</i> , 2021 ,	3.2	1
193	Prevalence and course of mood and anxiety disorders, and correlates of symptom severity in adolescents with type 1 diabetes: Results from diabetes LEAP. <i>Pediatric Diabetes</i> , 2021 , 22, 638-648	3.6	4
192	Symptoms of depression and anxiety in adults with type 1 diabetes: Associations with self-care behaviour, glycaemia and incident complications over four years - Results from diabetes MILES-Australia. <i>Journal of Affective Disorders</i> , 2021 , 282, 803-811	6.6	1
191	The impact of hypoglycaemia on quality of life outcomes among adults with type 1 diabetes: A systematic review. <i>Diabetes Research and Clinical Practice</i> , 2021 , 174, 108752	7.4	4
190	Hypoglycaemia and diabetes-specific quality of life in adolescents with type 1 diabetes. <i>Diabetic Medicine</i> , 2021 , 38, e14565	3.5	1
189	Does nocturnal hypoglycaemia really improve quality of life?. <i>Diabetologia</i> , 2021 , 64, 1893-1894	10.3	1
188	Data on diabetes-specific distress are needed to improve the quality of diabetes care. <i>Lancet, The</i> , 2021 , 397, 2149	40	1
187	Psychiatric disorders as risk factors for type 2 diabetes: An umbrella review of systematic reviews with and without meta-analyses. <i>Diabetes Research and Clinical Practice</i> , 2021 , 176, 108855	7.4	1
186	Depression Is Associated With Progression of Diabetic Nephropathy in Type 1 Diabetes. <i>Diabetes Care</i> , 2021 , 44, 174-180	14.6	2
185	Comment on van Duinkerken et al. Biopsychosocial factors associated with a current depressive episode in diabetes: The ELSA-Brasil study. <i>Diabetic Medicine</i> , 2021 , 38, e14418	3.5	
184	Facilitating participation in cardiovascular preventive initiatives among people with diabetes: a qualitative study. <i>BMC Public Health</i> , 2021 , 21, 203	4.1	
183	The division and transfer of care responsibilities in paediatric type 1 diabetes: A qualitative study on parental perspectives. <i>Journal of Advanced Nursing</i> , 2021 , 77, 1968-1979	3.1	2

(2020-2021)

182	The suitability of patient-reported outcome measures used to assess the impact of hypoglycaemia on quality of life in people with diabetes: a systematic review using COSMIN methods. <i>Diabetologia</i> , 2021 , 64, 1213-1225	10.3	4
181	The impact of hypoglycaemia on the quality of life of family members of adults with type 1 or type 2 diabetes: A qualitative systematic review. <i>Diabetic Medicine</i> , 2021 , 38, e14666	3.5	1
180	Prover again will I be carefree Pa qualitative study of the impact of hypoglycemia on quality of life among adults with type 1 diabetes. BMJ Open Diabetes Research and Care, 2021, 9,	4.5	1
179	Prevalence of and risk factors for sexual dysfunctions in adults with type 1 or type 2 diabetes: Results from Diabetes MILES - Flanders. <i>Diabetic Medicine</i> , 2021 , e14676	3.5	2
178	The bidirectional longitudinal association between depressive symptoms and HbA: A systematic review and meta-analysis. <i>Diabetic Medicine</i> , 2021 , e14671	3.5	4
177	Changes in quality of life following hypoglycaemia in adults with type 2 diabetes: A systematic review of longitudinal studies. <i>Diabetic Medicine</i> , 2021 , e14706	3.5	O
176	Overeating, binge eating, quality of life, emotional difficulties, and HbA in adolescents with type 1 diabetes: A Danish national survey. <i>Diabetes Research and Clinical Practice</i> , 2021 , 182, 109150	7.4	О
175	The prevalence of type 2 diabetes in people with psychiatric disorders: an umbrella review protocol. <i>Systematic Reviews</i> , 2020 , 9, 101	3	1
174	Correlates of fear of hypoglycemia among patients with type 1 and 2 diabetes mellitus in outpatient hospitals in Zambia. <i>International Journal of Diabetes in Developing Countries</i> , 2020 , 40, 619-6	526 ⁸	1
173	Psychological correlates of disordered eating in youth with type 1 diabetes: Results from diabetes MILES Youth-Australia. <i>Pediatric Diabetes</i> , 2020 , 21, 664-672	3.6	5
172	Reducing the burden of hypoglycaemia in people with diabetes through increased understanding: design of the Hypoglycaemia REdefining SOLutions for better liVEs (Hypo-RESOLVE) project. <i>Diabetic Medicine</i> , 2020 , 37, 1066-1073	3.5	12
171	Prevalence and correlates of diabetes distress, perceived stress and depressive symptoms among adults with early-onset Type 2 diabetes: cross-sectional survey results from the Danish DD2 study. <i>Diabetic Medicine</i> , 2020 , 37, 1679-1687	3.5	7
170	Depressive and anxiety symptoms and following of the Dutch Dietary Guidelines 2015 in adults with diabetes: Results from Diabetes MILES-The Netherlands. <i>Journal of Psychosomatic Research</i> , 2020 , 135, 110160	4.1	3
169	Back to the future: 25 years of © uidelines for encouraging psychological well-beingPamong people affected by diabetes. <i>Diabetic Medicine</i> , 2020 , 37, 1225-1229	3.5	16
168	How 25 years of psychosocial research has contributed to a better understanding of the links between depression and diabetes. <i>Diabetic Medicine</i> , 2020 , 37, 383-392	3.5	17
167	Psychosocial research in the diabetic foot: Are we making progress?. <i>Diabetes/Metabolism Research and Reviews</i> , 2020 , 36 Suppl 1, e3257	7.5	12
166	P wish my health professionals understood that itB not just all about your HbA!P Qualitative responses from the second Diabetes MILES - Australia (MILES-2) study. <i>Diabetic Medicine</i> , 2020 , 37, 971-	9851	15
165	Positive facilitators of diabetes management in emerging adults with type 1 diabetes-A qualitative analysis of blogs. <i>Endocrinology, Diabetes and Metabolism</i> , 2020 , 3, e00161	2.7	3

164	Social support and self-care outcomes in adults with diabetes: The mediating effects of self-efficacy and diabetes distress. Results of the second diabetes MILES - Australia (MILES-2) study. <i>Diabetes Research and Clinical Practice</i> , 2020 , 166, 108314	7.4	5
163	Psychosocial Moderators of the Impact of Diabetes Stigma: Results From the Second Diabetes MILES - Australia (MILES-2) Study. <i>Diabetes Care</i> , 2020 , 43, 2651-2659	14.6	7
162	Sustainable improvement of HbA and satisfaction with diabetes care after adding telemedicine in patients on adaptable insulin regimens: Results of the TeleDiabetes randomized controlled trial. Health Informatics Journal, 2020 , 26, 628-641	3	7
161	Prevalence of Insomnia (Symptoms) in T2D and Association With Metabolic Parameters and Glycemic Control: Meta-Analysis. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020 , 105,	5.6	11
160	The role of parental support for emerging adults with type 1 diabetes: A scoping review. <i>Pediatric Diabetes</i> , 2020 , 21, 995-1030	3.6	5
159	Longitudinal associations between depression and diabetes complications: a systematic review and meta-analysis. <i>Diabetic Medicine</i> , 2019 , 36, 1562-1572	3.5	66
158	What Do Adults with Type 2 Diabetes Want from the "Perfect" App? Results from the Second Diabetes MILES: Australia (MILES-2) Study. <i>Diabetes Technology and Therapeutics</i> , 2019 , 21, 393-399	8.1	17
157	Psychiatric disorders as risk factors for the development of type 2 diabetes mellitus: an umbrella review protocol. <i>BMJ Open</i> , 2019 , 9, e024981	3	3
156	Modeling Interactions Between Latent Variables in Research on Type D Personality: A Monte Carlo Simulation and Clinical Study of Depression and Anxiety. <i>Multivariate Behavioral Research</i> , 2019 , 54, 637	- 2 6 8 5	16
155	Assessing the perceived impact of diabetes on quality of life: Psychometric validation of the DAWN2 Impact of Diabetes Profile in the second Diabetes MILES - Australia (MILES-2) survey. <i>Diabetes Research and Clinical Practice</i> , 2019 , 150, 253-263	7.4	12
154	Anxiety is common and costly in T2DM - why psychology matters. <i>Nature Reviews Endocrinology</i> , 2019 , 15, 567-568	15.2	3
153	Study protocol of Diabetes LEAP: a longitudinal study examining emotional problems in adolescents with type 1 diabetes and their parents/caregivers. <i>BMC Pediatrics</i> , 2019 , 19, 377	2.6	1
152	Comment on the consensus report on the management of hyperglycaemia in Type 2 diabetes by the American Diabetes Association and the European Association for the Study of Diabetes. <i>Diabetic Medicine</i> , 2019 , 36, 911-912	3.5	1
151	Is Self-Compassion Related to Behavioural, Clinical and Emotional Outcomes in Adults with Diabetes? Results from the Second Diabetes MILESAustralia (MILES-2) Study. <i>Mindfulness</i> , 2019 , 10, 1222-1231	2.9	10
150	Prospective memory slips are associated with forgetting to take glucose-lowering therapies among adults with diabetes: results from the second Diabetes MILES - Australia (MILES-2) survey. <i>Diabetic Medicine</i> , 2019 , 36, 569-577	3.5	3
149	Mindfulness and fear of hypoglycaemia in parents of children with Type 1 diabetes: results from Diabetes MILES Youth - The Netherlands. <i>Diabetic Medicine</i> , 2018 , 35, 650-657	3.5	12
148	Diabetes distress is more strongly associated with HbA1c than depressive symptoms in adolescents with type 1 diabetes: Results from Diabetes MILES Youth-Australia. <i>Pediatric Diabetes</i> , 2018 , 19, 840-847	3 .6	37
147	Diabetes stigma is associated with negative treatment appraisals among adults with insulin-treated Type 2 diabetes: results from the second Diabetes MILES - Australia (MILES-2) survey. <i>Diabetic Medicine</i> 2018 35, 658-662	3.5	15

146	Comment on Young-Hyman et al. Psychosocial Care for People With Diabetes: A Position Statement of the American Diabetes Association. Diabetes Care 2016;39:2126-2140. <i>Diabetes Care</i> , 2018 , 41, e31-6	14 .6	3
145	What is the best measure for assessing diabetes distress? A comparison of the Problem Areas in Diabetes and Diabetes Distress Scale: results from Diabetes MILES-Australia. <i>Journal of Health Psychology</i> , 2018 , 23, 667-680	3.1	29
144	Identification of barriers to insulin therapy and approaches to overcoming them. <i>Diabetes, Obesity and Metabolism</i> , 2018 , 20, 488-496	6.7	110
143	More work is needed to better understand diabetes distress as a predictor of all-cause mortality in type 2 diabetes. <i>Diabetologia</i> , 2018 , 61, 2247-2248	10.3	1
142	Emotional Well-Being Factors Associated with Sexual Dysfunction in Adults with Type 1 or Type 2 Diabetes: Results from Diabetes MILESAustralia. <i>International Journal of Sexual Health</i> , 2018 , 30, 237-24	19	6
141	Type D personality and social relations in adults with diabetes: results from diabetes MILES - The Netherlands. <i>Psychology and Health</i> , 2018 , 33, 1456-1471	2.9	6
140	The associations between diabetes distress and self-efficacy, medication adherence, self-care activities and disease control depend on the way diabetes distress is measured: Comparing the DDS-17, DDS-2 and the PAID-5. <i>Diabetes Research and Clinical Practice</i> , 2018 , 142, 74-84	7.4	13
139	Advanced Glycation End Product (AGE) Accumulation in the Skin is Associated with Depression: The Maastricht Study. <i>Depression and Anxiety</i> , 2017 , 34, 59-67	8.4	19
138	Paediatric parenting stress in fathers and mothers of young children with Type 1 diabetes: a longitudinal study. <i>Diabetic Medicine</i> , 2017 , 34, 821-827	3.5	14
137	Cohort profiles of the cross-sectional and prospective participant groups in the second Diabetes MILES-Australia (MILES-2) study. <i>BMJ Open</i> , 2017 , 7, e012926	3	16
136	Strengths, Risk Factors, and Resilient Outcomes in Adolescents With Type 1 Diabetes: Results From Diabetes MILES Youth-Australia. <i>Diabetes Care</i> , 2017 , 40, 849-855	14.6	15
135	Fatigue in type 1 diabetes: an understudied problem. <i>Lancet Diabetes and Endocrinology,the</i> , 2017 , 5, 409-410	18.1	1
134	Diabetes Complications and Depressive Symptoms: Prospective Results From the Montreal Diabetes Health and Well-Being Study. <i>Psychosomatic Medicine</i> , 2017 , 79, 603-612	3.7	10
133	Assessing Psychological Insulin Resistance in Type 2 Diabetes: a Critical Comparison of Measures. <i>Current Diabetes Reports</i> , 2017 , 17, 46	5.6	7
132	Association of Microvascular Dysfunction With Late-Life Depression: A Systematic Review and Meta-analysis. <i>JAMA Psychiatry</i> , 2017 , 74, 729-739	14.5	138
131	Associations Between Physical Activity and Depressive Symptoms by Weight Status Among Adults With Type 2 Diabetes: Results From Diabetes MILES-Australia. <i>Journal of Physical Activity and Health</i> , 2017 , 14, 195-202	2.5	4
130	Severe hypoglycemia, impaired awareness of hypoglycemia, and self-monitoring in adults with type 1 diabetes: Results from Diabetes MILES-Australia. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 577-582	3.2	15
129	The Use of Mobile Applications Among Adults with Type 1 and Type 2 Diabetes: Results from the Second MILES-Australia (MILES-2) Study. <i>Diabetes Technology and Therapeutics</i> , 2017 , 19, 730-738	8.1	25

128	Depressive Symptoms Are Negatively Associated with Glucose Testing and Eating Meals on Time among Individuals with Diabetes in Zambia. <i>Diabetes and Metabolism Journal</i> , 2017 , 41, 440-448	5	3
127	Comment on Young-Hyman et al. Psychosocial Care for People With Diabetes: A Position Statement of the American Diabetes Association. Diabetes Care 2016;39:2126-2140. <i>Diabetes Care</i> , 2017 , 40, e127-	e ¹ 128	1
126	Gender differences in disordered eating behaviors and body dissatisfaction among adolescents with type 1 diabetes: Results from diabetes MILES youth-Australia. <i>International Journal of Eating Disorders</i> , 2017 , 50, 1183-1193	6.3	31
125	Cut Points for Identifying Clinically Significant Diabetes Distress in Adolescents With Type 1 Diabetes Using the PAID-T: Results From Diabetes MILES Youth-Australia. <i>Diabetes Care</i> , 2017 , 40, 1462	-1468	16
124	Diabetes in Sub-Saharan African Children: Risks, Care, and Challenges 2017 , 157-172		
123	Low vitamin D levels are not a contributing factor to higher prevalence of depressive symptoms in people with Type 2 diabetes mellitus: the Hoorn study. <i>Diabetic Medicine</i> , 2017 , 34, 577-581	3.5	4
122	The relationship between parenting stress and parent-child interaction with health outcomes in the youngest patients with type 1 diabetes (0-7 years). <i>European Journal of Pediatrics</i> , 2016 , 175, 329-38	4.1	21
121	Association of Type D personality with increased vulnerability to depression: Is there a role for inflammation or endothelial dysfunction? - The Maastricht Study. <i>Journal of Affective Disorders</i> , 2016 , 189, 118-25	6.6	32
120	Subjective Wellbeing Among Adults with Diabetes: Results from Diabetes MILESAustralia. Journal of Happiness Studies, 2016 , 17, 1205-1217	3.7	8
119	Diabetes MILES Youth-Australia: methods and sample characteristics of a national survey of the psychological aspects of living with type 1 diabetes in Australian youth and their parents. <i>BMC Psychology</i> , 2016 , 4, 42	2.8	19
118	Diabetes, Depression, and Cardiovascular Risk 2016 , 831-847		1
117	Effect of vitamin D supplementation on health status in non-vitamin D deficient people with type 2 diabetes mellitus. <i>Endocrine Connections</i> , 2016 , 5, 61-69	3.5	6
116	Review of research grant allocation to psychosocial studies in diabetes research. <i>Diabetic Medicine</i> , 2016 , 33, 1673-1676	3.5	5
115	The Use of Mobile Applications Among Adolescents with Type 1 Diabetes: Results from Diabetes MILES Youth-Australia. <i>Diabetes Technology and Therapeutics</i> , 2016 , 18, 813-819	8.1	11
114	Vitamin D status and health-related quality of life in patients with Type 2 diabetes. <i>Diabetic Medicine</i> , 2016 , 33, 300-6	3.5	5
113	Psychological and personality factors in type 2 diabetes mellitus, presenting the rationale and exploratory results from The Maastricht Study, a population-based cohort study. <i>BMC Psychiatry</i> , 2016 , 16, 17	4.2	34
112	Prospectively measured lifestyle factors and BMI explain differences in health-related quality of life between colorectal cancer patients with and without comorbid diabetes. <i>Supportive Care in Cancer</i> , 2016 , 24, 2591-601	3.9	12
111	Does self-efficacy mediate the cross-sectional relationship between perceived quality of health care and self-management of diabetes? Results from Diabetes MILES - Australia. <i>Psychology and Health</i> , 2016 , 31, 592-604	2.9	6

110	Associations of low grade inflammation and endothelial dysfunction with depression - The Maastricht Study. <i>Brain, Behavior, and Immunity</i> , 2016 , 56, 390-6	16.6	71
109	Explaining psychological insulin resistance in adults with non-insulin-treated type 2 diabetes: The roles of diabetes distress and current medication concerns. Results from Diabetes MILESAustralia. <i>Primary Care Diabetes</i> , 2016 , 10, 75-82	2.4	13
108	The impact of having both cancer and diabetes on patient-reported outcomes: a systematic review and directions for future research. <i>Journal of Cancer Survivorship</i> , 2016 , 10, 406-15	5.1	22
107	Anxiety and Depressive Symptoms as Predictors of All-Cause Mortality among People with Insulin-NaMe Type 2 Diabetes: 17-Year Follow-Up of the Second Nord-TrEdelag Health Survey (HUNT2), Norway. <i>PLoS ONE</i> , 2016 , 11, e0160861	3.7	8
106	Suicidal ideation reported by adults with Type 1 or Type 2 diabetes: results from Diabetes MILES-Australia. <i>Diabetic Medicine</i> , 2016 , 33, 1582-1589	3.5	16
105	Psychological Factors and Diabetes Mellitus 2016 , 819-839		
104	The Association of Mindful Parenting with Glycemic Control and Quality of Life in Adolescents with Type 1 Diabetes: Results from Diabetes MILES-The Netherlands. <i>Mindfulness</i> , 2016 , 7, 1227-1237	2.9	10
103	The impact of diabetes on neuropathic symptoms and receipt of chemotherapy among colorectal cancer patients: results from the PROFILES registry. <i>Journal of Cancer Survivorship</i> , 2015 , 9, 523-31	5.1	13
102	Negative appraisals of insulin therapy are common among adults with Type 2 diabetes using insulin: Results from Diabetes MILES - Australia cross-sectional survey. <i>Diabetic Medicine</i> , 2015 , 32, 1297-303	3.5	27
101	Effectiveness of a stepped care intervention for anxiety and depression in people with diabetes, asthma or COPD in primary care: A randomized controlled trial. <i>Journal of Affective Disorders</i> , 2015 , 184, 269-76	6.6	33
100	Stressful life events and incident metabolic syndrome: the Hoorn study. Stress, 2015, 18, 507-13	3	19
99	Anxiety, depression and timing of insulin treatment among people with type 2 diabetes: Nine-year follow-up of the Nord-TrEdelag Health Study, Norway. <i>Journal of Psychosomatic Research</i> , 2015 , 79, 309-15	4.1	12
98	Type D personality, suboptimal health behaviors and emotional distress in adults with diabetes: results from Diabetes MILES-The Netherlands. <i>Diabetes Research and Clinical Practice</i> , 2015 , 108, 94-10.	5 ^{7.4}	33
97	Impact of cancer on use of glucose-lowering drug treatment in individuals with diabetes: potential mechanisms. <i>Diabetologia</i> , 2015 , 58, 1378-9	10.3	3
96	The association between glucose-lowering drug use and mortality among breast cancer patients with type 2 diabetes. <i>Breast Cancer Research and Treatment</i> , 2015 , 150, 427-37	4.4	33
95	Living with type 1 diabetes is challenging for Zambian adolescents: qualitative data on stress, coping with stress and quality of care and life. <i>BMC Endocrine Disorders</i> , 2015 , 15, 20	3.3	30
94	Diabetes mellitus and comorbid depression in Zambia. <i>Diabetic Medicine</i> , 2015 , 32, 814-8	3.5	6
93	If it does not significantly change HbA1c levels why should we waste time on it? A plea for the prioritization of psychological well-being in people with diabetes. <i>Diabetic Medicine</i> , 2015 , 32, 155-63	3.5	52

92	Depression, anxiety and self-care behaviours of young adults with Type 2 diabetes: results from the International Diabetes Management and Impact for Long-term Empowerment and Success (MILES) Study. <i>Diabetic Medicine</i> , 2015 , 32, 133-40	3.5	44
91	Fear of hypoglycaemia in adults with Type diabetes: results from Diabetes MILES - The Netherlands. <i>Diabetic Medicine</i> , 2015 , 32, 1289-96	3.5	24
90	Psychological and pharmacological interventions for depression in patients with diabetes mellitus: an abridged Cochrane review: some critical comments. <i>Diabetic Medicine</i> , 2015 , 32, 839	3.5	1
89	Suicidal ideation or non-suicidal self-harm? A mismatch between the DSM-IV criterion and PHQ-9 item nine. <i>Diabetes Research and Clinical Practice</i> , 2015 , 108, e5-6	7.4	4
88	Diabetes, Depression, and Cardiovascular Risk 2015 , 1-17		
87	Intraperitoneal versus subcutaneous insulin therapy in the treatment of type 1 diabetes mellitus. <i>Netherlands Journal of Medicine</i> , 2015 , 73, 399-409	0.5	7
86	Qualitative observation instrument to measure the quality of parent-child interactions in young children with type 1 diabetes mellitus. <i>BMC Pediatrics</i> , 2014 , 14, 145	2.6	4
85	The individual and combined effect of colorectal cancer and diabetes on health-related quality of life and sexual functioning: results from the PROFILES registry. <i>Supportive Care in Cancer</i> , 2014 , 22, 307	1399	12
84	Endothelial dysfunction is associated with a greater depressive symptom score in a general elderly population: the Hoorn Study. <i>Psychological Medicine</i> , 2014 , 44, 1403-16	6.9	48
83	Comment on Schmitz et al. Recurrent subthreshold depression in type 2 diabetes: an important risk factor for poor health outcomes. Diabetes care 2014;37:970-978. <i>Diabetes Care</i> , 2014 , 37, e145	14.6	1
82	Severe obesity and diabetes self-care attitudes, behaviours and burden: implications for weight management from a matched case-controlled study. Results from Diabetes MILESAustralia. <i>Diabetic Medicine</i> , 2014 , 31, 232-40	3.5	12
81	Associations between economic hardship and markers of self-management in adults with type 2 diabetes: results from Diabetes MILES - Australia. <i>Australian and New Zealand Journal of Public Health</i> , 2014 , 38, 466-72	2.3	12
80	Effects of oxidative stress on fatty acid- and one-carbon-metabolism in psychiatric and cardiovascular disease comorbidity. <i>Acta Psychiatrica Scandinavica</i> , 2014 , 130, 163-80	6.5	83
79	Response to Kawada. Depressive symptoms and HbA1c in patients with Type 1 and Type 2 diabetes. <i>Diabetic Medicine</i> , 2014 , 31, 760-1	3.5	
78	Diabetes-specific emotional distress in people with Type 2 diabetes: a comparison between primary and secondary care. <i>Diabetic Medicine</i> , 2014 , 31, 1252-9	3.5	39
77	Toward defining a cutoff score for elevated fear of hypoglycemia on the hypoglycemia fear survey worry subscale in patients with type 2 diabetes. <i>Diabetes Care</i> , 2014 , 37, 102-8	14.6	31
76	Further investigation of the psychometric properties of the Insulin Treatment Appraisal Scale among insulin-using and non-insulin-using adults with type 2 diabetes: results from Diabetes MILES-Australia. <i>Health and Quality of Life Outcomes</i> , 2014 , 12, 87	3	14
75	Psychometric properties of the Exercise Self-efficacy Scale in Dutch Primary care patients with type 2 diabetes mellitus. <i>International Journal of Behavioral Medicine</i> , 2014 , 21, 394-401	2.6	18

(2013-2014)

74	Managing hypoglycemia in diabetes may be more fear management than glucose management: a practical guide for diabetes care providers. <i>Current Diabetes Reviews</i> , 2014 , 10, 364-70	2.7	29
73	Self-reported access to and quality of healthcare for diabetes: do the severely obese experience equal access?. <i>Australian Family Physician</i> , 2014 , 43, 552-6		1
72	Measuring and Assessing Depression in People with Diabetes: Implications for Clinical Practice 2013 , 199-209		2
71	Diabetes Attitudes, Wishes and Needs second study (DAWN2) cross-national benchmarking indicators for family members living with people with diabetes. <i>Diabetic Medicine</i> , 2013 , 30, 778-88	3.5	168
70	Does the shortage of diabetes specialists in regional and rural Australia matter? Results from Diabetes MILESAustralia. <i>Diabetes Research and Clinical Practice</i> , 2013 , 100, 222-9	7.4	11
69	Diabetes Attitudes Wishes and Needs 2 (DAWN2): a multinational, multi-stakeholder study of psychosocial issues in diabetes and person-centred diabetes care. <i>Diabetes Research and Clinical Practice</i> , 2013 , 99, 174-84	7.4	159
68	The impact of comorbidity on Health-Related Quality of Life among cancer survivors: analyses of data from the PROFILES registry. <i>Journal of Cancer Survivorship</i> , 2013 , 7, 602-13	5.1	79
67	Depressive symptoms, insulin sensitivity and insulin secretion in the RISC cohort study. <i>Diabetes and Metabolism</i> , 2013 , 39, 42-9	5.4	7
66	Severely obese people with diabetes experience impaired emotional well-being associated with socioeconomic disadvantage: results from diabetes MILES - Australia. <i>Diabetes Research and Clinical Practice</i> , 2013 , 101, 131-40	7.4	24
65	Ethnic aspects of emotional distress in patients with diabetesthe Amsterdam Health Monitor Study. <i>Diabetic Medicine</i> , 2013 , 30, e25-31	3.5	11
64	Psychometric and screening properties of the WHO-5 well-being index in adult outpatients with Type or Type diabetes mellitus. <i>Diabetic Medicine</i> , 2013 , 30, e63-9	3.5	119
63	Effects of exercise training on quality of life, symptoms of depression, symptoms of anxiety and emotional well-being in type 2 diabetes mellitus: a systematic review. <i>Diabetologia</i> , 2013 , 56, 1210-25	10.3	60
62	PS12 - 7. The quality of parent-child interaction and disease-related parenting stress is associated with HbA1c and quality of life in young children with type 1 diabetes <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2013 , 11, 180-180	Ο	
61	Screening for type 2 diabetes does not reduce mortality over 10 years. <i>Evidence-Based Medicine</i> , 2013 , 18, e44		
60	Differential associations between depressive symptoms and glycaemic control in outpatients with diabetes. <i>Diabetic Medicine</i> , 2013 , 30, e115-22	3.5	56
59	The association between diabetes and an episode of depressive symptoms in the 2002 World Health Survey: an analysis of 231,797 individuals from 47 countries. <i>Diabetic Medicine</i> , 2013 , 30, e208-1	4 ^{3·5}	50
58	Diabetes Attitudes, Wishes and Needs second study (DAWN2) cross-national benchmarking of diabetes-related psychosocial outcomes for people with diabetes. <i>Diabetic Medicine</i> , 2013 , 30, 767-77	3.5	405
57	Compromised quality of life in patients with both Type 1 diabetes mellitus and coeliac disease. <i>Diabetic Medicine</i> , 2013 , 30, 835-9	3.5	17

56	The effects of a mindfulness-based intervention on emotional distress, quality of life, and HbA(1c) in outpatients with diabetes (DiaMind): a randomized controlled trial. <i>Diabetes Care</i> , 2013 , 36, 823-30	14.6	105
55	Suboptimal glycemic control in type 2 diabetes: a key role for anhedonia?. <i>Journal of Psychiatric Research</i> , 2012 , 46, 549-54	5.2	37
54	Diabetes MILESAustralia (management and impact for long-term empowerment and success): methods and sample characteristics of a national survey of the psychological aspects of living with type 1 or type 2 diabetes in Australian adults. <i>BMC Public Health</i> , 2012 , 12, 120	4.1	46
53	The course of depressive symptoms in primary care patients with type 2 diabetes: results from the Diabetes, Depression, Type D Personality Zuidoost-Brabant (DiaDDZoB) Study. <i>Diabetologia</i> , 2012 , 55, 608-16	10.3	60
52	The longitudinal association between glycaemic control and health-related quality of life following insulin therapy optimisation in type 2 diabetes patients. A prospective observational study in secondary care. <i>Quality of Life Research</i> , 2012 , 21, 1359-65	3.7	21
51	Screening tools used for measuring depression among people with Type 1 and Type 2 diabetes: a systematic review. <i>Diabetic Medicine</i> , 2012 , 29, 164-75	3.5	92
50	Association of coexisting diabetes and depression with mortality after myocardial infarction. <i>Diabetes Care</i> , 2012 , 35, 503-9	14.6	35
49	Quality of life of children with type 1 diabetes: a systematic review. <i>Current Diabetes Reviews</i> , 2012 , 8, 434-43	2.7	62
48	Initiation of insulin glargine in patients with Type 2 diabetes in suboptimal glycaemic control positively impacts health-related quality of life. A prospective cohort study in primary care. <i>Diabetic Medicine</i> , 2011 , 28, 1096-102	3.5	26
47	Limited effect of screening for depression with written feedback in outpatients with diabetes mellitus: a randomised controlled trial. <i>Diabetologia</i> , 2011 , 54, 741-8	10.3	49
46	Quality of the parent-child interaction in young children with type 1 diabetes mellitus: study protocol. <i>BMC Pediatrics</i> , 2011 , 11, 28	2.6	7
45	Web-based depression treatment for type 1 and type 2 diabetic patients: a randomized, controlled trial. <i>Diabetes Care</i> , 2011 , 34, 320-5	14.6	156
44	PS11 - 59. Development of a new observational rating scale to measure the quality of parent-child interactions in young children with type 1 diabetes mellitus. <i>Nederlands Tijdschrift Voor Diabetologie</i> , 2011 , 9, 131-131	О	
43	Prevalence of depression in individuals with impaired glucose metabolism or undiagnosed diabetes: a systematic review and meta-analysis of the European Depression in Diabetes (EDID) Research Consortium. <i>Diabetes Care</i> , 2011 , 34, 752-62	14.6	157
42	Prevalence of comorbid depression is high in out-patients with Type 1 or Type 2 diabetes mellitus. Results from three out-patient clinics in the Netherlands. <i>Diabetic Medicine</i> , 2010 , 27, 217-24	3.5	110
41	Diabetes-specific emotional distress mediates the association between depressive symptoms and glycaemic control in Type 1 and Type 2 diabetes. <i>Diabetic Medicine</i> , 2010 , 27, 798-803	3.5	108
40	The Epidemiology of Depression and Diabetes 2010 , 1-27		8
39	Type 2 diabetes mellitus as a risk factor for the onset of depression: a systematic review and meta-analysis. <i>Diabetologia</i> , 2010 , 53, 2480-6	10.3	470

(2006-2010)

38	Eicosapentaenoic acid as an add-on to antidepressant medication for co-morbid major depression in patients with diabetes mellitus: a randomized, double-blind placebo-controlled study. <i>Journal of Affective Disorders</i> , 2010 , 126, 282-6	6.6	42
37	Does emotional stress cause type 2 diabetes mellitus? A review from the European Depression in Diabetes (EDID) Research Consortium. <i>Discovery Medicine</i> , 2010 , 9, 112-8	2.5	125
36	Depressive symptoms are associated with physical inactivity in patients with type 2 diabetes. The DIAZOB Primary Care Diabetes study. <i>Family Practice</i> , 2009 , 26, 171-3	1.9	45
35	Associations between vascular co-morbidities and depression in insulin-naive diabetes patients: the DIAZOB Primary Care Diabetes study. <i>Diabetologia</i> , 2009 , 52, 2056-63	10.3	23
34	Symptoms of depression and diabetes-specific emotional distress are associated with a negative appraisal of insulin therapy in insulin-nalle patients with Type 2 diabetes mellitus. A study from the European Depression in Diabetes [EDID] Research Consortium. <i>Diabetic Medicine</i> , 2009 , 26, 28-33	3.5	56
33	Online cognitieve gedragstherapie voor diabetespatiliten met depressieve klachten: een gerandomiseerd gecontroleerd onderzoek. <i>Psychologie and Gezondheid</i> , 2009 , 37, 222-228		2
32	Serious diabetes-specific emotional problems in patients with type 2 diabetes who have different levels of comorbid depression: a Polish study from the European Depression in Diabetes (EDID) Research Consortium. <i>European Psychiatry</i> , 2009 , 24, 425-30	6	39
31	Symptoms of depression in people with impaired glucose metabolism or Type 2 diabetes mellitus: The Hoorn Study. <i>Diabetic Medicine</i> , 2008 , 25, 843-9	3.5	40
30	Personality characteristics in adolescence predict long-term changes in body fatness. <i>Diabetes Research and Clinical Practice</i> , 2008 , 79, e10-3	7.4	1
29	Diabetes-related symptom distress in association with glucose metabolism and comorbidity: the Hoorn Study. <i>Diabetes Care</i> , 2008 , 31, 2268-70	14.6	38
28	Omissions impact measurements of patient-reported outcomes. <i>Quality of Life Research</i> , 2008 , 17, 963-4; author reply 965-6	3.7	1
27	Web-based cognitive behavioural therapy (W-CBT) for diabetes patients with co-morbid depression: design of a randomised controlled trial. <i>BMC Psychiatry</i> , 2008 , 8, 9	4.2	33
26	Psychosociale interventies bij mensen met diabetes. <i>Psychologie and Gezondheid</i> , 2008 , 36, 209-220		
25	Monitoring health related quality of life in adolescents with diabetes: a review of measures. <i>Archives of Disease in Childhood</i> , 2007 , 92, 434-9	2.2	52
24	Development and validation of the insulin treatment appraisal scale (ITAS) in patients with type 2 diabetes. <i>Health and Quality of Life Outcomes</i> , 2007 , 5, 69	3	83
23	No evidence for increased self-reported cognitive failure in Type 1 and Type 2 diabetes: a cross-sectional study. <i>Diabetic Medicine</i> , 2007 , 24, 735-40	3.5	11
22	Validation of the WHO-5 Well-Being Index in adolescents with type 1 diabetes. <i>Diabetes Care</i> , 2007 , 30, 2003-6	14.6	172
21	Depression as a risk factor for the onset of type 2 diabetes mellitus. A meta-analysis. <i>Diabetologia</i> , 2006 , 49, 837-45	10.3	653

20	Reply to comment on: Knol MJ, Twisk JWR, Beekman ATF, Heine RJ, Snoek FJ, Pouwer F (2006) Depression as a risk factor for the onset of type 2 diabetes mellitus. A meta-analysis. Diabetologia 49:837 B 45. <i>Diabetologia</i> , 2006 , 49, 2799-2800	10.3	3
19	Associations between depressive symptoms and insulin resistance: the Hoorn Study. <i>Diabetologia</i> , 2006 , 49, 2874-7	10.3	45
18	Fat food for a bad mood. Could we treat and prevent depression in Type 2 diabetes by means of omega-3 polyunsaturated fatty acids? A review of the evidence. <i>Diabetic Medicine</i> , 2005 , 22, 1465-75	3.5	19
17	Depressive symptoms, insulin resistance, and risk of diabetes in women at midlife: response to Everson-Rose et al. <i>Diabetes Care</i> , 2005 , 28, 1265-6; author reply 1266	14.6	2
16	The confidence in diabetes self-care scale: psychometric properties of a new measure of diabetes-specific self-efficacy in Dutch and US patients with type 1 diabetes. <i>Diabetes Care</i> , 2003 , 26, 713-8	14.6	124
15	to: M. Koopmanschap: Coping with Type 2 diabetes: the patientß perspective. Diabetologia 45:S18-S22. <i>Diabetologia</i> , 2003 , 46, 302-3	10.3	
14	Rates and risks for co-morbid depression in patients with Type 2 diabetes mellitus: results from a community-based study. <i>Diabetologia</i> , 2003 , 46, 892-8	10.3	150
13	Continuous intraperitoneal insulin infusion in patients with PorittlePdiabetes: favourable effects on glycaemic control and hospital stay. <i>Diabetic Medicine</i> , 2002 , 19, 496-501	3.5	28
12	PatientsPEvaluation of the Quality of Diabetes Care (PEQD): development and validation of a new instrument. <i>Quality and Safety in Health Care</i> , 2002 , 11, 131-6		23
11	Association between symptoms of depression and glycaemic control may be unstable across gender. <i>Diabetic Medicine</i> , 2001 , 18, 595-8	3.5	40
10	Monitoring of psychological well-being in outpatients with diabetes: effects on mood, HbA(1c), and the patientB evaluation of the quality of diabetes care: a randomized controlled trial. <i>Diabetes Care</i> , 2001 , 24, 1929-35	14.6	97
9	The well-being questionnaire: evidence for a three-factor structure with 12 items (W-BQ12). <i>Psychological Medicine</i> , 2000 , 30, 455-62	6.9	43
8	Psychological distress two years after diagnosis of breast cancer: frequency and prediction. <i>Patient Education and Counseling</i> , 2000 , 40, 209-17	3.1	166
7	Diabetes-related emotional distress in Dutch and U.S. diabetic patients: cross-cultural validity of the problem areas in diabetes scale. <i>Diabetes Care</i> , 2000 , 23, 1305-9	14.6	316
6	Diabetes Fear of Injecting and Self-Testing Questionnaire: a psychometric evaluation. <i>Diabetes Care</i> , 2000 , 23, 765-9	14.6	65
5	Factor structure of the Disability and Impact Profile in patients with multiple sclerosis. <i>Quality of Life Research</i> , 1999 , 8, 141-50	3.7	9
4	The 12-item well-being questionnaire. An evaluation of its validity and reliability in Dutch people with diabetes. <i>Diabetes Care</i> , 1999 , 22, 2004-10	14.6	66
3	A comparison of the standard and the computerized versions of the Well-being Questionnaire (WBQ) and the Diabetes Treatment Satisfaction Questionnaire (DTSQ). <i>Quality of Life Research</i> , 1998 , 7, 33-8	3.7	70

LIST OF PUBLICATIONS

Ceiling effect reduces the validity of the Diabetes Treatment Satisfaction Questionnaire. *Diabetes Care*, **1998**, 21, 2039

14.6 19

Mindfulness in Relation to Diet Quality in Adults with Type 1 and Type 2 Diabetes: Results from Diabetes MILES-The Netherlands. *Mindfulness*,1

2.9