

# Gil F Salles

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/630992/publications.pdf>

Version: 2024-02-01

103  
papers

5,023  
citations

125106

35  
h-index

111975

67  
g-index

103  
all docs

103  
docs citations

103  
times ranked

6459  
citing authors

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Prevalence and associated factors of aspiration and severe dysphagia in asymptomatic patients in the late period after open partial laryngectomy: a videofluoroscopic evaluation. <i>European Archives of Oto-Rhino-Laryngology</i> , 2022, , .  | 0.8 | 0         |
| 2  | Prognostic Value of Changes in Aortic Stiffness for Cardiovascular Outcomes and Mortality in Resistant Hypertension: a Cohort Study. <i>Hypertension</i> , 2022, 79, 447-456.  | 1.3 | 14        |
| 3  | Refractory Hypertension: a Narrative Systematic Review with Emphasis on Prognosis. <i>Current Hypertension Reports</i> , 2022, 24, 95-106.   | 1.5 | 4         |
| 4  | Differential effects of treatment targets on risks of adverse outcomes according to diabetes duration, age and complications: Can these characteristics be used to individualize diabetes treatment? The Rio de Janeiro type 2 diabetes cohort. <i>Journal of Diabetes and Its Complications</i> , 2022, 36, 108124. | 1.2 | 5         |
| 5  | Associations of the nocturnal blood pressure fall and morning surge with cardiovascular events and mortality in individuals with resistant hypertension. <i>Journal of Hypertension</i> , 2021, 39, 1177-1187.   | 0.3 | 14        |
| 6  | Traditional and non-traditional risk factors for peripheral artery disease development/progression in patients with type 2 diabetes: the Rio de Janeiro type 2 diabetes cohort study. <i>Cardiovascular Diabetology</i> , 2021, 20, 54.  | 2.7 | 17        |
| 7  | Diretrizes Brasileiras de Hipertens o Arterial â€“ 2020. <i>Arquivos Brasileiros De Cardiologia</i> , 2021, 116, 516-658.  | 0.3 | 340       |
| 8  | Importance of non-invasive liver fibrosis scores for mortality and complications development in individuals with type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2021, 35, 107879.   | 1.2 | 10        |
| 9  | Associations Between Achieved Ambulatory Blood Pressures and Its Changes With Adverse Outcomes in Resistant Hypertension: Was There a J-Curve for Ambulatory Blood Pressures?. <i>Hypertension</i> , 2021, 77, 1895-1905.  | 1.3 | 5         |
| 10 | Importance of hematological parameters for micro- and macrovascular outcomes in patients with type 2 diabetes: the Rio de Janeiro type 2 diabetes cohort study. <i>Cardiovascular Diabetology</i> , 2021, 20, 133.   | 2.7 | 16        |
| 11 | Prognostic impact of liver fibrosis and steatosis by transient elastography for cardiovascular and mortality outcomes in individuals with nonalcoholic fatty liver disease and type 2 diabetes: the Rio de Janeiro Cohort Study. <i>Cardiovascular Diabetology</i> , 2021, 20, 193.                                  | 2.7 | 17        |
| 12 | Prognostic Impact of Home Blood Pressures for Adverse Cardiovascular Outcomes and Mortality in Patients With Resistant Hypertension: A Prospective Cohort Study. <i>Hypertension</i> , 2021, 78, 1617-1627.  | 1.3 | 10        |
| 13 | Effect of Continuous Positive Airway Pressure on Weight and Local Adiposity in Adults with Obstructive Sleep Apnea: A Meta-Analysis. <i>Annals of the American Thoracic Society</i> , 2021, 18, 1717-1727.   | 1.5 | 12        |
| 14 | Prognostic impact of short-term ambulatory blood pressure variability for microvascular and macrovascular outcomes in patients with type 2 diabetes: the Rio de Janeiro Type 2 Diabetes Cohort Study. <i>Journal of Hypertension</i> , 2021, 39, 935-946.  | 0.3 | 4         |
| 15 | Prognostic Importance of Resistant Hypertension in Patients With Type 2 Diabetes: The Rio de Janeiro Type 2 Diabetes Cohort Study. <i>Diabetes Care</i> , 2020, 43, 219-227.   | 4.3 | 14        |
| 16 | Reply to Drs Mantovani and Zusi. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020, 30, 164-166.   | 1.1 | 0         |
| 17 | Refractory Hypertension and Risks of Adverse Cardiovascular Events and Mortality in Patients With Resistant Hypertension: A Prospective Cohort Study. <i>Journal of the American Heart Association</i> , 2020, 9, e017634.   | 1.6 | 20        |
| 18 | Prognostic importance of visit-to-visit blood pressure variability for micro- and macrovascular outcomes in patients with type 2 diabetes: The Rio de Janeiro Type 2 Diabetes Cohort Study. <i>Cardiovascular Diabetology</i> , 2020, 19, 50.  | 2.7 | 16        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | Prognostic Importance of On-Treatment Clinic and Ambulatory Blood Pressures in Resistant Hypertension. <i>Hypertension</i> , 2020, 75, 1184-1194.   | 1.3 | 17        |
| 20 | Effects of continuous positive airway pressure treatment on aortic stiffness in patients with resistant hypertension and obstructive sleep apnea: A randomized controlled trial. <i>Journal of Sleep Research</i> , 2020, 29, e12990.                                       | 1.7 | 12        |
| 21 | PNPLA3 gene polymorphism in Brazilian patients with type 2 diabetes: A prognostic marker beyond liver disease?. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2019, 29, 965-971.   | 1.1 | 15        |
| 22 | Efficacy of diacerein in reducing liver steatosis and fibrosis in patients with type 2 diabetes and non-alcoholic fatty liver disease: A randomized, placebo-controlled trial. <i>Diabetes, Obesity and Metabolism</i> , 2019, 21, 1266-1270.                               | 2.2 | 25        |
| 23 | Prognostic impact of carotid intima-media thickness and carotid plaques on the development of micro- and macrovascular complications in individuals with type 2 diabetes: the Rio de Janeiro type 2 diabetes cohort study. <i>Cardiovascular Diabetology</i> , 2019, 18, 2. | 2.7 | 37        |
| 24 | Rebuttal From Drs Cardoso and Salles. <i>Chest</i> , 2019, 155, 1102-1103.  | 0.4 | 0         |
| 25 | COUNTERPOINT: Should Sleep Studies Be Performed for All Patients With Poorly Controlled Hypertension? No. <i>Chest</i> , 2019, 155, 1097-1101.  | 0.4 | 4         |
| 26 | Prognostic Impact of Aortic Stiffness in Patients With Resistant Hypertension. <i>Hypertension</i> , 2019, 73, 728-735.   | 1.3 | 21        |
| 27 | Long-term visit-to-visit glycemic variability as predictor of micro- and macrovascular complications in patients with type 2 diabetes: The Rio de Janeiro Type 2 Diabetes Cohort Study. <i>Cardiovascular Diabetology</i> , 2018, 17, 33.                                   | 2.7 | 134       |
| 28 | Aortic stiffness and ambulatory blood pressure as predictors of diabetic kidney disease: a competing risks analysis from the Rio de Janeiro Type 2 Diabetes Cohort Study. <i>Diabetologia</i> , 2018, 61, 455-465.  | 2.9 | 26        |
| 29 | Treatment Adherence and Its Associated Factors in Patients with Type 2 Diabetes: Results from the Rio de Janeiro Type 2 Diabetes Cohort Study. <i>Journal of Diabetes Research</i> , 2018, 2018, 1-8.   | 1.0 | 46        |
| 30 | Prognostic impact of the ankle-brachial index on the development of micro- and macrovascular complications in individuals with type 2 diabetes: the Rio de Janeiro Type 2 Diabetes Cohort Study. <i>Diabetologia</i> , 2018, 61, 2266-2276.                                 | 2.9 | 15        |
| 31 | Effects of continuous positive airway pressure treatment on aldosterone excretion in patients with obstructive sleep apnoea and resistant hypertension. <i>Journal of Hypertension</i> , 2017, 35, 837-844.   | 0.3 | 23        |
| 32 | Polymorphisms of Leptin (G2548A) and Leptin Receptor (Q223R and K109R) Genes and Blood Pressure During Pregnancy and the Postpartum Period: A Cohort. <i>American Journal of Hypertension</i> , 2017, 30, 130-140.  | 1.0 | 9         |
| 33 | Efficacy and Safety of Diacerein in Patients With Inadequately Controlled Type 2 Diabetes: A Randomized Controlled Trial. <i>Diabetes Care</i> , 2017, 40, 1356-1363.   | 4.3 | 24        |
| 34 | Predictors of Development and Progression of Retinopathy in Patients with Type 2 Diabetes: Importance of Blood Pressure Parameters. <i>Scientific Reports</i> , 2017, 7, 4867.  | 1.6 | 50        |
| 35 | NAFLD and Increased Aortic Stiffness: Parallel or Common Physiopathological Mechanisms?. <i>International Journal of Molecular Sciences</i> , 2016, 17, 460.  | 1.8 | 33        |
| 36 | Aortic Stiffness as a Surrogate Endpoint to Micro- and Macrovascular Complications in Patients with Type 2 Diabetes. <i>International Journal of Molecular Sciences</i> , 2016, 17, 2044.   | 1.8 | 41        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 37 | Increasing aortic stiffness is predictive of advanced liver fibrosis in patients with type 2 diabetes: the Rio de Janeiro Type 2 Diabetes Cohort Study. <i>Liver International</i> , 2016, 36, 977-985.  | 1.9 | 19        |
| 38 | Prognostic Importance of C-reactive Protein in High Cardiovascular Risk Patients With Type 2 Diabetes Mellitus: The Rio de Janeiro Type 2 Diabetes Cohort Study. <i>Journal of the American Heart Association</i> , 2016, 5, .                                 | 1.6 | 22        |
| 39 | Are 2 Years Enough? Exploring Technical Skills Acquisition Among General Surgery Residents in Brazil. <i>Teaching and Learning in Medicine</i> , 2016, 28, 260-268.  | 1.3 | 1         |
| 40 | Prevalence of subclinical hypercortisolism in type 2 diabetic patients from the Rio de Janeiro Type 2 Diabetes Cohort Study. <i>Journal of Diabetes and Its Complications</i> , 2016, 30, 1032-1038.   | 1.2 | 11        |
| 41 | Aortic stiffness: is it time to be included into clinical diabetes management?. <i>Journal of Diabetes and Its Complications</i> , 2016, 30, 1207-1208.  | 1.2 | 3         |
| 42 | Prognostic Importance of Ambulatory Blood Pressure Monitoring in Resistant Hypertension: Is It All that Matters?. <i>Current Hypertension Reports</i> , 2016, 18, 85.  | 1.5 | 14        |
| 43 | Profile of disabilities and their associated factors in patients with type 2 diabetes evaluated by the Canadian occupational performance measure: the Rio De Janeiro type 2 diabetes cohort study. <i>Disability and Rehabilitation</i> , 2016, 38, 2095-2101. | 0.9 | 12        |
| 44 | Prognostic Effect of the Nocturnal Blood Pressure Fall in Hypertensive Patients. <i>Hypertension</i> , 2016, 67, 693-700.  | 1.3 | 399       |
| 45 | Correlates of aortic stiffness progression in patients with resistant hypertension. <i>Journal of Hypertension</i> , 2015, 33, 827-835.  | 0.3 | 15        |
| 46 | Nitroglycerin-mediated, but not flow-mediated vasodilation, is associated with blunted nocturnal blood pressure fall in patients with resistant hypertension. <i>Journal of Hypertension</i> , 2015, 33, 1666-1675.  | 0.3 | 15        |
| 47 | Increased aortic stiffness predicts future development and progression of peripheral neuropathy in patients with type 2 diabetes: the Rio de Janeiro Type 2 Diabetes Cohort Study. <i>Diabetologia</i> , 2015, 58, 2161-2168.                                  | 2.9 | 42        |
| 48 | Is Home Blood Pressure Monitoring Useful in the Management of Patients With Resistant Hypertension?. <i>American Journal of Hypertension</i> , 2015, 28, 190-199.  | 1.0 | 21        |
| 49 | Blood Pressure in Healthy Pregnancy and Factors Associated With No Mid-Trimester Blood Pressure Drop: A Prospective Cohort Study. <i>American Journal of Hypertension</i> , 2015, 28, 680-689.   | 1.0 | 20        |
| 50 | Correlates of Aortic Stiffness Progression in Patients With Type 2 Diabetes: Importance of Glycemic Control. <i>Diabetes Care</i> , 2015, 38, 897-904.   | 4.3 | 51        |
| 51 | Development of a risk score to predict sudden death in patients with Chaga's heart disease. <i>International Journal of Cardiology</i> , 2015, 187, 700-704.   | 0.8 | 48        |
| 52 | Prognostic impact of sex-specific ambulatory blood pressure interactions in 10 cohorts of 17,312 patients diagnosed with hypertension. <i>Journal of Hypertension</i> , 2015, 33, 212-220.   | 0.3 | 23        |
| 53 | Effects of Continuous Positive Airway Pressure Treatment on Clinic and Ambulatory Blood Pressures in Patients With Obstructive Sleep Apnea and Resistant Hypertension. <i>Hypertension</i> , 2015, 65, 736-742.  | 1.3 | 126       |
| 54 | Prognostic importance of baseline and serial glycosylated hemoglobin levels in high-risk patients with type 2 diabetes: the Rio de Janeiro Type 2 Diabetes Cohort Study. <i>Acta Diabetologica</i> , 2015, 52, 21-29.  | 1.2 | 22        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 55 | Non-alcoholic fatty liver disease and diabetes: From physiopathological interplay to diagnosis and treatment. <i>World Journal of Gastroenterology</i> , 2014, 20, 8377.                               | 1.4 | 76        |
| 56 | Diagnostic accuracy of the Berlin questionnaire in detecting obstructive sleep apnea in patients with resistant hypertension. <i>Journal of Hypertension</i> , 2014, 32, 2030-2037.                    | 0.3 | 30        |
| 57 | Prognostic impact from clinic, daytime, and night-time systolic blood pressure in nine cohorts of 13,844 patients with hypertension. <i>Journal of Hypertension</i> , 2014, 32, 2332-2340.             | 0.3 | 222       |
| 58 | Cardiovascular and Renal Complications in Patients with Resistant Hypertension. <i>Current Hypertension Reports</i> , 2014, 16, 471.   | 1.5 | 24        |
| 59 | Prevalence and Associated Factors of Obstructive Sleep Apnea in Patients with Resistant Hypertension. <i>American Journal of Hypertension</i> , 2014, 27, 1069-1078.                                   | 1.0 | 117       |
| 60 | Relationships between reduced heart rate variability and pre-clinical cardiovascular disease in patients with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2014, 106, 110-117.    | 1.1 | 18        |
| 61 | A reduced heart rate variability is independently associated with a blunted nocturnal blood pressure fall in patients with resistant hypertension. <i>Journal of Hypertension</i> , 2014, 32, 644-651. | 0.3 | 25        |
| 62 | Prognostic Impact of Aortic Stiffness in High-Risk Type 2 Diabetic Patients. <i>Diabetes Care</i> , 2013, 36, 3772-3778.   | 4.3 | 93        |
| 63 | Factors associated with abnormal T-wave axis and increased QRS-T angle in type 2 diabetes. <i>Acta Diabetologica</i> , 2013, 50, 919-925.  | 1.2 | 12        |
| 64 | How to use ambulatory blood pressure monitoring in resistant hypertension. <i>Hypertension Research</i> , 2013, 36, 385-389.   | 1.5 | 25        |
| 65 | Prognostic impact of clinic and ambulatory blood pressure components in high-risk type 2 diabetic patients. <i>Journal of Hypertension</i> , 2013, 31, 2176-2186.                                      | 0.3 | 37        |
| 66 | Prognostic Significance of Baseline Heart Rate and Its Interaction With Beta-Blocker Use in Resistant Hypertension: A Cohort Study. <i>American Journal of Hypertension</i> , 2013, 26, 218-226.       | 1.0 | 23        |
| 67 | Appropriate Time Interval to Repeat Ambulatory Blood Pressure Monitoring in Patients With White-Coat Resistant Hypertension. <i>Hypertension</i> , 2012, 59, 384-389.                                  | 1.3 | 47        |
| 68 | Thresholds of Ambulatory Blood Pressure Associated With Chronic Complications in Type 2 Diabetes. <i>American Journal of Hypertension</i> , 2012, 25, 82-88.   | 1.0 | 19        |
| 69 | Prognostic factors in resistant hypertension: implications for cardiovascular risk stratification and therapeutic management. <i>Expert Review of Cardiovascular Therapy</i> , 2012, 10, 735-745.      | 0.6 | 20        |
| 70 | Factors associated with carotid intima-media thickness and carotid plaques in type 2 diabetic patients. <i>Journal of Hypertension</i> , 2012, 30, 940-947.  | 0.3 | 28        |
| 71 | Prevalence and associated factors of subclinical hypercortisolism in patients with resistant hypertension. <i>Journal of Hypertension</i> , 2012, 30, 967-973.   | 0.3 | 37        |
| 72 | o posicionamento brasileiro sobre hipertenso arterial resistente. <i>Arquivos Brasileiros De Cardiologia</i> , 2012, 99, 576-585.   | 0.3 | 27        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 73 | Prognostic importance of baseline and serial changes in microalbuminuria in patients with resistant hypertension. <i>Atherosclerosis</i> , 2011, 216, 199-204.   | 0.4 | 32        |
| 74 | Prognostic significance of a reduced glomerular filtration rate and interaction with microalbuminuria in resistant hypertension. <i>Journal of Hypertension</i> , 2011, 29, 2014-2023.                 | 0.3 | 35        |
| 75 | Histopathological stages of nonalcoholic fatty liver disease in type 2 diabetes: prevalences and correlated factors. <i>Liver International</i> , 2011, 31, 700-706.                                   | 1.9 | 151       |
| 76 | Prognostic significance of baseline and serial changes in electrocardiographic strain pattern in resistant hypertension. <i>Journal of Hypertension</i> , 2010, 28, 1715-1723.                         | 0.3 | 16        |
| 77 | Prognostic impact of the ambulatory arterial stiffness index in resistant hypertension. <i>Journal of Hypertension</i> , 2010, 28, 1547-1553.  | 0.3 | 42        |
| 78 | Efficacy of Spironolactone Therapy in Patients With True Resistant Hypertension. <i>Hypertension</i> , 2010, 55, 147-152.  | 1.3 | 217       |
| 79 | Uncontrolled isolated office hypertension is associated with subclinical markers of cardiovascular disease in hypertensive type 2 diabetic patients. <i>Hypertension Research</i> , 2010, 33, 819-824. | 1.5 | 7         |
| 80 | Prognostic impact of baseline and serial changes in electrocardiographic left ventricular hypertrophy in resistant hypertension. <i>American Heart Journal</i> , 2010, 159, 833-840.                   | 1.2 | 35        |
| 81 | A blunted decrease in nocturnal blood pressure is independently associated with increased aortic stiffness in patients with resistant hypertension. <i>Hypertension Research</i> , 2009, 32, 591-596.  | 1.5 | 45        |
| 82 | Prevalence and associated factors of nonalcoholic fatty liver disease in patients with type 2 diabetes mellitus. <i>Liver International</i> , 2009, 29, 113-119.                                       | 1.9 | 427       |
| 83 | Microvascular degenerative complications are associated with increased aortic stiffness in type 2 diabetic patients. <i>Atherosclerosis</i> , 2009, 205, 472-476.                                      | 0.4 | 67        |
| 84 | Prognostic value of ventricular repolarization prolongation in resistant hypertension: a prospective cohort study. <i>Journal of Hypertension</i> , 2009, 27, 1094-1101.                               | 0.3 | 25        |
| 85 | Predictors of development and progression of microvascular complications in a cohort of Brazilian type 2 diabetic patients. <i>Journal of Diabetes and Its Complications</i> , 2008, 22, 164-170.      | 1.2 | 32        |
| 86 | Pattern of 24-Hour Ambulatory Blood Pressure Monitoring in Type 2 Diabetic Patients with Cardiovascular Dysautonomy. <i>Hypertension Research</i> , 2008, 31, 865-872.                                 | 1.5 | 38        |
| 87 | Recent Ventricular Repolarization Markers in Resistant Hypertension: Are They Different from the Traditional QT Interval?. <i>American Journal of Hypertension</i> , 2008, 21, 47-53.                  | 1.0 | 37        |
| 88 | Ambulatory Arterial Stiffness Index or Pulse Pressure: Which Correlates Better with Arterial Stiffness in Resistant Hypertension?. <i>Hypertension Research</i> , 2008, 31, 607-613.                   | 1.5 | 40        |
| 89 | Prognostic Influence of Office and Ambulatory Blood Pressures in Resistant Hypertension. <i>Archives of Internal Medicine</i> , 2008, 168, 2340.   | 4.3 | 264       |
| 90 | Pulse pressure or dipping pattern: which one is a better cardiovascular risk marker in resistant hypertension?. <i>Journal of Hypertension</i> , 2008, 26, 878-884.                                    | 0.3 | 41        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 91  | Relation of Left Ventricular Hypertrophy With Systemic Inflammation and Endothelial Damage in Resistant Hypertension. <i>Hypertension</i> , 2007, 50, 723-728.  | 1.3 | 81        |
| 92  | Electrocardiographic Strain Pattern and Cardiovascular Prognosis in Hypertension. <i>American Journal of Hypertension</i> , 2007, 20, 1005-1006.  | 1.0 | 0         |
| 93  | Macro and microvascular complications are determinants of increased infection-related mortality in Brazilian type 2 diabetes mellitus patients. <i>Diabetes Research and Clinical Practice</i> , 2007, 75, 51-58.                     | 1.1 | 20        |
| 94  | Importance of the Electrocardiographic Strain Pattern in Patients With Resistant Hypertension. <i>Hypertension</i> , 2006, 48, 437-442.   | 1.3 | 35        |
| 95  | Combined QT Interval and Voltage Criteria Improve Left Ventricular Hypertrophy Detection in Resistant Hypertension. <i>Hypertension</i> , 2005, 46, 1207-1212.  | 1.3 | 39        |
| 96  | Mortality and Predictors of Mortality in a Cohort of Brazilian Type 2 Diabetic Patients. <i>Diabetes Care</i> , 2004, 27, 1299-1305.  | 4.3 | 51        |
| 97  | T-wave axis deviation as an independent predictor of mortality in chronic Chagas' disease. <i>American Journal of Cardiology</i> , 2004, 93, 1136-1140.   | 0.7 | 34        |
| 98  | Prognostic value of QT interval parameters in type 2 diabetes mellitus. <i>Journal of Diabetes and Its Complications</i> , 2003, 17, 169-178.   | 1.2 | 70        |
| 99  | Electrocardiographic Ventricular Repolarization Parameters in Chronic Chagas' Disease as Predictors of Asymptomatic Left Ventricular Systolic Dysfunction. <i>PACE - Pacing and Clinical Electrophysiology</i> , 2003, 26, 1326-1335. | 0.5 | 20        |
| 100 | Prognostic Value of QT Interval Parameters for Mortality Risk Stratification in Chagas's Disease. <i>Circulation</i> , 2003, 108, 305-312.  | 1.6 | 125       |
| 101 | QTc Interval Prolongation Is a Predictor of Future Strokes in Patients With Type 2 Diabetes Mellitus. <i>Stroke</i> , 2003, 34, 2187-2194.  | 1.0 | 44        |
| 102 | Twenty-four hour ambulatory blood pressure monitoring pattern of resistant hypertension. <i>Blood Pressure Monitoring</i> , 2003, 8, 181-185.   | 0.4 | 113       |
| 103 | Clinical determinants of increased QT dispersion in patients with diabetes mellitus. <i>International Journal of Cardiology</i> , 2001, 79, 253-262.  | 0.8 | 46        |