

# Christoph HÃ¶rweg

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6323303/publications.pdf>

Version: 2024-02-01

13  
papers

98  
citations

1478505

6  
h-index

1372567

10  
g-index

22  
all docs

22  
docs citations

22  
times ranked

121  
citing authors

#	ARTICLE	IF	CITATIONS
1	Wherefrom and whereabouts of an alien: the American liver fluke <i>Fascioloides magna</i> in Austria: an overview. <i>Wiener Klinische Wochenschrift</i> , 2014, 126, 23-31.	1.9	18
2	Recovery of <i>Fascioloides magna</i> (Digenea) population in spite of treatment programme? Screening of <i>Galba truncatula</i> (Gastropoda, Lymnaeidae) from Lower Austria. <i>Veterinary Parasitology</i> , 2012, 187, 445-451.	1.8	13
3	A new species of the South East Asian genus <i>Sarax</i> Simon, 1892 (Arachnida: Amblypygi: Charinidae) and synonymization of <i>Sarax mediterraneus</i> Delle Cave, 1986. <i>Zootaxa</i> , 2015, 4012, 542-52.	0.5	13
4	Cercarial dermatitis in Austria: Questionnaires as useful tools to estimate risk factors?. <i>Wiener Klinische Wochenschrift</i> , 2006, 118, 77-80.	1.9	12
5	Genetic homogeneity of <i>Fascioloides magna</i> in Austria. <i>Veterinary Parasitology</i> , 2017, 243, 75-78.	1.8	8
6	First Record of <i>Trichobilharzia physellae</i> (Talbot, 1936) in Europe, a Possible Causative Agent of Cercarial Dermatitis. <i>Pathogens</i> , 2021, 10, 1473.	2.8	8
7	The whip spider collection (Arachnida, Amblypygi) held in the Natural History Museum Vienna, Austria. <i>Arachnologische Mitteilungen</i> , 2013, 46, 47-53.	0.3	6
8	Reconditioning of the Nalepa collection of eriophyoid mites (Acariformes, Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50,462 Td (Er	0.5	6
9	First confirmed record of <i>Trichobilharzia franki</i> MÄ¼ller & Kimmig, 1994, from <i>Radix auricularia</i> (Linnaeus, 1758) for Austria. <i>Parasitology Research</i> , 2020, 119, 4135-4141.	1.6	5
10	Supplementary description of three species from the subfamily Cecidophyinae (Eriophyoidea: Eriophyidae) from the Nalepa collection. <i>Systematic and Applied Acarology</i> , 2018, 23, 838.	0.5	3
11	Redescriptions of three species of the genus <i>Neobisium</i> from the Balkans and the Carpathians (Arachnida: Pseudoscorpiones). <i>Acta Zoologica Academiae Scientiarum Hungaricae</i> , 2017, 63, 277-292.	0.5	3
12	Supplementary descriptions of seven eriophyoid mite species (Acari: Eriophyoidea) recovered from the Viennese Nalepa collection and comparison with Japanese species. <i>Acarologia</i> , 2022, 62, 273-301.	0.6	1
13	Synanthropic is best: <i>Nuctenea umbratica</i> (Araneae: Araneidae) and <i>Steatoda bipunctata</i> (Araneae:) Tj ETQq1 1 0.784314 rgBT /Overl 56, 32-35.	0.3	0