Per-Olof Hansson

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/633032/publications.pdf

Version: 2024-02-01

		430754	276775
58	1,863	18	41
papers	citations	h-index	g-index
60	60	60	2103
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	SCORE2 risk prediction algorithms: new models to estimate 10-year risk of cardiovascular disease in Europe. European Heart Journal, 2021, 42, 2439-2454.	1.0	491
2	Smoking and Abdominal Obesity. Archives of Internal Medicine, 1999, 159, 1886.	4.3	269
3	Deep Vein Thrombosis and Pulmonary Embolism in the General Population. Archives of Internal Medicine, 1997, 157, 1665.	4. 3	264
4	Ischemic Stroke in Children and Young Adults With Congenital Heart Disease. Journal of the American Heart Association, $2016, 5, \ldots$	1.6	81
5	Adherence of self-monitoring of blood glucose in persons with type 1 diabetes in Sweden. BMJ Open Diabetes Research and Care, 2017, 5, e000342.	1.2	70
6	Atrial Fibrillation Burden in Young Patients With Congenital Heart Disease. Circulation, 2018, 137, 928-937.	1.6	67
7	Long-Term Risk of Hemorrhagic Stroke in Young Patients With Congenital Heart Disease. Stroke, 2018, 49, 1155-1162.	1.0	38
8	Low aerobic capacity in middle-aged men associated with increased mortality rates during 45 years of follow-up. European Journal of Preventive Cardiology, 2016, 23, 1557-1564.	0.8	37
9	Impact of changes in heart rate with age on all-cause death and cardiovascular events in 50-year-old men from the general population. Open Heart, 2019, 6, e000856.	0.9	37
10	Risk of falling in a stroke unit after acute stroke: The Fall Study of Gothenburg (FallsGOT). Clinical Rehabilitation, 2018, 32, 398-409.	1.0	34
11	Secular changes in cardiovascular risk factors in Swedish 50-year-old men over a 50-year period: The study of men born in 1913, 1923, 1933, 1943, 1953 and 1963. European Journal of Preventive Cardiology, 2017, 24, 612-620.	0.8	31
12	The risk of atrial fibrillation in the general male population: a lifetime follow-up of 50-year-old men. Europace, 2015, 17, 1018-1022.	0.7	28
13	Association between left atrial enlargement and obstructive sleep apnea in a general population of 71â€yearâ€old men. Journal of Sleep Research, 2018, 27, 254-260.	1.7	27
14	Early prediction of falls after stroke: a 12-month follow-up of 490 patients in The Fall Study of Gothenburg (FallsGOT). Clinical Rehabilitation, 2019, 33, 773-783.	1.0	27
15	Fear of falling in acute stroke: The Fall Study of Gothenburg (FallsGOT). Topics in Stroke Rehabilitation, 2018, 25, 256-260.	1.0	24
16	Risk factors for subarachnoid haemorrhage: a nationwide cohort of 950Â000 adults. International Journal of Epidemiology, 2019, 48, 2018-2025.	0.9	21
17	Physical Activity Levels and Their Associations With Postural Control in the First Year After Stroke. Physical Therapy, 2016, 96, 1389-1396.	1.1	19
18	The clinical consequences of a pre-hospital diagnosis of stroke by the emergency medical service system. A pilot study. Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine, 2012, 20, 48.	1.1	18

#	Article	IF	CITATIONS
19	mHealth Self-Report Monitoring in Competitive Middle- and Long-Distance Runners: Qualitative Study of Long-Term Use Intentions Using the Technology Acceptance Model. JMIR MHealth and UHealth, 2018, 6, e10270.	1.8	17
20	Although Coronary Mortality Has Decreased, Rates of Cardiovascular Disease Remain High: 21 Years of Followâ€Up Comparing Cohorts of Men Born in 1913 With Men Born in 1943. Journal of the American Heart Association, 2018, 7, .	1.6	15
21	Atrial natriuretic peptide as a predictor of atrial fibrillation in a male population study. The Study of Men Born in 1913 and 1923. International Journal of Cardiology, 2014, 171, 44-48.	0.8	13
22	Lung function, functional capacity, and respiratory symptoms at discharge from hospital in patients with acute pulmonary embolism: A cross-sectional study. Physiotherapy Theory and Practice, 2018, 34, 194-201.	0.6	13
23	Prevalence and risk factors of aortic stenosis and aortic sclerosis: a 21-year follow-up of middle-aged men. Scandinavian Cardiovascular Journal, 2020, 54, 115-123.	0.4	13
24	Obesity in adolescent men increases the risk of venous thromboembolism in adult life. Journal of Internal Medicine, 2020, 287, 734-745.	2.7	13
25	Recurrent stroke in patients with patent foramen ovale: An observational prospective study of percutaneous closure of PFO versus non-closure. International Journal of Cardiology, 2015, 195, 293-299.	0.8	12
26	Genetic variation at the human connexin 43 locus but not at the connexin 40 locus is associated with left bundle branch block. Open Heart, 2015, 2, e000187.	0.9	11
27	Pulmonary embolism and deep vein thrombosisâ€"comorbidities and temporary provoking factors in a registerâ€based study of 1.48 million people. Research and Practice in Thrombosis and Haemostasis, 2022, 6, e12714.	1.0	11
28	Quality of life after percutaneous closure of patent foramen ovale in patients after cryptogenic stroke compared to a normative sample. International Journal of Cardiology, 2018, 257, 46-49.	0.8	9
29	Prehospital assessment of suspected stroke and TIA: An observational study. Acta Neurologica Scandinavica, 2019, 140, 93-99.	1.0	8
30	High prevalence of cardiac dysfunction or overt heart failure in 71-year-old men: A 21-year follow-up of "The Study of men born in 1943― European Journal of Preventive Cardiology, 2020, 27, 717-725.	0.8	8
31	Determinants of Recurrent Falls Poststroke: A 1-Year Follow-up of the Fall Study of Gothenburg. Archives of Physical Medicine and Rehabilitation, 2020, 101, 1541-1548.	0.5	8
32	Physical activity and respiratory symptoms after pulmonary embolism. A longitudinal observational study. Thrombosis Research, 2020, 189, 55-60.	0.8	8
33	Atrial fibrillation in the elderly general population: a 30-year follow-up from 70 to 100 years of age. Scandinavian Cardiovascular Journal, 2020, 54, 232-238.	0.4	8
34	Determinants of Stroke in a General Male Population. Stroke, 2018, 49, 2830-2836.	1.0	7
35	Prehospital identification of factors associated with death during oneâ€year followâ€up after acute stroke. Brain and Behavior, 2018, 8, e00987.	1.0	7
36	Hyperparathyroidism in men – morbidity and mortality during 21 years' follow-up. Scandinavian Journal of Clinical and Laboratory Investigation, 2020, 80, 6-13.	0.6	7

#	Article	IF	CITATIONS
37	Prehospital recognition of stroke is associated with a lower risk of death. Acta Neurologica Scandinavica, 2022, 146, 126-136.	1.0	7
38	Determinants of falls after stroke based on data on 5065 patients from the Swedish VÃststroke and Riksstroke Registers. Scientific Reports, 2021, 11, 24035.	1.6	7
39	Drug Treatment, Postural Control, and Falls: An Observational Cohort Study of 504 Patients With Acute Stroke, the Fall Study of Gothenburg. Archives of Physical Medicine and Rehabilitation, 2019, 100, 1267-1273.	0.5	6
40	Prediction of fear of falling at $6\text{\^{A}}$ months after stroke based on 279 individuals from the Fall Study of Gothenburg. Scientific Reports, 2021, 11, 13503.	1.6	6
41	Atrial fibrillation and risk of venous thromboembolism: a Swedish Nationwide Registry Study. Europace, 2021, 23, 1913-1921.	0.7	6
42	Long-term risk of stroke and myocardial infarction in middle-aged men with a hypertensive response to exercise: a 44-year follow-up study. Journal of Hypertension, 2021, 39, 503-510.	0.3	6
43	Seven-year follow-up of percutaneous closure of patent foramen ovale. International Journal of Cardiology Heart & Vessels, 2013, 1, 32-36.	0.5	5
44	The incidence of atrial fibrillation and the added value of thumb ECG for detecting new cases. Scandinavian Cardiovascular Journal, 2018, 52, 256-261.	0.4	5
45	Natriuretic and Inflammatory Biomarkers as Risk Predictors of Heart Failure in Middle-Aged Men From the General Population: A 21-Year Follow-Up. Journal of Cardiac Failure, 2018, 24, 594-600.	0.7	5
46	Cardiovascular risk factors in relation to dietary patterns in 50-year-old men and women: a feasibility study of a short FFQ. Public Health Nutrition, 2019, 22, 645-653.	1.1	5
47	A negative Tâ€wave in electrocardiogram at 50 years predicted lifetime mortality in a random populationâ€based cohort. Clinical Cardiology, 2020, 43, 1279-1285.	0.7	5
48	High-normal blood pressure conferred higher risk of cardiovascular disease in a random population sample of 50-year-old men. Medicine (United States), 2020, 99, e19895.	0.4	5
49	SBP and antihypertensive treatment in the acute phase after stroke and its impact on the risk of falling. Journal of Hypertension, 2019, 37, 1032-1039.	0.3	4
50	Incremental changes in QRS duration as predictor for cardiovascular disease: a 21-year follow-up of a randomly selected general population. Scientific Reports, 2021, 11, 13652.	1.6	4
51	Rationale for a Swedish cohort consortium. Upsala Journal of Medical Sciences, 2019, 124, 21-28.	0.4	3
52	Secular trends in cardiovascular risk factors among women aged 45–54 years in Gothenburg, Sweden, from 1980 to 2014. BMC Public Health, 2020, 20, 1042.	1.2	3
53	The impact of time-updated resting heart rate on cause-specific mortality in a random middle-aged male population: a lifetime follow-up. Clinical Research in Cardiology, 2021, 110, 822-830.	1.5	3
54	Multi-modality biomarkers in the early prediction of ischaemic heart disease in middle-aged men during a 21-year follow-up. BMC Cardiovascular Disorders, 2021, 21, 65.	0.7	3

#	Article	IF	CITATIONS
55	Body iron stores had no impact on coronary heart disease outcomes: a middle-aged male cohort from the general population with 21-year follow-up. Open Heart, 2022, 9, e001928.	0.9	3
56	Lifetime risk of stroke in the general male population. Acta Neurologica Scandinavica, 2020, 142, 30-36.	1.0	0
57	Abstract WP197: Recurrent Falls After Stroke: A One-Year Follow-Up of the Fall Study of Gothenburg. Stroke, 2019, 50, .	1.0	O
58	Midâ€life extrapyramidal symptoms predict cognitive impairment 23 years later. Acta Neurologica Scandinavica, 2022, 145, 305-313.	1.0	0