

# Ryota Asahina

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/6333885/ryota-asahina-publications-by-citations.pdf>

**Version:** 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

13  
papers

60  
citations

3  
h-index

7  
g-index

14  
ext. papers

96  
ext. citations

1.7  
avg, IF

2.47  
L-index

#	Paper	IF	Citations
13	A review of the roles of keratinocyte-derived cytokines and chemokines in the pathogenesis of atopic dermatitis in humans and dogs. <i>Veterinary Dermatology</i> , <b>2017</b> , 28, 16-e5	1.8	31
12	Transcription of thymic stromal lymphopoietin via Toll-like receptor 2 in canine keratinocytes: a possible association of <i>Staphylococcus</i> spp. in the deterioration of allergic inflammation in canine atopic dermatitis. <i>Veterinary Dermatology</i> , <b>2016</b> , 27, 184-e46	1.8	8
11	Gene transcription of pro-inflammatory cytokines and chemokines induced by IL-17A in canine keratinocytes. <i>Veterinary Dermatology</i> , <b>2015</b> , 26, 426-31, e100	1.8	6
10	A review of the roles of keratinocyte-derived cytokines and chemokines in the pathogenesis of atopic dermatitis in humans and dogs <b>2017</b> , 15-25		3
9	Expression of IL-33 in chronic lesional skin of canine atopic dermatitis. <i>Veterinary Dermatology</i> , <b>2018</b> , 29, 246-e91	1.8	3
8	Expression of ZO-1 and claudin-1 in a 3D epidermal equivalent using canine progenitor epidermal keratinocytes. <i>Veterinary Dermatology</i> , <b>2018</b> , 29, 288	1.8	3
7	Th17 cells increase during maturation in peripheral blood of healthy dogs. <i>Veterinary Immunology and Immunopathology</i> , <b>2019</b> , 209, 17-21	2	2
6	Characterization of a novel canine T-cell line established from a dog with cutaneous T-cell lymphoma. <i>Journal of Dermatological Science</i> , <b>2017</b> , 88, 254-256	4.3	1
5	Serum canine thymus and activation-regulated chemokine (TARC/CCL17) concentrations correlate with disease severity and therapeutic responses in dogs with atopic dermatitis. <i>Veterinary Dermatology</i> , <b>2020</b> , 31, 446-455	1.8	1
4	Transcriptional analysis of the IL-33 receptor suppression of tumourigenicity 2 and its effects on canine Type 2 T helper cells: a preliminary study. <i>Veterinary Dermatology</i> , <b>2018</b> , 29, 112-e45	1.8	1
3	Cutaneous Liver X Receptor Activation Prevents the Formation of Imiquimod-Induced Psoriatic Dermatitis. <i>Journal of Investigative Dermatology</i> , <b>2021</b> ,	4.3	1
2	Phenotypic analysis of mice xenografted with canine epitheliotropic cutaneous T-cell lymphoma cells. <i>Veterinary Dermatology</i> , <b>2018</b> , 29, 517-e172	1.8	0
1	Narrow-band ultraviolet B therapy attenuates cutaneous T-cell responses in hapten-induced, experimental contact dermatitis in beagles. <i>Veterinary Dermatology</i> , <b>2021</b> , 32, 605-e161	1.8	