
RafaÅ, Ficek

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6334030/publications.pdf Version: 2024-02-01



RAFAL FICER

#	Article	IF	CITATIONS
1	Calcification of coronary arteries and abdominal aorta in relation to traditional and novel risk factors of atherosclerosis in hemodialysis patients. BMC Nephrology, 2013, 14, 10.	1.8	47
2	Relationship between plasma levels of zonulin, bacterial lipopolysaccharides, d-lactate and markers of inflammation in haemodialysis patients. International Urology and Nephrology, 2017, 49, 717-725.	1.4	26
3	Relationship between plasma levels of sclerostin, calcium–phosphate disturbances, established markers of bone turnover, and inflammation in haemodialysis patients. International Urology and Nephrology, 2019, 51, 519-526.	1.4	17
4	Functional deficiency of vitamin K in hemodialysis patients in Upper Silesia in Poland. International Urology and Nephrology, 2016, 48, 765-771.	1.4	12
5	Plasma Concentrations of Tumor Necrosis Factor Alpha May Predict the Outcome of Patients with Acute Renal Failure. Kidney and Blood Pressure Research, 2006, 29, 203-209.	2.0	11
6	Daily intake and serum concentration of menaquinone-4 (MK-4) in haemodialysis patients with chronic kidney disease. Clinical Biochemistry, 2015, 48, 1246-1251.	1.9	11
7	Diabetes decreases patency of tunneled catheters in hemodialysis patients after first effective thrombolysis with urokinase. Renal Failure, 2018, 40, 384-389.	2.1	7
8	Arterial stiffness but not endothelial dysfunction is associated with multidrug antihypertensive therapy and nondipper blood pressure pattern in kidney transplant recipients. Medicine (United) Tj ETQq0 0 0 rg	;BT1 /.O verlo	ock610 Tf 50 4
9	Bronchial asthma control after argon plasma coagulation turbinectomy in patients with chronic rhinitis. European Archives of Oto-Rhino-Laryngology, 2014, 271, 1581-1587.	1.6	5
10	Number of Regularly Prescribed Drugs and Intrapatient Tacrolimus Trough Levels Variability in Stable Kidney Transplant Recipients. Journal of Clinical Medicine, 2020, 9, 1926.	2.4	5
11	N-Terminal Prohormone of Brain Natriuretic Peptide but not C-Terminal Pre-Pro Vasopressin (Copeptin) Level is Associated with the Response to Antihypertensive Therapy in Haemodialysis Patients. Kidney and Blood Pressure Research, 2017, 42, 1013-1022.	2.0	4
12	Nutrient intake assessed with Diet History Questionnaire II, in relation to long-term calcium-phosphate control in hemodialysis patients with end-stage renal failure. Advances in Clinical and Experimental Medicine, 2018, 27, 217-224.	1.4	4

13	Guanylins – Are They of Nephrological Relevance?. Nephron, 2000, 84, 201-205.	1.8	1

Are plasma concentrations of leptin and carnitine interrelated in haemodialysis patients?. Nephrology 0.7 0 Dialysis Transplantation, 2001, 16, 434-435.