

Simon Gilbody

List of Publications by Citations

Source: <https://exaly.com/author-pdf/6337009/simon-gilbody-publications-by-citations.pdf>
Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

378 papers	19,754 citations	70 h-index	134 g-index
427 ext. papers	24,462 ext. citations	5.5 avg, IF	7.02 L-index

#	Paper	IF	Citations
378	Collaborative care for depression: a cumulative meta-analysis and review of longer-term outcomes. <i>Archives of Internal Medicine</i> , 2006 , 166, 2314-21		913
377	Optimal cut-off score for diagnosing depression with the Patient Health Questionnaire (PHQ-9): a meta-analysis. <i>Cmaj</i> , 2012 , 184, E191-6	3.5	892
376	Screening for depression in medical settings with the Patient Health Questionnaire (PHQ): a diagnostic meta-analysis. <i>Journal of General Internal Medicine</i> , 2007 , 22, 1596-602	4	767
375	Loneliness and social isolation as risk factors for coronary heart disease and stroke: systematic review and meta-analysis of longitudinal observational studies. <i>Heart</i> , 2016 , 102, 1009-16	5.1	650
374	Stepped care in psychological therapies: access, effectiveness and efficiency. Narrative literature review. <i>British Journal of Psychiatry</i> , 2005 , 186, 11-7	5.4	609
373	Educational and organizational interventions to improve the management of depression in primary care: a systematic review. <i>JAMA - Journal of the American Medical Association</i> , 2003 , 289, 3145-51	27.4	501
372	Collaborative care for depression and anxiety problems. <i>The Cochrane Library</i> , 2012 , 10, CD006525	5.2	451
371	Evidence-based guidelines for treating depressive disorders with antidepressants: A revision of the 2008 British Association for Psychopharmacology guidelines. <i>Journal of Psychopharmacology</i> , 2015 , 29, 459-525	4.6	381
370	Accuracy of Patient Health Questionnaire-9 (PHQ-9) for screening to detect major depression: individual participant data meta-analysis. <i>BMJ, The</i> , 2019 , 365, l1476	5.9	374
369	Efficacy of Self-guided Internet-Based Cognitive Behavioral Therapy in the Treatment of Depressive Symptoms: A Meta-analysis of Individual Participant Data. <i>JAMA Psychiatry</i> , 2017 , 74, 351-359	14.5	346
368	Screening and case-finding instruments for depression: a meta-analysis. <i>Cmaj</i> , 2008 , 178, 997-1003	3.5	312
367	A diagnostic meta-analysis of the Patient Health Questionnaire-9 (PHQ-9) algorithm scoring method as a screen for depression. <i>General Hospital Psychiatry</i> , 2015 , 37, 67-75	5.6	311
366	Systematic review and meta-analysis of transdiagnostic psychological treatments for anxiety and depressive disorders in adulthood. <i>Clinical Psychology Review</i> , 2015 , 40, 91-110	10.8	297
365	What makes self-help interventions effective in the management of depressive symptoms? Meta-analysis and meta-regression. <i>Psychological Medicine</i> , 2007 , 37, 1217-28	6.9	293
364	Collaborative care for depression in primary care. Making sense of a complex intervention: systematic review and meta-regression. <i>British Journal of Psychiatry</i> , 2006 , 189, 484-93	5.4	293
363	Cost and Outcome of Behavioural Activation versus Cognitive Behavioural Therapy for Depression (COBRA): a randomised, controlled, non-inferiority trial. <i>Lancet, The</i> , 2016 , 388, 871-80	4.0	282
362	A meta-analysis of randomized trials of behavioural treatment of depression. <i>Psychological Medicine</i> , 2008 , 38, 611-23	6.9	279

361	Barriers to the uptake of computerized cognitive behavioural therapy: a systematic review of the quantitative and qualitative evidence. <i>Psychological Medicine</i> , 2009 , 39, 705-12	6.9	267
360	Vitamin K and the prevention of fractures: systematic review and meta-analysis of randomized controlled trials. <i>Archives of Internal Medicine</i> , 2006 , 166, 1256-61		262
359	Publication and related biases.. <i>Health Technology Assessment</i> , 2000 , 4,	4.4	261
358	Behavioural activation for depression; an update of meta-analysis of effectiveness and sub group analysis. <i>PLoS ONE</i> , 2014 , 9, e100100	3.7	259
357	Methylenetetrahydrofolate reductase (MTHFR) genetic polymorphisms and psychiatric disorders: a HuGE review. <i>American Journal of Epidemiology</i> , 2007 , 165, 1-13	3.8	247
356	Computerised cognitive behaviour therapy (cCBT) as treatment for depression in primary care (REEACT trial): large scale pragmatic randomised controlled trial. <i>BMJ, The</i> , 2015 , 351, h5627	5.9	235
355	Routinely administered questionnaires for depression and anxiety: systematic review. <i>BMJ: British Medical Journal</i> , 2001 , 322, 406-9		227
354	The Hospital Anxiety and Depression Scale: a diagnostic meta-analysis of case-finding ability. <i>Journal of Psychosomatic Research</i> , 2010 , 69, 371-8	4.1	223
353	Influence of initial severity of depression on effectiveness of low intensity interventions: meta-analysis of individual patient data. <i>BMJ, The</i> , 2013 , 346, f540	5.9	182
352	A systematic review of quantitative and qualitative research on the role and effectiveness of written information available to patients about individual medicines. <i>Health Technology Assessment</i> , 2007 , 11, iii, 1-160	4.4	176
351	Smoking cessation in severe mental illness: what works?. <i>Addiction</i> , 2010 , 105, 1176-89	4.6	174
350	Can we predict suicide and non-fatal self-harm with the Beck Hopelessness Scale? A meta-analysis. <i>Psychological Medicine</i> , 2007 , 37, 769-78	6.9	172
349	Is low folate a risk factor for depression? A meta-analysis and exploration of heterogeneity. <i>Journal of Epidemiology and Community Health</i> , 2007 , 61, 631-7	5.1	162
348	Screening and case finding for major depressive disorder using the Patient Health Questionnaire (PHQ-9): a meta-analysis. <i>General Hospital Psychiatry</i> , 2015 , 37, 567-76	5.6	156
347	Managing common mental health disorders in primary care: conceptual models and evidence base. <i>BMJ, The</i> , 2005 , 330, 839-42	5.9	143
346	Rethinking recommendations for screening for depression in primary care. <i>Cmaj</i> , 2012 , 184, 413-8	3.5	142
345	Behavioural activation delivered by the non-specialist: phase II randomised controlled trial. <i>British Journal of Psychiatry</i> , 2011 , 198, 66-72	5.4	141
344	A systematic review, psychometric analysis and qualitative assessment of generic preference-based measures of health in mental health populations and the estimation of mapping functions from widely used specific measures. <i>Health Technology Assessment</i> , 2014 , 18, vii-viii, xiii-xxv, 1-188	4.4	141

343	Methods to identify postnatal depression in primary care: an integrated evidence synthesis and value of information analysis. <i>Health Technology Assessment</i> , 2009 , 13, 1-145, 147-230	4.4	139
342	Should we screen for depression?. <i>BMJ, The</i> , 2006 , 332, 1027-30	5.9	138
341	Benefits and harms of direct to consumer advertising: a systematic review. <i>Quality and Safety in Health Care</i> , 2005 , 14, 246-50		138
340	Clinical effectiveness of collaborative care for depression in UK primary care (CADET): cluster randomised controlled trial. <i>BMJ, The</i> , 2013 , 347, f4913	5.9	136
339	Costs and consequences of enhanced primary care for depression: systematic review of randomised economic evaluations. <i>British Journal of Psychiatry</i> , 2006 , 189, 297-308	5.4	134
338	A meta-review of "lifestyle psychiatry": the role of exercise, smoking, diet and sleep in the prevention and treatment of mental disorders. <i>World Psychiatry</i> , 2020 , 19, 360-380	14.4	132
337	Prevalence of HIV, hepatitis B, and hepatitis C in people with severe mental illness: a systematic review and meta-analysis. <i>Lancet Psychiatry, the</i> , 2016 , 3, 40-48	23.3	127
336	Defining successful treatment outcome in depression using the PHQ-9: a comparison of methods. <i>Journal of Affective Disorders</i> , 2010 , 127, 122-9	6.6	127
335	The CORE-10: A short measure of psychological distress for routine use in the psychological therapies. <i>Counselling and Psychotherapy Research</i> , 2013 , 13, 3-13	1.3	125
334	Diagnosing depression in primary care using self-completed instruments: UK validation of PHQ-9 and CORE-OM. <i>British Journal of General Practice</i> , 2007 , 57, 650-2	1.6	125
333	Informing efficient randomised controlled trials: exploration of challenges in developing progression criteria for internal pilot studies. <i>BMJ Open</i> , 2017 , 7, e013537	3	122
332	Loneliness, social isolation and social relationships: what are we measuring? A novel framework for classifying and comparing tools. <i>BMJ Open</i> , 2016 , 6, e010799	3	122
331	Collaborative care for depression in UK primary care: a randomized controlled trial. <i>Psychological Medicine</i> , 2008 , 38, 279-87	6.9	120
330	Psychotherapy mediated by remote communication technologies: a meta-analytic review. <i>BMC Psychiatry</i> , 2008 , 8, 60	4.2	117
329	Characteristics of effective collaborative care for treatment of depression: a systematic review and meta-regression of 74 randomised controlled trials. <i>PLoS ONE</i> , 2014 , 9, e108114	3.7	115
328	The diagnostic accuracy of brief versions of the Geriatric Depression Scale: a systematic review and meta-analysis. <i>International Journal of Geriatric Psychiatry</i> , 2016 , 31, 837-57	3.9	111
327	Qualitative meta-synthesis of user experience of computerised therapy for depression and anxiety. <i>PLoS ONE</i> , 2014 , 9, e84323	3.7	107
326	Estimating the sample mean and standard deviation from commonly reported quantiles in meta-analysis. <i>Statistical Methods in Medical Research</i> , 2020 , 962280219889080	2.3	104

325	Screening and case finding instruments for depression. <i>The Cochrane Library</i> , 2005 , CD002792	5.2	102
324	Internet-Based Cognitive Behavioral Therapy for Depression: A Systematic Review and Individual Patient Data Network Meta-analysis. <i>JAMA Psychiatry</i> , 2021 , 78, 361-371	14.5	97
323	Fathers' views and experiences of their own mental health during pregnancy and the first postnatal year: a qualitative interview study of men participating in the UK Born and Bred in Yorkshire (BaBY) cohort. <i>BMC Pregnancy and Childbirth</i> , 2017 , 17, 45	3.2	90
322	Delivering stepped care: an analysis of implementation in routine practice. <i>Implementation Science</i> , 2012 , 7, 3	8.4	90
321	Screening for postnatal depression in primary care: cost effectiveness analysis. <i>BMJ, The</i> , 2009 , 339, b5293	5.3	87
320	Accuracy of the PHQ-2 Alone and in Combination With the PHQ-9 for Screening to Detect Major Depression: Systematic Review and Meta-analysis. <i>JAMA - Journal of the American Medical Association</i> , 2020 , 323, 2290-2300	27.4	86
319	Loneliness, social isolation and risk of cardiovascular disease in the English Longitudinal Study of Ageing. <i>European Journal of Preventive Cardiology</i> , 2018 , 25, 1387-1396	3.9	86
318	Identifying depression with the PHQ-2: A diagnostic meta-analysis. <i>Journal of Affective Disorders</i> , 2016 , 203, 382-395	6.6	83
317	Routine use of patient reported outcome measures (PROMs) for improving treatment of common mental health disorders in adults. <i>The Cochrane Library</i> , 2016 , 7, CD011119	5.2	83
316	Newer atypical antipsychotic medication in comparison to clozapine: a systematic review of randomized trials. <i>Schizophrenia Research</i> , 2002 , 56, 1-10	3.6	81
315	Post-it notes to improve questionnaire response rates in RCTs - findings from a randomised sub-study. <i>Trials</i> , 2015 , 16,	2.8	78
314	Prognostic models for predicting relapse or recurrence of depression. <i>The Cochrane Library</i> , 2019 ,	5.2	78
313	Routine administration of Health Related Quality of Life (HRQoL) and needs assessment instruments to improve psychological outcome--a systematic review. <i>Psychological Medicine</i> , 2002 , 32, 1345-56	6.9	77
312	Smoking cessation for people with severe mental illness (SCIMITAR+): a pragmatic randomised controlled trial. <i>Lancet Psychiatry, the</i> , 2019 , 6, 379-390	23.3	75
311	Outcomes research in mental health. Systematic review. <i>British Journal of Psychiatry</i> , 2002 , 181, 8-16	5.4	74
310	Psychological and pharmacological interventions for posttraumatic stress disorder and comorbid mental health problems following complex traumatic events: Systematic review and component network meta-analysis. <i>PLoS Medicine</i> , 2020 , 17, e1003262	11.6	71
309	A systematic review and economic model of the clinical effectiveness and cost-effectiveness of interventions for preventing relapse in people with bipolar disorder. <i>Health Technology Assessment</i> , 2007 , 11, iii-iv, ix-206	4.4	70
308	Depression screening and patient outcomes in pregnancy or postpartum: a systematic review. <i>Journal of Psychosomatic Research</i> , 2014 , 76, 433-46	4.1	69

307	Nurse-delivered collaborative care for depression and long-term physical conditions: a systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2013 , 149, 14-22	6.6	69
306	Improving the detection and management of depression in primary care. <i>Quality and Safety in Health Care</i> , 2003 , 12, 149-55		67
305	The causes, consequences and detection of publication bias in psychiatry. <i>Acta Psychiatrica Scandinavica</i> , 2000 , 102, 241-9	6.5	66
304	Equivalency of the diagnostic accuracy of the PHQ-8 and PHQ-9: a systematic review and individual participant data meta-analysis. <i>Psychological Medicine</i> , 2020 , 50, 1368-1380	6.9	65
303	Smoking cessation in severe mental ill health: what works? an updated systematic review and meta-analysis. <i>BMC Psychiatry</i> , 2017 , 17, 252	4.2	64
302	Modelling publication bias in meta-analysis: a review. <i>Statistical Methods in Medical Research</i> , 2000 , 9, 421-445	2.3	63
301	Effect of Collaborative Care vs Usual Care on Depressive Symptoms in Older Adults With Subthreshold Depression: The CASPER Randomized Clinical Trial. <i>JAMA - Journal of the American Medical Association</i> , 2017 , 317, 728-737	27.4	61
300	Early changes, attrition, and dose-response in low intensity psychological interventions. <i>British Journal of Clinical Psychology</i> , 2014 , 53, 114-30	3.6	61
299	Is self-guided internet-based cognitive behavioural therapy (iCBT) harmful? An individual participant data meta-analysis. <i>Psychological Medicine</i> , 2018 , 48, 2456-2466	6.9	60
298	Bespoke smoking cessation for people with severe mental ill health (SCIMITAR): a pilot randomised controlled trial. <i>Lancet Psychiatry</i> , 2015 , 2, 395-402	23.3	59
297	Patient Health Questionnaire-9 scores do not accurately estimate depression prevalence: individual participant data meta-analysis. <i>Journal of Clinical Epidemiology</i> , 2020 , 122, 115-128.e1	5.7	58
296	Telephone-supported computerised cognitive-behavioural therapy: REEACT-2 large-scale pragmatic randomised controlled trial. <i>British Journal of Psychiatry</i> , 2017 , 210, 362-367	5.4	57
295	How durable is the effect of low intensity CBT for depression and anxiety? Remission and relapse in a longitudinal cohort study. <i>Behaviour Research and Therapy</i> , 2017 , 94, 1-8	5.2	55
294	Association Between Chronic Physical Conditions and the Effectiveness of Collaborative Care for Depression: An Individual Participant Data Meta-analysis. <i>JAMA Psychiatry</i> , 2016 , 73, 978-89	14.5	54
293	Screening for postnatal depression Is it acceptable to women and healthcare professionals? A systematic review and meta-synthesis. <i>Journal of Reproductive and Infant Psychology</i> , 2010 , 28, 328-344	2.9	54
292	Developing a U.K. protocol for collaborative care: a qualitative study. <i>General Hospital Psychiatry</i> , 2006 , 28, 296-305	5.6	54
291	The diagnostic accuracy of the Patient Health Questionnaire-2 (PHQ-2), Patient Health Questionnaire-8 (PHQ-8), and Patient Health Questionnaire-9 (PHQ-9) for detecting major depression: protocol for a systematic review and individual patient data meta-analyses. <i>Systematic Reviews</i> , 2014 , 3, 124	3	53
290	The clinical effectiveness of sertraline in primary care and the role of depression severity and duration (PANDA): a pragmatic, double-blind, placebo-controlled randomised trial. <i>Lancet Psychiatry</i> , 2019 , 6, 903-914	23.3	52

289	Feedback-informed treatment versus usual psychological treatment for depression and anxiety: a multisite, open-label, cluster randomised controlled trial. <i>Lancet Psychiatry</i> , 2018 , 5, 564-572	23.3	52
288	Effectiveness of interventions designed to promote patient involvement to enhance safety: a systematic review. <i>BMJ Quality and Safety</i> , 2010 , 19, e10	5.4	51
287	Brief case finding tools for anxiety disorders: validation of GAD-7 and GAD-2 in addictions treatment. <i>Drug and Alcohol Dependence</i> , 2012 , 125, 37-42	4.9	50
286	The Effects of Meditation, Yoga, and Mindfulness on Depression, Anxiety, and Stress in Tertiary Education Students: A Meta-Analysis. <i>Frontiers in Psychiatry</i> , 2019 , 10, 193	5	47
285	Diagnostic accuracy of the Whooley questions for the identification of depression: a diagnostic meta-analysis. <i>BMJ Open</i> , 2015 , 5, e008913	3	47
284	Is it clinically and cost effective to screen for postnatal depression: a systematic review of controlled clinical trials and economic evidence. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2009 , 116, 1019-27	3.7	46
283	The accessibility and acceptability of self-management support interventions for men with long term conditions: a systematic review and meta-synthesis of qualitative studies. <i>BMC Public Health</i> , 2014 , 14, 1230	4.1	45
282	Instruments to identify post-natal depression: Which methods have been the most extensively validated, in what setting and in which language?. <i>International Journal of Psychiatry in Clinical Practice</i> , 2010 , 14, 72-6	2.4	44
281	What work has to be done to implement collaborative care for depression? Process evaluation of a trial utilizing the Normalization Process Model. <i>Implementation Science</i> , 2010 , 5, 15	8.4	44
280	Patient experience of computerised therapy for depression in primary care. <i>BMJ Open</i> , 2015 , 5, e0085813		43
279	Can we improve the morale of staff working in psychiatric units? A systematic review. <i>Journal of Mental Health</i> , 2006 , 15, 7-17	2.7	43
278	Improving the delivery and organisation of mental health services: beyond the conventional randomised controlled trial. <i>British Journal of Psychiatry</i> , 2002 , 180, 13-8	5.4	43
277	On poverty, politics and psychology: the socioeconomic gradient of mental healthcare utilisation and outcomes. <i>British Journal of Psychiatry</i> , 2016 , 209, 429-430	5.4	42
276	Vegetarianism in young women: another means of weight control?. <i>International Journal of Eating Disorders</i> , 1999 , 26, 87-90	6.3	42
275	The Accuracy of the Patient Health Questionnaire-9 Algorithm for Screening to Detect Major Depression: An Individual Participant Data Meta-Analysis. <i>Psychotherapy and Psychosomatics</i> , 2020 , 89, 25-37	9.4	42
274	How reliable is depression screening in alcohol and drug users? A validation of brief and ultra-brief questionnaires. <i>Journal of Affective Disorders</i> , 2011 , 134, 266-71	6.6	41
273	A randomised controlled trial of computerised cognitive behaviour therapy for the treatment of depression in primary care: the Randomised Evaluation of the Effectiveness and Acceptability of Computerised Therapy (REEACT) trial. <i>Health Technology Assessment</i> , 2015 , 19, viii, xxi-171	4.4	40
272	Prevalence and incidence of postnatal depression: what can systematic reviews tell us?. <i>Archives of Women's Mental Health</i> , 2010 , 13, 295-305	5	38

271	Randomized controlled trials in schizophrenia: a critical perspective on the literature. <i>Acta Psychiatrica Scandinavica</i> , 2002 , 105, 243-51	6.5	38
270	Probability of major depression diagnostic classification using semi-structured versus fully structured diagnostic interviews. <i>British Journal of Psychiatry</i> , 2018 , 212, 377-385	5.4	36
269	Collaborative care for Screen-Positive ElDeRs with major depression (CASPER plus): a multicentred randomised controlled trial of clinical effectiveness and cost-effectiveness. <i>Health Technology Assessment</i> , 2017 , 21, 1-252	4.4	36
268	Improving the efficiency of psychological treatment using outcome feedback technology. <i>Behaviour Research and Therapy</i> , 2017 , 99, 89-97	5.2	35
267	Diagnostic accuracy of the Edinburgh Postnatal Depression Scale (EPDS) for detecting major depression in pregnant and postnatal women: protocol for a systematic review and individual patient data meta-analyses. <i>BMJ Open</i> , 2015 , 5, e009742	3	35
266	The effectiveness of behavioural therapy for the treatment of depression in older adults: a meta-analysis. <i>International Journal of Geriatric Psychiatry</i> , 2011 , 26, 1211-20	3.9	35
265	Cost-effectiveness of collaborative care for depression in UK primary care: economic evaluation of a randomised controlled trial (CADET). <i>PLoS ONE</i> , 2014 , 9, e104225	3.7	33
264	Predicting persistent depressive symptoms in older adults: A machine learning approach to personalised mental healthcare. <i>Journal of Affective Disorders</i> , 2019 , 246, 857-860	6.6	33
263	Selective Cutoff Reporting in Studies of Diagnostic Test Accuracy: A Comparison of Conventional and Individual-Patient-Data Meta-Analyses of the Patient Health Questionnaire-9 Depression Screening Tool. <i>American Journal of Epidemiology</i> , 2017 , 185, 954-964	3.8	32
262	The prevalence of nursing staff stress on adult acute psychiatric in-patient wards. A systematic review. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2006 , 41, 34-43	4.5	32
261	Risperidone versus other atypical antipsychotic medication for schizophrenia. <i>Cochrane Database of Systematic Reviews</i> , 2000 , CD002306		32
260	A review and critical appraisal of measures of therapist-patient interactions in mental health settings. <i>Health Technology Assessment</i> , 2008 , 12, iii, ix-47	4.4	31
259	Psychological distress during pregnancy in a multi-ethnic community: findings from the born in Bradford cohort study. <i>PLoS ONE</i> , 2013 , 8, e60693	3.7	30
258	Cost utility of behavioural activation delivered by the non-specialist. <i>British Journal of Psychiatry</i> , 2011 , 199, 510-1	5.4	30
257	Clinical effectiveness and cost-effectiveness of collaborative care for depression in UK primary care (CADET): a cluster randomised controlled trial. <i>Health Technology Assessment</i> , 2016 , 20, 1-192	4.4	30
256	Evaluation of ethnic disparities in detection of depression and anxiety in primary care during the maternal period: combined analysis of routine and cohort data. <i>British Journal of Psychiatry</i> , 2016 , 208, 453-61	5.4	30
255	Determinants of hospital length of stay for people with serious mental illness in England and implications for payment systems: a regression analysis. <i>BMC Health Services Research</i> , 2015 , 15, 439	2.9	29
254	Putting the 'Q' in depression QALYs: a comparison of utility measurement using EQ-5D and SF-6D health related quality of life measures. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2009 , 44, 569-78	4.5	28

253	Comparison of large versus smaller randomized trials for mental health-related interventions. <i>American Journal of Psychiatry</i> , 2005 , 162, 578-84	11.9	28
252	Usefulness of PHQ-9 in primary care to determine meaningful symptoms of low mood: a qualitative study. <i>British Journal of General Practice</i> , 2016 , 66, e78-84	1.6	27
251	Collaborative Depression Trial (CADET): multi-centre randomised controlled trial of collaborative care for depression--study protocol. <i>BMC Health Services Research</i> , 2009 , 9, 188	2.9	27
250	Clinical effectiveness, cost-effectiveness and acceptability of low-intensity interventions in the management of obsessive-compulsive disorder: the Obsessive-Compulsive Treatment Efficacy randomised controlled Trial (OCTET). <i>Health Technology Assessment</i> , 2017 , 21, 1-132	4.4	27
249	Dismantling, optimising, and personalising internet cognitive behavioural therapy for depression: a systematic review and component network meta-analysis using individual participant data. <i>Lancet Psychiatry</i> , 2021 , 8, 500-511	23.3	27
248	Cost-effectiveness of combining systematic identification and treatment of co-morbid major depression for people with chronic diseases: the example of cancer. <i>Psychological Medicine</i> , 2014 , 44, 1451-60	6.9	26
247	A randomised evaluation of CollABorative care and active surveillance for Screen-Positive ElDeRs with sub-threshold depression (CASPER): study protocol for a randomized controlled trial. <i>Trials</i> , 2011 , 12, 225	2.8	26
246	Varenicline for smoking cessation and reduction in people with severe mental illnesses: systematic review and meta-analysis. <i>Addiction</i> , 2016 , 111, 1554-67	4.6	26
245	The effectiveness of self-management support interventions for men with long-term conditions: a systematic review and meta-analysis. <i>BMJ Open</i> , 2015 , 5, e006620	3	25
244	Diagnostic accuracy of case-finding questions to identify perinatal depression. <i>Cmaj</i> , 2012 , 184, E424-30	3.5	25
243	Comparison of major depression diagnostic classification probability using the SCID, CIDI, and MINI diagnostic interviews among women in pregnancy or postpartum: An individual participant data meta-analysis. <i>International Journal of Methods in Psychiatric Research</i> , 2019 , 28, e1803	4.3	24
242	Improving health and productivity of depressed workers: a pilot randomized controlled trial of telephone cognitive behavioral therapy delivery in workplace settings. <i>General Hospital Psychiatry</i> , 2010 , 32, 337-40	5.6	24
241	Effectiveness and adherence of telephone-administered psychotherapy for depression: A systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2020 , 260, 514-526	6.6	24
240	The Effectiveness of Pharmacological and Non-Pharmacological Interventions for Improving Glycaemic Control in Adults with Severe Mental Illness: A Systematic Review and Meta-Analysis. <i>PLoS ONE</i> , 2017 , 12, e0168549	3.7	23
239	Outcome measures and needs assessment tools for schizophrenia and related disorders. <i>The Cochrane Library</i> , 2003 , CD003081	5.2	23
238	Feasibility randomized controlled trial of cognitive and behavioral interventions for depression symptoms in patients accessing drug and alcohol treatment. <i>Journal of Substance Abuse Treatment</i> , 2015 , 55, 6-14	4.2	22
237	Assessing the quality of diagnostic studies using psychometric instruments: applying QUADAS. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2009 , 44, 300-7	4.5	22
236	Cluster randomized trials produced similar results to individually randomized trials in a meta-analysis of enhanced care for depression. <i>Journal of Clinical Epidemiology</i> , 2008 , 61, 160-168	5.7	22

235	Smoking Cessation Intervention for severe Mental Ill Health Trial (SCIMITAR): a pilot randomised control trial of the clinical effectiveness and cost-effectiveness of a bespoke smoking cessation service. <i>Health Technology Assessment</i> , 2015 , 19, 1-148, v-vi	4.4	22
234	Clinical effectiveness and cost-effectiveness of parenting interventions for children with severe attachment problems: a systematic review and meta-analysis. <i>Health Technology Assessment</i> , 2015 , 19, vii-xxviii, 1-347	4.4	22
233	Probability of major depression diagnostic classification based on the SCID, CIDI and MINI diagnostic interviews controlling for Hospital Anxiety and Depression Scale - Depression subscale scores: An individual participant data meta-analysis of 73 primary studies. <i>Journal of Psychosomatic Research</i> , 2020 , 129, 109892	4.1	22
232	Detecting and predicting self-harm behaviour in prisoners: a prospective psychometric analysis of three instruments. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2009 , 44, 853-61	4.5	20
231	The second Randomised Evaluation of the Effectiveness, cost-effectiveness and Acceptability of Computerised Therapy (REEACT-2) trial: does the provision of telephone support enhance the effectiveness of computer-delivered cognitive behaviour therapy? A randomised controlled trial. <i>Health Technology Assessment</i> , 2016 , 20, 1-64	4.4	20
230	Diagnostic accuracy of the Depression subscale of the Hospital Anxiety and Depression Scale (HADS-D) for detecting major depression: protocol for a systematic review and individual patient data meta-analyses. <i>BMJ Open</i> , 2016 , 6, e011913	3	19
229	Cost and outcome of behavioural activation versus cognitive behaviour therapy for depression (COBRA): study protocol for a randomised controlled trial. <i>Trials</i> , 2014 , 15, 29	2.8	18
228	Sudden gains in behavioural activation for depression. <i>Behaviour Research and Therapy</i> , 2014 , 60, 34-8	5.2	18
227	The psychometric properties of the subscales of the GHQ-28 in a multi-ethnic maternal sample: results from the Born in Bradford cohort. <i>BMC Psychiatry</i> , 2013 , 13, 55	4.2	18
226	Translating research into practice in Leeds and Bradford (TRiPLaB): a protocol for a programme of research. <i>Implementation Science</i> , 2010 , 5, 37	8.4	18
225	Evidence-based medicine: An improved format for journal clubs. <i>Psychiatric Bulletin</i> , 1996 , 20, 673-675		18
224	Variation and ethnic inequalities in treatment of common mental disorders before, during and after pregnancy: combined analysis of routine and research data in the Born in Bradford cohort. <i>BMC Psychiatry</i> , 2016 , 16, 99	4.2	17
223	Revealing hidden depression in older people: a qualitative study within a randomised controlled trial. <i>BMC Family Practice</i> , 2015 , 16, 142	2.6	17
222	Relation between hostility and coronary heart disease. Evidence does not support link. <i>BMJ: British Medical Journal</i> , 1999 , 319, 917-8		17
221	Impact of long-term medical conditions on the outcomes of psychological therapy for depression and anxiety. <i>British Journal of Psychiatry</i> , 2017 , 210, 47-53	5.4	16
220	Benchmarking routine psychological services: a discussion of challenges and methods. <i>Behavioural and Cognitive Psychotherapy</i> , 2014 , 42, 16-30	2.1	16
219	Combining mirtazapine with SSRIs or SNRIs for treatment-resistant depression: the MIR RCT. <i>Health Technology Assessment</i> , 2018 , 22, 1-136	4.4	16
218	Maintenance or Discontinuation of Antidepressants in Primary Care. <i>New England Journal of Medicine</i> , 2021 , 385, 1257-1267	59.2	16

217	Heterogeneity in patient-reported outcomes following low-intensity mental health interventions: a multilevel analysis. <i>PLoS ONE</i> , 2014 , 9, e99658	3.7	15
216	Adapting manualized Behavioural Activation treatment for older adults with depression. <i>The Cognitive Behaviour Therapist</i> , 2014 , 7,	2.9	15
215	Validity of two case finding questions to detect postnatal depression: a review of diagnostic test accuracy. <i>Journal of Affective Disorders</i> , 2011 , 133, 388-97	6.6	15
214	Impact of family practice continuity of care on unplanned hospital use for people with serious mental illness. <i>Health Services Research</i> , 2019 , 54, 1316-1325	3.4	15
213	How much change is enough? Evidence from a longitudinal study on depression in UK primary care. <i>Psychological Medicine</i> , 2020 , 1-8	6.9	14
212	Low-intensity cognitive-behaviour therapy interventions for obsessive-compulsive disorder compared to waiting list for therapist-led cognitive-behaviour therapy: 3-arm randomised controlled trial of clinical effectiveness. <i>PLoS Medicine</i> , 2017 , 14, e1002337	11.6	14
211	A randomised controlled trial assessing the severity and duration of depressive symptoms associated with a clinically significant response to sertraline versus placebo, in people presenting to primary care with depression (PANDA trial): study protocol for a randomised controlled trial. <i>Trials</i> , 2017 , 18, 496	2.8	14
210	Identifying primary care quality indicators for people with serious mental illness: a systematic review. <i>British Journal of General Practice</i> , 2017 , 67, e519-e530	1.6	14
209	Does the quality and outcomes framework reduce psychiatric admissions in people with serious mental illness? A regression analysis. <i>BMJ Open</i> , 2015 , 5, e007342	3	14
208	Promoting HIV and sexual safety behaviour in people with severe mental illness: a systematic review of behavioural interventions. <i>International Journal of Mental Health Nursing</i> , 2014 , 23, 344-54	3.8	14
207	Can occupational therapist-led home environmental assessment prevent falls in older people? A modified cohort randomised controlled trial protocol. <i>BMJ Open</i> , 2018 , 8, e022488	3	14
206	Nature-based outdoor activities for mental and physical health: Systematic review and meta-analysis. <i>SSM - Population Health</i> , 2021 , 16, 100934	3.8	14
205	Computerised cognitive behaviour therapy for depression in adolescents: study protocol for a feasibility randomised controlled trial. <i>BMJ Open</i> , 2014 , 4, e006488	3	13
204	Differences in use of abbreviated mental test score by geriatricians and psychiatrists. <i>BMJ: British Medical Journal</i> , 1996 , 313, 465		13
203	Cost and Outcome of Behavioural Activation (COBRA): a randomised controlled trial of behavioural activation versus cognitive-behavioural therapy for depression. <i>Health Technology Assessment</i> , 2017 , 21, 1-366	4.4	13
202	Making the journey with me: a qualitative study of experiences of a bespoke mental health smoking cessation intervention for service users with serious mental illness. <i>BMC Psychiatry</i> , 2016 , 16, 193	4.2	13
201	Diagnostic accuracy of the Geriatric Depression Scale-30, Geriatric Depression Scale-15, Geriatric Depression Scale-5 and Geriatric Depression Scale-4 for detecting major depression: protocol for a systematic review and individual participant data meta-analysis. <i>BMJ Open</i> , 2018 , 8, e026598	3	13
200	Maternal psychological distress in primary care and association with child behavioural outcomes at age three. <i>European Child and Adolescent Psychiatry</i> , 2016 , 25, 601-13	5.5	12

199	Association Between Antipsychotic Polypharmacy and Outcomes for People With Serious Mental Illness in England. <i>Psychiatric Services</i> , 2019 , 70, 650-656	3.3	12
198	Smoking Cessation Intervention for Severe Mental Ill Health Trial (SCIMITAR+): study protocol for a randomised controlled trial. <i>Trials</i> , 2017 , 18, 44	2.8	12
197	CASPER plus (Collaborative care in Screen-Positive Elderly with major depressive disorder): study protocol for a randomised controlled trial. <i>Trials</i> , 2014 , 15, 451	2.8	12
196	Screening and case finding for depression in offender populations: a systematic review of diagnostic properties. <i>Journal of Affective Disorders</i> , 2011 , 128, 72-82	6.6	12
195	Efficacy of beta-blocker supplementation for schizophrenia: a systematic review of randomized trials. <i>Schizophrenia Research</i> , 2000 , 41, 341-7	3.6	12
194	The role of electronic journals in reducing publication bias. <i>Informatics for Health and Social Care</i> , 1999 , 24, 223-9		12
193	Life adversity in depressed and non-depressed older adults: A cross-sectional comparison of the brief LTE-Q questionnaire and life events and difficulties interview as part of the CASPER study. <i>Journal of Affective Disorders</i> , 2016 , 193, 31-8	6.6	12
192	Factors associated with regular physical activity participation among people with severe mental ill health. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2019 , 54, 887-895	4.5	12
191	Shortening self-report mental health symptom measures through optimal test assembly methods: Development and validation of the Patient Health Questionnaire-Depression-4. <i>Depression and Anxiety</i> , 2019 , 36, 82-92	8.4	12
190	Cost-effectiveness of computerized cognitive-behavioural therapy for the treatment of depression in primary care: findings from the Randomised Evaluation of the Effectiveness and Acceptability of Computerised Therapy (REEACT) trial. <i>Psychological Medicine</i> , 2017 , 47, 1825-1835	6.9	11
189	Are there researcher allegiance effects in diagnostic validation studies of the PHQ-9? A systematic review and meta-analysis. <i>BMJ Open</i> , 2017 , 7, e015247	3	11
188	Lost information? The fate of papers presented at the 40th society for Social Medicine Conference. <i>Journal of Epidemiology and Community Health</i> , 1999 , 53, 442-3	5.1	11
187	What factors indicate prognosis for adults with depression in primary care? A protocol for meta-analyses of individual patient data using the Dep-GP database. <i>Wellcome Open Research</i> , 2019 , 4, 69	4.8	11
186	The associations between loneliness, social exclusion and pain in the general population: A N=502,528 cross-sectional UK Biobank study. <i>Journal of Psychiatric Research</i> , 2020 , 130, 68-74	5.2	11
185	Identification of depression in women during pregnancy and the early postnatal period using the Whooley questions and the Edinburgh Postnatal Depression Scale: protocol for the Born and Bred in Yorkshire: PeriNatal Depression Diagnostic Accuracy (BaBY PaNDA) study. <i>BMJ Open</i> , 2016 , 6, e011223	3	11
184	Depression prevalence based on the Edinburgh Postnatal Depression Scale compared to Structured Clinical Interview for DSM Disorders classification: Systematic review and individual participant data meta-analysis. <i>International Journal of Methods in Psychiatric Research</i> , 2021 , 30, e1860	4.3	11
183	Role of age, gender and marital status in prognosis for adults with depression: An individual patient data meta-analysis. <i>Epidemiology and Psychiatric Sciences</i> , 2021 , 30, e42	5.1	11
182	Exploring why people with SMI smoke and why they may want to quit: baseline data from the SCIMITAR RCT. <i>Journal of Psychiatric and Mental Health Nursing</i> , 2016 , 23, 282-9	2.4	10

181	Randomized trials with concurrent economic evaluations reported unrepresentatively large clinical effect sizes. <i>Journal of Clinical Epidemiology</i> , 2007 , 60, 781-6	5.7	10
180	Influence of methodology on outcomes of randomised clozapine trials. <i>Pharmacopsychiatry</i> , 2000 , 33, 54-9	2	10
179	Pharmacy-based management for depression in adults. <i>The Cochrane Library</i> , 2019 , 12, CD013299	5.2	10
178	GPs' attitudes towards digital technologies for depression: an online survey in primary care. <i>British Journal of General Practice</i> , 2019 , 69, e164-e170	1.6	10
177	Impact of telephone delivered case-management on the effectiveness of collaborative care for depression and anti-depressant use: A systematic review and meta-regression. <i>PLoS ONE</i> , 2019 , 14, e0217948	3.7	9
176	Mirtazapine added to selective serotonin reuptake inhibitors for treatment-resistant depression in primary care (MIR trial): study protocol for a randomised controlled trial. <i>Trials</i> , 2016 , 17, 66	2.8	9
175	Successful recruitment to trials: findings from the SCIMITAR+ Trial. <i>Trials</i> , 2018 , 19, 53	2.8	9
174	Obsessive Compulsive Treatment Efficacy Trial (OCTET) comparing the clinical and cost effectiveness of self-managed therapies: study protocol for a randomised controlled trial. <i>Trials</i> , 2014 , 15, 278	2.8	9
173	Review: disease management programmes improve detection and care of people with depression. <i>Evidence-Based Mental Health</i> , 2004 , 7, 80	11.1	9
172	Assessing the risk of self-harm in an adult offender population: an incidence cohort study. <i>Health Technology Assessment</i> , 2014 , 18, 1-151, vii-viii	4.4	9
171	What factors indicate prognosis for adults with depression in primary care? A protocol for meta-analyses of individual patient data using the Dep-GP database. <i>Wellcome Open Research</i> , 2019 , 4, 69	4.8	9
170	Probability of Major Depression Classification Based on the SCID, CIDI, and MINI Diagnostic Interviews: A Synthesis of Three Individual Participant Data Meta-Analyses. <i>Psychotherapy and Psychosomatics</i> , 2021 , 90, 28-40	9.4	9
169	The HARMONIC trial: study protocol for a randomised controlled feasibility trial of Shaping Healthy Minds-a modular transdiagnostic intervention for mood, stressor-related and anxiety disorders in adults. <i>BMJ Open</i> , 2018 , 8, e024546	3	9
168	The Current View of Evidence and Evidence-Based Practice1-20		9
167	Should methodological filters for diagnostic test accuracy studies be used in systematic reviews of psychometric instruments? A case study involving screening for postnatal depression. <i>Systematic Reviews</i> , 2012 , 1, 9	3	8
166	Frailty and depression predict instrumental activities of daily living in older adults: A population-based longitudinal study using the CARE75+ cohort. <i>PLoS ONE</i> , 2020 , 15, e0243972	3.7	8
165	Identifying perinatal depression with case-finding instruments: a mixed-methods study (BaBY PaNDA [Born and Bred in Yorkshire PeriNatal Depression Diagnostic Accuracy]). <i>Health Services and Delivery Research</i> , 2018 , 6, 1-210	1.5	8
164	The role of relapse prevention for depression in collaborative care: A systematic review. <i>Journal of Affective Disorders</i> , 2020 , 265, 618-644	6.6	8

163	Smoking cessation in severe mental illness: combined long-term quit rates from the UK SCIMITAR trials programme. <i>British Journal of Psychiatry</i> , 2021 , 218, 95-97	5.4	8
162	A randomised controlled trial assessing the use of citalopram, sertraline, fluoxetine and mirtazapine in preventing relapse in primary care patients who are taking long-term maintenance antidepressants (ANTLER: ANTidepressants to prevent reLapse in dEpReSSion): study protocol for a randomised controlled trial. <i>Trials</i> , 2019 , 20, 319	2.8	7
161	Diagnostic accuracy of the Whooley depression tool in older adults in UK primary care. <i>Journal of Affective Disorders</i> , 2015 , 182, 39-43	6.6	7
160	Acceptability of mental health screening in routine addictions treatment. <i>General Hospital Psychiatry</i> , 2012 , 34, 415-22	5.6	7
159	Collaborative care and active surveillance for Screen-Positive ElDeRs with subthreshold depression (CASPER): a multicentred randomised controlled trial of clinical effectiveness and cost-effectiveness. <i>Health Technology Assessment</i> , 2017 , 21, 1-196	4.4	7
158	The contribution of depressive 'disorder characteristics' to determinations of prognosis for adults with depression: an individual patient data meta-analysis. <i>Psychological Medicine</i> , 2021 , 51, 1068-1081	6.9	7
157	Predicting prognosis for adults with depression using individual symptom data: a comparison of modelling approaches. <i>Psychological Medicine</i> , 2021 , 1-11	6.9	7
156	How should we implement collaborative care for older people with depression? A qualitative study using normalisation process theory within the CASPER plus trial. <i>BMC Family Practice</i> , 2018 , 19, 116	2.6	7
155	REDUCE (Reviewing long-term antidepressant use by careful monitoring in everyday practice) internet and telephone support to people coming off long-term antidepressants: protocol for a randomised controlled trial. <i>Trials</i> , 2020 , 21, 419	2.8	6
154	Relapse and Recurrence of Common Mental Health Problems after Low Intensity Cognitive Behavioural Therapy: The WYLOW Longitudinal Cohort Study. <i>Psychotherapy and Psychosomatics</i> , 2018 , 87, 116-117	9.4	6
153	Depression, anxiety and comorbid substance use: association patterns in outpatient addictions treatment. <i>Mental Health and Substance Use: Dual Diagnosis</i> , 2013 , 6, 59-75		6
152	Collaborative care for depression and anxiety problems 2007 ,		6
151	Prescribing antidepressants in general practice 1997 , 314, 826-826		6
150	Nobody ever expects the Spanish Inquisition[Python, 1991). <i>Psychiatric Bulletin</i> , 2001 , 25, 291-292		6
149	A bespoke smoking cessation service compared with treatment as usual for people with severe mental ill health: the SCIMITAR+ RCT. <i>Health Technology Assessment</i> , 2019 , 23, 1-116	4.4	6
148	Health of mothers of children with a life-limiting condition: a comparative cohort study. <i>Archives of Disease in Childhood</i> , 2021 , 106, 987-993	2.2	6
147	Shared goals for mental health research: what, why and when for the 2020s. <i>Journal of Mental Health</i> , 2021 , 1-9	2.7	6
146	Living with diabetes alongside a severe mental illness: A qualitative exploration with people with severe mental illness, family members and healthcare staff. <i>Diabetic Medicine</i> , 2021 , 38, e14562	3.5	6

145	Mitigating the impacts of COVID-19: where are the mental health trials?. <i>Lancet Psychiatry,the</i> , 2021 , 8, 647-650	23.3	6
144	Use of the Internet and Digital Devices Among People With Severe Mental Ill Health During the COVID-19 Pandemic Restrictions. <i>Frontiers in Psychiatry</i> , 2021 , 12, 732735	5	6
143	A systematic review of mental health measurement scales for evaluating the effects of mental health prevention interventions. <i>European Journal of Public Health</i> , 2020 , 30, 539-545	2.1	5
142	The ethics of Trials within Cohorts(TwCs): 2nd international symposium. <i>Trials</i> , 2017 , 18,	2.8	5
141	Decision curve analysis for assessing the usefulness of tests for making decisions to treat: an application to tests for prodromal psychosis. <i>Psychological Assessment</i> , 2013 , 25, 730-7	5.3	5
140	User-defined outcomes in mental health: A qualitative study and consensus development exercise. <i>Journal of Mental Health</i> , 2009 , 18, 415-423	2.7	5
139	IMPACT collaborative care programme reduces suicide ideation in depressed older adults. <i>Evidence-Based Mental Health</i> , 2007 , 10, 51	11.1	5
138	Publication bias and meta-analysis. <i>British Journal of Psychiatry</i> , 1995 , 167, 266	5.4	5
137	Loneliness among people with severe mental illness during the COVID-19 pandemic: Results from a linked UK population cohort study.. <i>PLoS ONE</i> , 2022 , 17, e0262363	3.7	5
136	A systematic review and metaethnography to identify how effective, cost-effective, accessible and acceptable self-management support interventions are for men with long-term conditions (SELF-MAN). <i>Health Services and Delivery Research</i> , 2015 , 3, 1-302	1.5	5
135	Screening for psychological and mental health difficulties in young people who offend: a systematic review and decision model. <i>Health Technology Assessment</i> , 2015 , 19, 1-128	4.4	5
134	Rapid systematic review of systematic reviews: what befriending, social support and low intensity psychosocial interventions, delivered remotely, may reduce social isolation and loneliness among older adults and how?. <i>F1000Research</i> , 9 , 1368	3.6	5
133	Barriers and facilitators to implementing perinatal mental health care in health and social care settings: a systematic review. <i>Lancet Psychiatry,the</i> , 2021 , 8, 521-534	23.3	5
132	Cost-effectiveness of feedback-informed psychological treatment: Evidence from the IAPT-FIT trial. <i>Behaviour Research and Therapy</i> , 2021 , 142, 103873	5.2	5
131	Cost-effectiveness of a specialist smoking cessation package compared with standard smoking cessation services for people with severe mental illness in England: a trial-based economic evaluation from the SCIMITAR+ study. <i>Addiction</i> , 2020 , 115, 2113-2122	4.6	5
130	Is social support pre-treatment associated with prognosis for adults with depression in primary care?. <i>Acta Psychiatrica Scandinavica</i> , 2021 , 143, 392-405	6.5	5
129	Individual participant data (IPD) meta-analysis of psychological relapse prevention interventions versus control for patients in remission from depression: a protocol. <i>BMJ Open</i> , 2020 , 10, e034158	3	4
128	Psychometric properties of the Treatment Outcomes Profile (TOP) psychological health scale. <i>Mental Health and Substance Use: Dual Diagnosis</i> , 2013 , 6, 140-149		4

127	An randomized controlled trial of Post-it [®] notes did not increase postal response rates in older depressed participants. <i>Journal of Evaluation in Clinical Practice</i> , 2017 , 23, 102-107	2.5	4
126	The acceptability of case-finding questions to identify perinatal depression. <i>British Journal of Midwifery</i> , 2015 , 23, 630-638	0.3	4
125	Mental health promotion in high risk groups. <i>Quality and Safety in Health Care</i> , 1997 , 6, 219-25		4
124	Drug treatments for schizophrenia. <i>Quality in Health Care: QHC</i> , 2000 , 9, 73-9		4
123	Investigating SOcial Competence and Isolation in children with Autism taking part in LEGO-based therapy clubs In School Environments (I-SOCIALISE): study protocol. <i>BMJ Open</i> , 2019 , 9, e030471	3	4
122	Do higher primary care practice performance scores predict lower rates of emergency admissions for persons with serious mental illness? An analysis of secondary panel data. <i>Health Services and Delivery Research</i> , 2015 , 3, 1-68	1.5	4
121	The digital divide: amplifying health inequalities for people with severe mental illness in the time of COVID-19. <i>British Journal of Psychiatry</i> , 2021 , 219, 529-531	5.4	4
120	Prognostic models for predicting relapse or recurrence of major depressive disorder in adults. <i>The Cochrane Library</i> , 2021 , 5, CD013491	5.2	4
119	Disentangling psychopathology, substance use and dependence: a factor analysis. <i>BMC Psychiatry</i> , 2016 , 16, 281	4.2	4
118	How can we estimate QALYs based on PHQ-9 scores? Equipercentile linking analysis of PHQ-9 and EQ-5D. <i>Evidence-Based Mental Health</i> , 2021 ,	11.1	4
117	Do care plans and annual reviews of physical health influence unplanned hospital utilisation for people with serious mental illness? Analysis of linked longitudinal primary and secondary healthcare records in England. <i>BMJ Open</i> , 2018 , 8, e023135	3	4
116	Should Google offer an online screening test for depression?. <i>BMJ, The</i> , 2017 , 358, j4144	5.9	3
115	Can support workers from AgeUK deliver an intervention to support older people with anxiety and depression? A qualitative evaluation. <i>BMC Family Practice</i> , 2019 , 20, 16	2.6	3
114	Community Pharmacies Mood Intervention Study (CHEMIST): feasibility and external pilot randomised controlled trial protocol. <i>Pilot and Feasibility Studies</i> , 2019 , 5, 71	1.9	3
113	Everything you wanted to know about e-cigarettes and vaping but were afraid to ask: a guide for mental health clinicians. <i>BJ Psych Advances</i> , 2019 , 25, 279-286	0.8	3
112	A feasibility study for NON-Traditional providers to support the management of Elderly People with Anxiety and Depression: The NOTEPAD study Protocol. <i>Trials</i> , 2018 , 19, 172	2.8	3
111	German EstSmoke: estimating adult smoking-related costs and consequences of smoking cessation for Germany. <i>Addiction</i> , 2018 , 113, 125-136	4.6	3
110	Routine use of patient reported outcome measures (PROMs) for improving treatment of common mental health disorders in adults 2014 ,		3

109	Depression in older adults: collaborative care model seems effective. <i>Evidence-Based Mental Health</i> , 2008 , 11, 44	11.1	3
108	Planning and Conducting Systematic Reviews150-179		3
107	Misleading meta-analysis. Subject to many potential biases. <i>BMJ: British Medical Journal</i> , 1995 , 311, 1303-4		3
106	Optimised patient information materials and recruitment to a study of behavioural activation in older adults: an embedded study within a trial. <i>F1000Research</i> , 2020 , 9, 417	3.6	3
105	Interventions for adults with a history of complex traumatic events: the INCiTE mixed-methods systematic review. <i>Health Technology Assessment</i> , 2020 , 24, 1-312	4.4	3
104	Exploring Severe Mental Illness and Diabetes: Protocol for a Longitudinal, Observational, and Qualitative Mixed Methods Study. <i>JMIR Research Protocols</i> , 2019 , 8, 13407	2	3
103	Antidepressant treatment with sertraline for adults with depressive symptoms in primary care: the PANDA research programme including RCT. <i>Programme Grants for Applied Research</i> , 2019 , 7, 1-108	1.5	3
102	Behavioural activation to prevent depression and loneliness among socially isolated older people with long-term conditions: The BASIL COVID-19 pilot randomised controlled trial. <i>PLoS Medicine</i> , 2021 , 18, e1003779	11.6	3
101	Overestimation of Postpartum Depression Prevalence Based on a 5-item Version of the EPDS: Systematic Review and Individual Participant Data Meta-analysis. <i>Canadian Journal of Psychiatry</i> , 2020 , 65, 835-844	4.8	3
100	Antidepressants for major depression disorder in older people: a network meta-analysis. <i>The Cochrane Library</i> , 2019 ,	5.2	3
99	Preventing depression among older people living in rural areas: A randomised controlled trial of behavioural activation in collaborative care. <i>International Journal of Geriatric Psychiatry</i> , 2021 , 36, 530-539	3.9	3
98	Data-driven methods distort optimal cutoffs and accuracy estimates of depression screening tools: a simulation study using individual participant data. <i>Journal of Clinical Epidemiology</i> , 2021 , 137, 137-147	5.7	3
97	Pharmacy-based management for depression in adults. <i>The Cochrane Library</i> ,	5.2	3
96	Can we improve recruitment to trials and informed consent by improving participant information sheets? - A nested RCT. <i>Trials</i> , 2011 , 12,	2.8	2
95	Recruiting ahead of target: What worked in the REEACT trial?. <i>Trials</i> , 2011 , 12,	2.8	2
94	Re-reflecting the article by Thombs and colleagues: Figure 1:. <i>Cmaj</i> , 2012 , 184, 438.2-439	3.5	2
93	Collaborative care for depression in UK primary care: a randomized controlled trial [Corrigendum. <i>Psychological Medicine</i> , 2009 , 39, 701-701	6.9	2
92	Rational decision-making in psychiatry: evidence-based psychiatry is just the start. <i>British Journal of Psychiatry</i> , 1997 , 171, 226-7	5.4	2

91	A Patient Stratification Approach to Identifying the Likelihood of Continued Chronic Depression and Relapse Following Treatment for Depression.. <i>Journal of Personalized Medicine</i> , 2021 , 11,	3.6	2
90	NICE work?: Potential and problems for psychiatry in the new National Health Service. <i>Psychiatric Bulletin</i> , 1999 , 23, 513-516		2
89	SCIMITAR+ Trial: A randomised study within a trial (SWAT) of a contingent financial reward to improve trial follow-up. <i>F1000Research</i> , 2019 , 8, 1937	3.6	2
88	Rapid systematic review of systematic reviews: what befriending, social support and low intensity psychosocial interventions, delivered remotely, are effective in reducing social isolation and loneliness among older adults? How do they work?. <i>F1000Research</i> , 2019 , 9, 1368	3.6	2
87	SCIMITAR+ Trial: A randomised study within a trial (SWAT) of a contingent financial reward to improve trial follow-up. <i>F1000Research</i> , 2019 , 8, 1937	3.6	2
86	What does the rise and rise of electronic nicotine delivery systems mean for mental health services?. <i>Lancet Psychiatry</i> , 2020 , 7, 11-13	23.3	2
85	Conversion and neuro-inflammation disorder observational study (CANDO). Protocol of a feasibility study. <i>European Journal of Psychiatry</i> , 2020 , 34, 164-172	1	2
84	Supporting smoking cessation and preventing relapse following a stay in a smoke-free setting: a meta-analysis and investigation of effective behaviour change techniques. <i>Addiction</i> , 2021 , 116, 2978-2994	4.6	2
83	Investigating smoking and nicotine dependence among people with severe mental illness during the COVID-19 pandemic: analysis of linked data from a UK Closing the Gap cohort. <i>BJPsych Open</i> , 2021 , 7, e86	5	2
82	The importance of transdiagnostic symptom level assessment to understanding prognosis for depressed adults: analysis of data from six randomised control trials. <i>BMC Medicine</i> , 2021 , 19, 109	11.4	2
81	Can We Prevent Depression in At-Risk Older Adults Using Self-Help? The UK SHARD Trial of Behavioral Activation. <i>American Journal of Geriatric Psychiatry</i> , 2021 ,	6.5	2
80	Home hazard assessment and environmental modification to prevent falls in older people: the OTIS trial. <i>F1000Research</i> , 2020 , 10, 500	3.6	2
79	Behavioural activation in nursing homes to treat depression (BAN-Dep): study protocol for a pragmatic randomised controlled trial. <i>BMJ Open</i> , 2019 , 9, e032421	3	2
78	Impact of severe mental illness on healthcare use and health outcomes for people with type 2 diabetes: a longitudinal observational study in England. <i>British Journal of General Practice</i> , 2021 , 71, e565-e573	1.6	2
77	Home environmental assessments and modification delivered by occupational therapists to reduce falls in people aged 65 years and over: the OTIS RCT. <i>Health Technology Assessment</i> , 2021 , 25, 1-118	4.4	2
76	Socioeconomic Indicators of Treatment Prognosis for Adults With Depression: A Systematic Review and Individual Patient Data Meta-analysis.. <i>JAMA Psychiatry</i> , 2022 ,	14.5	2
75	Cognitive or behavioural interventions (or both) to prevent or mitigate loneliness in adolescents, adults, and older adults. <i>The Cochrane Library</i> , 2020 ,	5.2	1
74	E-cigarette use and associated factors among smokers with severe mental illness. <i>Addictive Behaviors</i> , 2020 , 108, 106456	4.2	1

73	Commentary on: Suicide Screening Tools for Use with Incarcerated Offenders. <i>Archives of Suicide Research</i> , 2019 , 23, 523-525	2.3	1
72	The European pharmaceutical industry: coming to a screen near you?. <i>Journal of the Royal Society of Medicine</i> , 2008 , 101, 485-8	2.3	1
71	Vitamin K and the Prevention of FracturesReply. <i>Archives of Internal Medicine</i> , 2007 , 167, 94		1
70	4 factors were predictive of extended duration in hospital in acute psychiatric admission. <i>Evidence-Based Mental Health</i> , 2002 , 5, 64	11.1	1
69	'Number needed to treat'. <i>British Journal of Psychiatry</i> , 1996 , 169, 113	5.4	1
68	Deliberate self harm. <i>Emergency Medicine Journal</i> , 1996 , 13, 432	1.5	1
67	A Year Into the Pandemic: The Diversity of Experience Amongst People With Severe Mental Ill Health.. <i>Frontiers in Psychiatry</i> , 2021 , 12, 794585	5	1
66	Cost-utility analysis of LEGO based therapy for school children and young people with autism spectrum disorder: results from a randomised controlled trial.. <i>BMJ Open</i> , 2022 , 12, e056347	3	1
65	The feasibility and acceptability of a brief psychological intervention for adults with long-term health conditions and subthreshold depression delivered via community pharmacies: a mixed methods evaluation-the Community Pharmacies Mood Intervention Study (CHEMIST).. <i>Pilot and Feasibility Studies</i> , 2022 , 8, 27	1.9	1
64	Cost-Utility Analysis of Discontinuing Antidepressants in England Primary Care Patients Compared with Long-Term Maintenance: The ANTLER Study. <i>Applied Health Economics and Health Policy</i> , 2021 , 1	3.4	1
63	The association between primary care quality and health-care use, costs and outcomes for people with serious mental illness: a retrospective observational study. <i>Health Services and Delivery Research</i> , 2020 , 8, 1-126	1.5	1
62	Collaborative case management to aid return to work after long-term sickness absence: a pilot randomised controlled trial. <i>Public Health Research</i> , 2018 , 6, 1-76	1.7	1
61	Health risk behaviours among people with severe mental ill health during the COVID-19 pandemic: Analysis of linked cohort data. <i>PLoS ONE</i> , 2021 , 16, e0258349	3.7	1
60	A systematic review of behavioural smoking cessation interventions for people with severe mental ill health-what works?. <i>Addiction</i> , 2021 ,	4.6	1
59	Non-traditional support workers delivering a brief psychosocial intervention for older people with anxiety and depression: the NOTEPAD feasibility study. <i>Health Services and Delivery Research</i> , 2019 , 7, 1-120	1.5	1
58	What factors indicate prognosis for adults with depression in primary care? A protocol for meta-analyses of individual patient data using the Dep-GP database. <i>Wellcome Open Research</i> , 2021 , 4, 69	4.8	1
57	Depression in primary care1-16		1
56	United Kingdom perspective127-136		1

55	The Management of Depression and Anxiety in Older People: Evidence-Based Psychological Interventions, Stepped Care and Collaborative Care 2016 , 79-92		1
54	Identifying determinants of diabetes risk and outcomes for people with severe mental illness: a mixed-methods study. <i>Health Services and Delivery Research</i> , 2021 , 9, 1-194	1.5	1
53	Use of the Internet and digital devices among people with severe mental ill health during the COVID-19 pandemic restrictions		1
52	The development and validation of a prognostic model to PREDICT Relapse of depression in adult patients in primary care: protocol for the PREDICTR study. <i>Diagnostic and Prognostic Research</i> , 2021 , 5, 12	5.5	1
51	Still 'being bothered about Billy': managing the physical health of people with severe mental illness. <i>British Journal of General Practice</i> , 2021 , 71, 373-376	1.6	1
50	Loneliness among people with severe mental ill health during the COVID-19 pandemic: results from a linked UK population cohort study		1
49	Healthcare resource use and costs for people with type 2 diabetes mellitus with and without severe mental illness in England: longitudinal matched-cohort study using the Clinical Practice Research Datalink.. <i>British Journal of Psychiatry</i> , 2021 , 1-8	5.4	1
48	Shortening the Edinburgh postnatal depression scale using optimal test assembly methods: Development of the EPDS-Dep-5. <i>Acta Psychiatrica Scandinavica</i> , 2021 , 143, 348-362	6.5	1
47	Depressive Disorders in Primary Care: A Review 271-318		1
46	Factors that influence participation in physical activity for anxiety or depression: a synthesis of qualitative evidence. <i>The Cochrane Library</i> , 2020 ,	5.2	0
45	Predicting relapse or recurrence of depression: systematic review of prognostic models.. <i>British Journal of Psychiatry</i> , 2022 , 1-11	5.4	0
44	Antidepressant medication to prevent depression relapse in primary care: the ANTLE RCT. <i>Health Technology Assessment</i> , 2021 , 25, 1-62	4.4	0
43	Life events and treatment prognosis for depression: A systematic review and individual patient data meta-analysis.. <i>Journal of Affective Disorders</i> , 2021 , 299, 298-308	6.6	0
42	Standard smoking cessation services in sites participating in the SCIMITAR+ trial for people with severe mental ill health. <i>BJPsych Bulletin</i> , 2019 , 6-11	1.2	0
41	Health of mothers of children with a life-limiting condition: a protocol for comparative cohort study using the Clinical Practice Research Datalink. <i>BMJ Open</i> , 2020 , 10, e034024	3	0
40	Effectiveness of collaborative care in reducing suicidal ideation: An individual participant data meta-analysis. <i>General Hospital Psychiatry</i> , 2021 , 71, 27-35	5.6	0
39	Long-Term Cost-Effectiveness of Smoking Cessation Interventions in People With Mental Disorders: A Dynamic Decision Analytical Model. <i>Value in Health</i> , 2021 , 24, 1263-1272	3.3	0
38	A psychological intervention by community pharmacies to prevent depression in adults with subthreshold depression and long-term conditions: the CHEMIST pilot RCT. <i>Public Health Research</i> , 2022 , 10, 1-160	1.7	0

37	Behavioural Activation for Social IsoLation (BASIL+) trial (Behavioural activation to mitigate depression and loneliness among older people with long-term conditions): Protocol for a fully-powered pragmatic randomised controlled trial.. <i>PLoS ONE</i> , 2022 , 17, e0263856	3.7	o
36	The DIAMONDS intervention to support self-management of type 2 diabetes in people with severe mental illness: Study protocol for a single-group feasibility study. <i>SSM Mental Health</i> , 2022 , 2, 100086		o
35	Cognitive, behavioural or cognitive-behavioural self-help interventions for subclinical depression in older adults: A systematic review and meta-analysis.. <i>Journal of Affective Disorders</i> , 2022 , 308, 384-390	6.6	o
34	Challenges and solutions to nicotine replacement therapy access: observations from SCIMITAR. <i>BJPsych Open</i> , 2020 , 6, e120	5	
33	Out of pocket expenses in obsessive compulsive disorder. <i>Journal of Mental Health</i> , 2020 , 1-6	2.7	
32	Computerised Cognitive Behaviour Therapy for Depression Management: A Cost-Effectiveness Analysis. <i>Value in Health</i> , 2014 , 17, A459	3.3	
31	M9 Lung Health of OPIate Users (LHOPI): A pilot study to assess the respiratory health of opiate misusers attending a community substance misuse clinic. <i>Thorax</i> , 2015 , 70, A230.2-A231	7.3	
30	P1-294 Risk factors for poor mental health during pregnancy from the born in Bradford cohort: early results from a bi-ethnic sample. <i>Journal of Epidemiology and Community Health</i> , 2011 , 65, A148-A148	5.1	
29	P1-496 An examination of the psychometric properties of the GHQ-28 subscales in a bi-ethnic antenatal sample in the UK: preliminary results from the born in Bradford cohort. <i>Journal of Epidemiology and Community Health</i> , 2011 , 65, A204-A204	5.1	
28	Linking evidence to practice31-54		
27	A stepped care intervention is effective for elderly people with subthreshold depression or anxiety. <i>Evidence-Based Mental Health</i> , 2009 , 12, 115	11.1	
26	Telephone treatment support improves outcomes for depressed employees. <i>Evidence-Based Mental Health</i> , 2008 , 11, 47	11.1	
25	A smoking ban in psychiatric units: threat or opportunity?. <i>Mental Health in Family Medicine</i> , 2008 , 5, 129-33		
24	Cost-effectiveness of brief structured interventions to discontinue long-term benzodiazepine use: an economic analysis alongside a randomised controlled trial. <i>HRB Open Research</i> , 2020 , 3, 33	1.2	
23	Collaborative care for depression in older adults: How much is enough?. <i>Behaviour Research and Therapy</i> , 2020 , 135, 103725	5.2	
22	Infection in people with severe mental illness - Authors' reply. <i>Lancet Psychiatry,the</i> , 2016 , 3, 204-5	23.3	
21	Improving Access to Care386-400		
20	Collaborative Care and Stepped Care401-419		

- 19 Multiple versus single risk behaviour interventions for people with severe mental illness: a network meta-analysis and qualitative synthesis **2022**, 10, 1-186
- 18 Models of care for depression 17-30
- 17 Anatomy of a review 55-72
- 16 Collaborative care 87-102
- 15 Referral 103-110
- 14 Summary of the evidence 111-116
- 13 Making it happen 117-126
- 12 United States perspective 137-144
- 11 Psychological and pharmacological interventions for posttraumatic stress disorder and comorbid mental health problems following complex traumatic events: Systematic review and component network meta-analysis **2020**, 17, e1003262
- 10 Psychological and pharmacological interventions for posttraumatic stress disorder and comorbid mental health problems following complex traumatic events: Systematic review and component network meta-analysis **2020**, 17, e1003262
- 9 Psychological and pharmacological interventions for posttraumatic stress disorder and comorbid mental health problems following complex traumatic events: Systematic review and component network meta-analysis **2020**, 17, e1003262
- 8 Psychological and pharmacological interventions for posttraumatic stress disorder and comorbid mental health problems following complex traumatic events: Systematic review and component network meta-analysis **2020**, 17, e1003262
- 7 Psychological and pharmacological interventions for posttraumatic stress disorder and comorbid mental health problems following complex traumatic events: Systematic review and component network meta-analysis **2020**, 17, e1003262
- 6 Frailty and depression predict instrumental activities of daily living in older adults: A population-based longitudinal study using the CARE75+ cohort **2020**, 15, e0243972
- 5 Frailty and depression predict instrumental activities of daily living in older adults: A population-based longitudinal study using the CARE75+ cohort **2020**, 15, e0243972
- 4 Frailty and depression predict instrumental activities of daily living in older adults: A population-based longitudinal study using the CARE75+ cohort **2020**, 15, e0243972
- 3 Frailty and depression predict instrumental activities of daily living in older adults: A population-based longitudinal study using the CARE75+ cohort **2020**, 15, e0243972
- 2 Frailty and depression predict instrumental activities of daily living in older adults: A population-based longitudinal study using the CARE75+ cohort **2020**, 15, e0243972

- 1 Frailty and depression predict instrumental activities of daily living in older adults: A population-based longitudinal study using the CARE75+ cohort **2020**, 15, e0243972