Simon Gilbody

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/6337009/simon-gilbody-publications-by-year.pdf

Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

378	19,754	70	134
papers	citations	h-index	g-index
427	24,462 ext. citations	5.5	7.02
ext. papers		avg, IF	L-index

#	Paper	IF	Citations
378	Loneliness among people with severe mental illness during the COVID-19 pandemic: Results from a linked UK population cohort study <i>PLoS ONE</i> , 2022 , 17, e0262363	3.7	5
377	Cost-utility analysis of LEGO based therapy for school children and young people with autism spectrum disorder: results from a randomised controlled trial <i>BMJ Open</i> , 2022 , 12, e056347	3	1
376	Predicting relapse or recurrence of depression: systematic review of prognostic models <i>British Journal of Psychiatry</i> , 2022 , 1-11	5.4	O
375	The feasibility and acceptability of a brief psychological intervention for adults with long-term health conditions and subthreshold depression delivered via community pharmacies: a mixed methods evaluation-the Community Pharmacies Mood Intervention Study (CHEMIST) Pilot and	1.9	1
374	A psychological intervention by community pharmacies to prevent depression in adults with subthreshold depression and long-term conditions: the CHEMIST pilot RCT. <i>Public Health Research</i> , 2022 , 10, 1-160	1.7	O
373	Socioeconomic Indicators of Treatment Prognosis for Adults With Depression: A Systematic Review and Individual Patient Data Meta-analysis <i>JAMA Psychiatry</i> , 2022 ,	14.5	2
372	Behavioural Activation for Social IsoLation (BASIL+) trial (Behavioural activation to mitigate depression and loneliness among older people with long-term conditions): Protocol for a fully-powered pragmatic randomised controlled trial <i>PLoS ONE</i> , 2022 , 17, e0263856	3.7	O
371	The DIAMONDS intervention to support self-management of type 2 diabetes in people with severe mental illness: Study protocol for a single-group feasibility study. <i>SSM Mental Health</i> , 2022 , 2, 100086		O
370	Multiple versus single risk behaviour interventions for people with severe mental illness: a network meta-analysis and qualitative synthesis 2022 , 10, 1-186		
369	Cognitive, behavioural or cognitive-behavioural self-help interventions for subclinical depression in older adults: A systematic review and meta-analysis <i>Journal of Affective Disorders</i> , 2022 , 308, 384-390	6.6	O
368	A Year Into the Pandemic: The Diversity of Experience Amongst People With Severe Mental Ill Health <i>Frontiers in Psychiatry</i> , 2021 , 12, 794585	5	1
367	A Patient Stratification Approach to Identifying the Likelihood of Continued Chronic Depression and Relapse Following Treatment for Depression <i>Journal of Personalized Medicine</i> , 2021 , 11,	3.6	2
366	Cost-Utility Analysis of Discontinuing Antidepressants in England Primary Care Patients Compared with Long-Term Maintenance: The ANTLER Study. <i>Applied Health Economics and Health Policy</i> , 2021 , 1	3.4	1
365	Antidepressant medication to prevent depression relapse in primary care: the ANTLER RCT. <i>Health Technology Assessment</i> , 2021 , 25, 1-62	4.4	O
364	Life events and treatment prognosis for depression: A systematic review and individual patient data meta-analysis <i>Journal of Affective Disorders</i> , 2021 , 299, 298-308	6.6	O
363	Health risk behaviours among people with severe mental ill health during the COVID-19 pandemic: Analysis of linked cohort data. <i>PLoS ONE</i> , 2021 , 16, e0258349	3.7	1
362	A systematic review of behavioural smoking cessation interventions for people with severe mental ill health-what works?. <i>Addiction</i> , 2021 ,	4.6	1

(2021-2021)

361	Behavioural activation to prevent depression and loneliness among socially isolated older people with long-term conditions: The BASIL COVID-19 pilot randomised controlled trial. <i>PLoS Medicine</i> , 2021 , 18, e1003779	11.6	3	
360	Supporting smoking cessation and preventing relapse following a stay in a smoke-free setting: a meta-analysis and investigation of effective behaviour change techniques. <i>Addiction</i> , 2021 , 116, 2978-2	2944	2	
359	Health of mothers of children with a life-limiting condition: a comparative cohort study. <i>Archives of Disease in Childhood</i> , 2021 , 106, 987-993	2.2	6	
358	Investigating smoking and nicotine dependence among people with severe mental illness during the COVID-19 pandemic: analysis of linked data from a UK Closing the Gap cohort. <i>BJPsych Open</i> , 2021 , 7, e86	5	2	
357	Internet-Based Cognitive Behavioral Therapy for Depression: A Systematic Review and Individual Patient Data Network Meta-analysis. <i>JAMA Psychiatry</i> , 2021 , 78, 361-371	14.5	97	
356	The contribution of depressive 'disorder characteristics' to determinations of prognosis for adults with depression: an individual patient data meta-analysis. <i>Psychological Medicine</i> , 2021 , 51, 1068-1081	6.9	7	
355	The digital divide: amplifying health inequalities for people with severe mental illness in the time of COVID-19. <i>British Journal of Psychiatry</i> , 2021 , 219, 529-531	5.4	4	
354	Predicting prognosis for adults with depression using individual symptom data: a comparison of modelling approaches. <i>Psychological Medicine</i> , 2021 , 1-11	6.9	7	
353	Shared goals for mental health research: what, why and when for the 2020s. <i>Journal of Mental Health</i> , 2021 , 1-9	2.7	6	
352	Living with diabetes alongside a severe mental illness: A qualitative exploration with people with severe mental illness, family members and healthcare staff. <i>Diabetic Medicine</i> , 2021 , 38, e14562	3.5	6	
351	The importance of transdiagnostic symptom level assessment to understanding prognosis for depressed adults: analysis of data from six randomised control trials. <i>BMC Medicine</i> , 2021 , 19, 109	11.4	2	
350	Prognostic models for predicting relapse or recurrence of major depressive disorder in adults. <i>The Cochrane Library</i> , 2021 , 5, CD013491	5.2	4	
349	Identifying determinants of diabetes risk and outcomes for people with severe mental illness: a mixed-methods study. <i>Health Services and Delivery Research</i> , 2021 , 9, 1-194	1.5	1	
348	Can We Prevent Depression in At-Risk Older Adults Using Self-Help? The UK SHARD Trial of Behavioral Activation. <i>American Journal of Geriatric Psychiatry</i> , 2021 ,	6.5	2	
347	Barriers and facilitators to implementing perinatal mental health care in health and social care settings: a systematic review. <i>Lancet Psychiatry,the</i> , 2021 , 8, 521-534	23.3	5	
346	Dismantling, optimising, and personalising internet cognitive behavioural therapy for depression: a systematic review and component network meta-analysis using individual participant data. <i>Lancet Psychiatry,the</i> , 2021 , 8, 500-511	23.3	27	
345	Cost-effectiveness of feedback-informed psychological treatment: Evidence from the IAPT-FIT trial. <i>Behaviour Research and Therapy</i> , 2021 , 142, 103873	5.2	5	
344	Effectiveness of collaborative care in reducing suicidal ideation: An individual participant data meta-analysis. <i>General Hospital Psychiatry</i> , 2021 , 71, 27-35	5.6	О	

343	The development and validation of a prognostic model to PREDICT Relapse of depression in adult patients in primary care: protocol for the PREDICTR study. <i>Diagnostic and Prognostic Research</i> , 2021 , 5, 12	5.5	1
342	Depression prevalence based on the Edinburgh Postnatal Depression Scale compared to Structured Clinical Interview for DSM Disorders classification: Systematic review and individual participant data meta-analysis. <i>International Journal of Methods in Psychiatric Research</i> , 2021 , 30, e1860	4.3	11
341	Preventing depression among older people living in rural areas: A randomised controlled trial of behavioural activation in collaborative care. <i>International Journal of Geriatric Psychiatry</i> , 2021 , 36, 530-5	39 9	3
340	Probability of Major Depression Classification Based on the SCID, CIDI, and MINI Diagnostic Interviews: A Synthesis of Three Individual Participant Data Meta-Analyses. <i>Psychotherapy and Psychosomatics</i> , 2021 , 90, 28-40	9.4	9
339	Smoking cessation in severe mental illness: combined long-term quit rates from the UK SCIMITAR trials programme. <i>British Journal of Psychiatry</i> , 2021 , 218, 95-97	5.4	8
338	Impact of severe mental illness on healthcare use and health outcomes for people with type 2 diabetes: a longitudinal observational study in England. <i>British Journal of General Practice</i> , 2021 , 71, e56	5-657	3 ²
337	Is social support pre-treatment associated with prognosis for adults with depression in primary care?. <i>Acta Psychiatrica Scandinavica</i> , 2021 , 143, 392-405	6.5	5
336	How can we estimate QALYs based on PHQ-9 scores? Equipercentile linking analysis of PHQ-9 and EQ-5D. <i>Evidence-Based Mental Health</i> , 2021 ,	11.1	4
335	Home environmental assessments and modification delivered by occupational therapists to reduce falls in people aged 65 years and over: the OTIS RCT. <i>Health Technology Assessment</i> , 2021 , 25, 1-118	4.4	2
334	Still 'being bothered about Billy': managing the physical health of people with severe mental illness. <i>British Journal of General Practice</i> , 2021 , 71, 373-376	1.6	1
333	Mitigating the impacts of COVID-19: where are the mental health trials?. <i>Lancet Psychiatry,the</i> , 2021 , 8, 647-650	23.3	6
332	Long-Term Cost-Effectiveness of Smoking Cessation Interventions in People With Mental Disorders: A Dynamic Decision Analytical Model. <i>Value in Health</i> , 2021 , 24, 1263-1272	3.3	O
331	Data-driven methods distort optimal cutoffs and accuracy estimates of depression screening tools: a simulation study using individual participant data. <i>Journal of Clinical Epidemiology</i> , 2021 , 137, 137-147	5.7	3
330	Healthcare resource use and costs for people with type 2 diabetes mellitus with and without severe mental illness in England: longitudinal matched-cohort study using the Clinical Practice Research Datalink <i>British Journal of Psychiatry</i> , 2021 , 1-8	5.4	1
329	Use of the Internet and Digital Devices Among People With Severe Mental Ill Health During the COVID-19 Pandemic Restrictions. <i>Frontiers in Psychiatry</i> , 2021 , 12, 732735	5	6
328	Maintenance or Discontinuation of Antidepressants in Primary Care. <i>New England Journal of Medicine</i> , 2021 , 385, 1257-1267	59.2	16
327	Nature-based outdoor activities for mental and physical health: Systematic review and meta-analysis. <i>SSM - Population Health</i> , 2021 , 16, 100934	3.8	14
326	Role of age, gender and marital status in prognosis for adults with depression: An individual patient data meta-analysis. <i>Epidemiology and Psychiatric Sciences</i> , 2021 , 30, e42	5.1	11

(2020-2021)

325	Shortening the Edinburgh postnatal depression scale using optimal test assembly methods: Development of the EPDS-Dep-5. <i>Acta Psychiatrica Scandinavica</i> , 2021 , 143, 348-362	6.5	1
324	Cognitive or behavioural interventions (or both) to prevent or mitigate loneliness in adolescents, adults, and older adults. <i>The Cochrane Library</i> , 2020 ,	5.2	1
323	Challenges and solutions to nicotine replacement therapy access: observations from SCIMITAR. <i>BJPsych Open</i> , 2020 , 6, e120	5	
322	How much change is enough? Evidence from a longitudinal study on depression in UK primary care. <i>Psychological Medicine</i> , 2020 , 1-8	6.9	14
321	Out of pocket expenses in obsessive compulsive disorder. <i>Journal of Mental Health</i> , 2020 , 1-6	2.7	
320	REDUCE (Reviewing long-term antidepressant use by careful monitoring in everyday practice) internet and telephone support to people coming off long-term antidepressants: protocol for a randomised controlled trial. <i>Trials</i> , 2020 , 21, 419	2.8	6
319	Individual participant data (IPD) meta-analysis of psychological relapse prevention interventions versus control for patients in remission from depression: a protocol. <i>BMJ Open</i> , 2020 , 10, e034158	3	4
318	Accuracy of the PHQ-2 Alone and in Combination With the PHQ-9 for Screening to Detect Major Depression: Systematic Review and Meta-analysis. <i>JAMA - Journal of the American Medical Association</i> , 2020 , 323, 2290-2300	27.4	86
317	E-cigarette use and associated factors among smokers with severe mental illness. <i>Addictive Behaviors</i> , 2020 , 108, 106456	4.2	1
316	Factors that influence participation in physical activity for anxiety or depression: a synthesis of qualitative evidence. <i>The Cochrane Library</i> , 2020 ,	5.2	O
315	Patient Health Questionnaire-9 scores do not accurately estimate depression prevalence: individual participant data meta-analysis. <i>Journal of Clinical Epidemiology</i> , 2020 , 122, 115-128.e1	5.7	58
314	Estimating the sample mean and standard deviation from commonly reported quantiles in meta-analysis. <i>Statistical Methods in Medical Research</i> , 2020 , 962280219889080	2.3	104
313	A systematic review of mental health measurement scales for evaluating the effects of mental health prevention interventions. <i>European Journal of Public Health</i> , 2020 , 30, 539-545	2.1	5
312	Optimised patient information materials and recruitment to a study of behavioural activation in older adults: an embedded study within a trial. <i>F1000Research</i> , 2020 , 9, 417	3.6	3
311	Frailty and depression predict instrumental activities of daily living in older adults: A population-based longitudinal study using the CARE75+ cohort. <i>PLoS ONE</i> , 2020 , 15, e0243972	3.7	8
310	Interventions for adults with a history of complex traumatic events: the INCiTE mixed-methods systematic review. <i>Health Technology Assessment</i> , 2020 , 24, 1-312	4.4	3
309	The association between primary care quality and health-care use, costs and outcomes for people with serious mental illness: a retrospective observational study. <i>Health Services and Delivery Research</i> , 2020 , 8, 1-126	1.5	1
308	Cost-effectiveness of brief structured interventions to discontinue long-term benzodiazepine use: an economic analysis alongside a randomised controlled trial. <i>HRB Open Research</i> , 2020 , 3, 33	1.2	

307	What does the rise and rise of electronic nicotine delivery systems mean for mental health services?. <i>Lancet Psychiatry,the</i> , 2020 , 7, 11-13	23.3	2
306	Probability of major depression diagnostic classification based on the SCID, CIDI and MINI diagnostic interviews controlling for Hospital Anxiety and Depression Scale - Depression subscale scores: An individual participant data meta-analysis of 73 primary studies. <i>Journal of Psychosomatic</i>	4.1	22
305	The role of relapse prevention for depression in collaborative care: A systematic review. <i>Journal of Affective Disorders</i> , 2020 , 265, 618-644	6.6	8
304	Collaborative care for depression in older adults: How much is enough?. <i>Behaviour Research and Therapy</i> , 2020 , 135, 103725	5.2	
303	Health of mothers of children with a life-limiting condition: a protocol for comparative cohort study using the Clinical Practice Research Datalink. <i>BMJ Open</i> , 2020 , 10, e034024	3	O
302	The associations between loneliness, social exclusion and pain in the general population: A N=502,528 cross-sectional UK Biobank study. <i>Journal of Psychiatric Research</i> , 2020 , 130, 68-74	5.2	11
301	Conversion and neuro-inflammation disorder observational study (CANDO). Protocol of a feasibility study. <i>European Journal of Psychiatry</i> , 2020 , 34, 164-172	1	2
300	Overestimation of Postpartum Depression Prevalence Based on a 5-item Version of the EPDS: Systematic Review and Individual Participant Data Meta-analysis. <i>Canadian Journal of Psychiatry</i> , 2020 , 65, 835-844	4.8	3
299	A meta-review of "lifestyle psychiatry": the role of exercise, smoking, diet and sleep in the prevention and treatment of mental disorders. <i>World Psychiatry</i> , 2020 , 19, 360-380	14.4	132
298	Psychological and pharmacological interventions for posttraumatic stress disorder and comorbid mental health problems following complex traumatic events: Systematic review and component network meta-analysis. <i>PLoS Medicine</i> , 2020 , 17, e1003262	11.6	71
297	Effectiveness and adherence of telephone-administered psychotherapy for depression: A systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2020 , 260, 514-526	6.6	24
296	The Accuracy of the Patient Health Questionnaire-9 Algorithm for Screening to Detect Major Depression: An Individual Participant Data Meta-Analysis. <i>Psychotherapy and Psychosomatics</i> , 2020 , 89, 25-37	9.4	42
295	Equivalency of the diagnostic accuracy of the PHQ-8 and PHQ-9: a systematic review and individual participant data meta-analysis. <i>Psychological Medicine</i> , 2020 , 50, 1368-1380	6.9	65
294	Cost-effectiveness of a specialist smoking cessation package compared with standard smoking cessation services for people with severe mental illness in England: a trial-based economic evaluation from the SCIMITAR+ study. <i>Addiction</i> , 2020 , 115, 2113-2122	4.6	5
293	Psychological and pharmacological interventions for posttraumatic stress disorder and comorbid mental health problems following complex traumatic events: Systematic review and component network meta-analysis 2020 , 17, e1003262		
292	Psychological and pharmacological interventions for posttraumatic stress disorder and comorbid mental health problems following complex traumatic events: Systematic review and component network meta-analysis 2020 , 17, e1003262		
291	Psychological and pharmacological interventions for posttraumatic stress disorder and comorbid mental health problems following complex traumatic events: Systematic review and component network meta-analysis 2020 , 17, e1003262		
290	Psychological and pharmacological interventions for posttraumatic stress disorder and comorbid mental health problems following complex traumatic events: Systematic review and component network meta-analysis 2020 , 17, e1003262		

(2019-2020)

289	Psychological and pharmacological interventions for posttraumatic stress disorder and comorbid mental health problems following complex traumatic events: Systematic review and component network meta-analysis 2020 , 17, e1003262		
288	Frailty and depression predict instrumental activities of daily living in older adults: A population-based longitudinal study using the CARE75+ cohort 2020 , 15, e0243972		
287	Frailty and depression predict instrumental activities of daily living in older adults: A population-based longitudinal study using the CARE75+ cohort 2020 , 15, e0243972		
286	Frailty and depression predict instrumental activities of daily living in older adults: A population-based longitudinal study using the CARE75+ cohort 2020 , 15, e0243972		
285	Frailty and depression predict instrumental activities of daily living in older adults: A population-based longitudinal study using the CARE75+ cohort 2020 , 15, e0243972		
284	Frailty and depression predict instrumental activities of daily living in older adults: A population-based longitudinal study using the CARE75+ cohort 2020 , 15, e0243972		
283	Frailty and depression predict instrumental activities of daily living in older adults: A population-based longitudinal study using the CARE75+ cohort 2020 , 15, e0243972		
282	Comparison of major depression diagnostic classification probability using the SCID, CIDI, and MINI diagnostic interviews among women in pregnancy or postpartum: An individual participant data meta-analysis. <i>International Journal of Methods in Psychiatric Research</i> , 2019 , 28, e1803	4.3	24
281	The clinical effectiveness of sertraline in primary care and the role of depression severity and duration (PANDA): a pragmatic, double-blind, placebo-controlled randomised trial. <i>Lancet Psychiatry,the</i> , 2019 , 6, 903-914	23.3	52
280	Can support workers from AgeUK deliver an intervention to support older people with anxiety and depression? A qualitative evaluation. <i>BMC Family Practice</i> , 2019 , 20, 16	2.6	3
279	Impact of telephone delivered case-management on the effectiveness of collaborative care for depression and anti-depressant use: A systematic review and meta-regression. <i>PLoS ONE</i> , 2019 , 14, e02	137948	9
278	A randomised controlled trial assessing the use of citalopram, sertraline, fluoxetine and mirtazapine in preventing relapse in primary care patients who are taking long-term maintenance antidepressants (ANTLER: ANTidepressants to prevent reLapse in dEpRession): study protocol for a	2.8	7
277	Community Pharmacies Mood Intervention Study (CHEMIST): feasibility and external pilot randomised controlled trial protocol. <i>Pilot and Feasibility Studies</i> , 2019 , 5, 71	1.9	3
276	Everything you wanted to know about e-cigarettes and vaping but were afraid to ask: a guide for mental health clinicians. <i>BJ Psych Advances</i> , 2019 , 25, 279-286	0.8	3
275	Association Between Antipsychotic Polypharmacy and Outcomes for People With Serious Mental Illness in England. <i>Psychiatric Services</i> , 2019 , 70, 650-656	3.3	12
274	Smoking cessation for people with severe mental illness (SCIMITAR+): a pragmatic randomised controlled trial. <i>Lancet Psychiatry,the</i> , 2019 , 6, 379-390	23.3	75
273	Accuracy of Patient Health Questionnaire-9 (PHQ-9) for screening to detect major depression: individual participant data meta-analysis. <i>BMJ, The</i> , 2019 , 365, l1476	5.9	374
272	The Effects of Meditation, Yoga, and Mindfulness on Depression, Anxiety, and Stress in Tertiary Education Students: A Meta-Analysis. <i>Frontiers in Psychiatry</i> , 2019 , 10, 193	5	47

271	Commentary on: Suicide Screening Tools for Use with Incarcerated Offenders. <i>Archives of Suicide Research</i> , 2019 , 23, 523-525	2.3	1
270	Investigating SOcial Competence and Isolation in children with Autism taking part in LEGO-based therapy clubs In School Environments (I-SOCIALISE): study protocol. <i>BMJ Open</i> , 2019 , 9, e030471	3	4
269	SCIMITAR+ Trial: A randomised study within a trial (SWAT) of a contingent financial reward to improve trial follow-up. <i>F1000Research</i> , 2019 , 8, 1937	3.6	2
268	What factors indicate prognosis for adults with depression in primary care? A protocol for meta-analyses of individual patient data using the Dep-GP database. <i>Wellcome Open Research</i> , 2019 , 4, 69	4.8	11
267	Exploring Severe Mental Illness and Diabetes: Protocol for a Longitudinal, Observational, and Qualitative Mixed Methods Study. <i>JMIR Research Protocols</i> , 2019 , 8, 13407	2	3
266	A bespoke smoking cessation service compared with treatment as usual for people with severe mental ill health: the SCIMITAR+ RCT. <i>Health Technology Assessment</i> , 2019 , 23, 1-116	4.4	6
265	Antidepressant treatment with sertraline for adults with depressive symptoms in primary care: the PANDA research programme including RCT. <i>Programme Grants for Applied Research</i> , 2019 , 7, 1-108	1.5	3
264	What factors indicate prognosis for adults with depression in primary care? A protocol for meta-analyses of individual patient data using the Dep-GP database. <i>Wellcome Open Research</i> , 2019 , 4, 69	4.8	9
263	Non-traditional support workers delivering a brief psychosocial intervention for older people with anxiety and depression: the NOTEPAD feasibility study. <i>Health Services and Delivery Research</i> , 2019 , 7, 1-120	1.5	1
262	SCIMITAR+ Trial: A randomised study within a trial (SWAT) of a contingent financial reward to improve trial follow-up. <i>F1000Research</i> , 2019 , 8, 1937	3.6	2
261	Standard smoking cessation services in sites participating in the SCIMITAR+ trial for people with severe mental ill health. <i>BJPsych Bulletin</i> , 2019 , 6-11	1.2	О
260	Behavioural activation in nursing homes to treat depression (BAN-Dep): study protocol for a pragmatic randomised controlled trial. <i>BMJ Open</i> , 2019 , 9, e032421	3	2
259	Prognostic models for predicting relapse or recurrence of depression. The Cochrane Library, 2019,	5.2	78
258	Antidepressants for major depression disorder in older people: a network meta-analysis. <i>The Cochrane Library</i> , 2019 ,	5.2	3
257	Pharmacy-based management for depression in adults. <i>The Cochrane Library</i> , 2019 , 12, CD013299	5.2	10
256	Impact of family practice continuity of care on unplanned hospital use for people with serious mental illness. <i>Health Services Research</i> , 2019 , 54, 1316-1325	3.4	15
255	Factors associated with regular physical activity participation among people with severe mental ill health. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2019 , 54, 887-895	4.5	12
254	Predicting persistent depressive symptoms in older adults: A machine learning approach to personalised mental healthcare. <i>Journal of Affective Disorders</i> , 2019 , 246, 857-860	6.6	33

253	GPs' attitudes towards digital technologies for depression: an online survey in primary care. <i>British Journal of General Practice</i> , 2019 , 69, e164-e170	1.6	10
252	Shortening self-report mental health symptom measures through optimal test assembly methods: Development and validation of the Patient Health Questionnaire-Depression-4. <i>Depression and Anxiety</i> , 2019 , 36, 82-92	8.4	12
251	Relapse and Recurrence of Common Mental Health Problems after Low Intensity Cognitive Behavioural Therapy: The WYLOW Longitudinal Cohort Study. <i>Psychotherapy and Psychosomatics</i> , 2018 , 87, 116-117	9.4	6
250	A feasibility study for NOn-Traditional providers to support the management of Elderly People with Anxiety and Depression: The NOTEPAD study Protocol. <i>Trials</i> , 2018 , 19, 172	2.8	3
249	Probability of major depression diagnostic classification using semi-structured versus fully structured diagnostic interviews. <i>British Journal of Psychiatry</i> , 2018 , 212, 377-385	5.4	36
248	Is self-guided internet-based cognitive behavioural therapy (iCBT) harmful? An individual participant data meta-analysis. <i>Psychological Medicine</i> , 2018 , 48, 2456-2466	6.9	60
247	German EstSmoke: estimating adult smoking-related costs and consequences of smoking cessation for Germany. <i>Addiction</i> , 2018 , 113, 125-136	4.6	3
246	Feedback-informed treatment versus usual psychological treatment for depression and anxiety: a multisite, open-label, cluster randomised controlled trial. <i>Lancet Psychiatry,the</i> , 2018 , 5, 564-572	23.3	52
245	Loneliness, social isolation and risk of cardiovascular disease in the English Longitudinal Study of Ageing. <i>European Journal of Preventive Cardiology</i> , 2018 , 25, 1387-1396	3.9	86
244	Successful recruitment to trials: findings from the SCIMITAR+ Trial. <i>Trials</i> , 2018 , 19, 53	2.8	9
244	Successful recruitment to trials: findings from the SCIMITAR+ Trial. <i>Trials</i> , 2018 , 19, 53 Identifying perinatal depression with case-finding instruments: a mixed-methods study (BaBY PaNDA Born and Bred in Yorkshire PeriNatal Depression Diagnostic Accuracy). <i>Health Services and Delivery Research</i> , 2018 , 6, 1-210	2.8	9
	Identifying perinatal depression with case-finding instruments: a mixed-methods study (BaBY PaNDA Born and Bred in Yorkshire PeriNatal Depression Diagnostic Accuracy). <i>Health Services and</i>		
243	Identifying perinatal depression with case-finding instruments: a mixed-methods study (BaBY PaNDA Born and Bred in Yorkshire PeriNatal Depression Diagnostic Accuracy). <i>Health Services and Delivery Research</i> , 2018 , 6, 1-210 Combining mirtazapine with SSRIs or SNRIs for treatment-resistant depression: the MIR RCT. <i>Health</i>	1.5	8
243	Identifying perinatal depression with case-finding instruments: a mixed-methods study (BaBY PaNDA [Born and Bred in Yorkshire PeriNatal Depression Diagnostic Accuracy). <i>Health Services and Delivery Research</i> , 2018 , 6, 1-210 Combining mirtazapine with SSRIs or SNRIs for treatment-resistant depression: the MIR RCT. <i>Health Technology Assessment</i> , 2018 , 22, 1-136 Collaborative case management to aid return to work after long-term sickness absence: a pilot	1.5 4.4	8
243 242 241	Identifying perinatal depression with case-finding instruments: a mixed-methods study (BaBY PaNDA IBorn and Bred in Yorkshire PeriNatal Depression Diagnostic Accuracy). <i>Health Services and Delivery Research</i> , 2018 , 6, 1-210 Combining mirtazapine with SSRIs or SNRIs for treatment-resistant depression: the MIR RCT. <i>Health Technology Assessment</i> , 2018 , 22, 1-136 Collaborative case management to aid return to work after long-term sickness absence: a pilot randomised controlled trial. <i>Public Health Research</i> , 2018 , 6, 1-76 Do care plans and annual reviews of physical health influence unplanned hospital utilisation for people with serious mental illness? Analysis of linked longitudinal primary and secondary	1.5 4.4 1.7	8 16 1
243 242 241 240	Identifying perinatal depression with case-finding instruments: a mixed-methods study (BaBY PaNDA [Born and Bred in Yorkshire PeriNatal Depression Diagnostic Accuracy). <i>Health Services and Delivery Research</i> , 2018 , 6, 1-210 Combining mirtazapine with SSRIs or SNRIs for treatment-resistant depression: the MIR RCT. <i>Health Technology Assessment</i> , 2018 , 22, 1-136 Collaborative case management to aid return to work after long-term sickness absence: a pilot randomised controlled trial. <i>Public Health Research</i> , 2018 , 6, 1-76 Do care plans and annual reviews of physical health influence unplanned hospital utilisation for people with serious mental illness? Analysis of linked longitudinal primary and secondary healthcare records in England. <i>BMJ Open</i> , 2018 , 8, e023135 Diagnostic accuracy of the Geriatric Depression Scale-30, Geriatric Depression Scale-15, Geriatric Depression Scale-5 and Geriatric Depression Scale-4 for detecting major depression: protocol for a	1.5 4.4 1.7	8 16 1
243 242 241 240 239	Identifying perinatal depression with case-finding instruments: a mixed-methods study (BaBY PaNDA IBorn and Bred in Yorkshire PeriNatal Depression Diagnostic Accuracy). <i>Health Services and Delivery Research</i> , 2018 , 6, 1-210 Combining mirtazapine with SSRIs or SNRIs for treatment-resistant depression: the MIR RCT. <i>Health Technology Assessment</i> , 2018 , 22, 1-136 Collaborative case management to aid return to work after long-term sickness absence: a pilot randomised controlled trial. <i>Public Health Research</i> , 2018 , 6, 1-76 Do care plans and annual reviews of physical health influence unplanned hospital utilisation for people with serious mental illness? Analysis of linked longitudinal primary and secondary healthcare records in England. <i>BMJ Open</i> , 2018 , 8, e023135 Diagnostic accuracy of the Geriatric Depression Scale-30, Geriatric Depression Scale-15, Geriatric Depression Scale-5 and Geriatric Depression Scale-4 for detecting major depression: protocol for a systematic review and individual participant data meta-analysis. <i>BMJ Open</i> , 2018 , 8, e026598 The HARMONIC trial: study protocol for a randomised controlled feasibility trial of Shaping Healthy Minds-a modular transdiagnostic intervention for mood, stressor-related and anxiety disorders in	1.5 4.4 1.7 3	8 16 1 4

235	Effect of Collaborative Care vs Usual Care on Depressive Symptoms in Older Adults With Subthreshold Depression: The CASPER Randomized Clinical Trial. <i>JAMA - Journal of the American Medical Association</i> , 2017 , 317, 728-737	27.4	61
234	Efficacy of Self-guided Internet-Based Cognitive Behavioral Therapy in the Treatment of Depressive Symptoms: A Meta-analysis of Individual Participant Data. <i>JAMA Psychiatry</i> , 2017 , 74, 351-3	5 9 4·5	346
233	Cost-effectiveness of computerized cognitive-behavioural therapy for the treatment of depression in primary care: findings from the Randomised Evaluation of the Effectiveness and Acceptability of Computerised Therapy (REEACT) trial. <i>Psychological Medicine</i> , 2017 , 47, 1825-1835	6.9	11
232	Telephone-supported computerised cognitive-behavioural therapy: REEACT-2 large-scale pragmatic randomised controlled trial. <i>British Journal of Psychiatry</i> , 2017 , 210, 362-367	5.4	57
231	How durable is the effect of low intensity CBT for depression and anxiety? Remission and relapse in a longitudinal cohort study. <i>Behaviour Research and Therapy</i> , 2017 , 94, 1-8	5.2	55
230	Selective Cutoff Reporting in Studies of Diagnostic Test Accuracy: A Comparison of Conventional and Individual-Patient-Data Meta-Analyses of the Patient Health Questionnaire-9 Depression Screening Tool. <i>American Journal of Epidemiology</i> , 2017 , 185, 954-964	3.8	32
229	Fathers' views and experiences of their own mental health during pregnancy and the first postnatal year: a qualitative interview study of men participating in the UK Born and Bred in Yorkshire (BaBY) cohort. <i>BMC Pregnancy and Childbirth</i> , 2017 , 17, 45	3.2	90
228	Impact of long-term medical conditions on the outcomes of psychological therapy for depression and anxiety. <i>British Journal of Psychiatry</i> , 2017 , 210, 47-53	5.4	16
227	Are there researcher allegiance effects in diagnostic validation studies of the PHQ-9? A systematic review and meta-analysis. <i>BMJ Open</i> , 2017 , 7, e015247	3	11
226	Improving the efficiency of psychological treatment using outcome feedback technology. <i>Behaviour Research and Therapy</i> , 2017 , 99, 89-97	5.2	35
225	Should Google offer an online screening test for depression?. <i>BMJ, The</i> , 2017 , 358, j4144	5.9	3
224	The Effectiveness of Pharmacological and Non-Pharmacological Interventions for Improving Glycaemic Control in Adults with Severe Mental Illness: A Systematic Review and Meta-Analysis. <i>PLoS ONE</i> , 2017 , 12, e0168549	3.7	23
223	Low-intensity cognitive-behaviour therapy interventions for obsessive-compulsive disorder compared to waiting list for therapist-led cognitive-behaviour therapy: 3-arm randomised controlled trial of clinical effectiveness. <i>PLoS Medicine</i> , 2017 , 14, e1002337	11.6	14
222	The ethics of II rials within Cohorts [(TwiCs): 2nd international symposium. <i>Trials</i> , 2017 , 18,	2.8	5
221	A randomised controlled trial assessing the severity and duration of depressive symptoms associated with a clinically significant response to sertraline versus placebo, in people presenting to primary care with depression (PANDA trial): study protocol for a randomised controlled trial.	2.8	14
220	Trials, 2017 , 18, 496 Smoking cessation in severe mental ill health: what works? an updated systematic review and meta-analysis. <i>BMC Psychiatry</i> , 2017 , 17, 252	4.2	64
219	Identifying primary care quality indicators for people with serious mental illness: a systematic review. <i>British Journal of General Practice</i> , 2017 , 67, e519-e530	1.6	14
218	An randomized controlled trial of Post-it notes did not increase postal response rates in older depressed participants. <i>Journal of Evaluation in Clinical Practice</i> , 2017 , 23, 102-107	2.5	4

217	Smoking Cessation Intervention for Severe Mental Ill Health Trial (SCIMITAR+): study protocol for a randomised controlled trial. <i>Trials</i> , 2017 , 18, 44	2.8	12
216	Informing efficient randomised controlled trials: exploration of challenges in developing progression criteria for internal pilot studies. <i>BMJ Open</i> , 2017 , 7, e013537	3	122
215	CollAborative care and active surveillance for Screen-Positive EldeRs with subthreshold depression (CASPER): a multicentred randomised controlled trial of clinical effectiveness and cost-effectiveness. <i>Health Technology Assessment</i> , 2017 , 21, 1-196	4.4	7
214	Clinical effectiveness, cost-effectiveness and acceptability of low-intensity interventions in the management of obsessive-compulsive disorder: the Obsessive-Compulsive Treatment Efficacy randomised controlled Trial (OCTET). <i>Health Technology Assessment</i> , 2017 , 21, 1-132	4.4	27
213	Cost and Outcome of BehaviouRal Activation (COBRA): a randomised controlled trial of behavioural activation versus cognitive-behavioural therapy for depression. <i>Health Technology Assessment</i> , 2017 , 21, 1-366	4.4	13
212	CollAborative care for Screen-Positive EldeRs with major depression (CASPER plus): a multicentred randomised controlled trial of clinical effectiveness and cost-effectiveness. <i>Health Technology Assessment</i> , 2017 , 21, 1-252	4.4	36
211	Maternal psychological distress in primary care and association with child behavioural outcomes at age three. <i>European Child and Adolescent Psychiatry</i> , 2016 , 25, 601-13	5.5	12
2 10	Prevalence of HIV, hepatitis B, and hepatitis C in people with severe mental illness: a systematic review and meta-analysis. <i>Lancet Psychiatry,the</i> , 2016 , 3, 40-48	23.3	127
209	Association Between Chronic Physical Conditions and the Effectiveness of Collaborative Care for Depression: An Individual Participant Data Meta-analysis. <i>JAMA Psychiatry</i> , 2016 , 73, 978-89	14.5	54
208	On poverty, politics and psychology: the socioeconomic gradient of mental healthcare utilisation and outcomes. <i>British Journal of Psychiatry</i> , 2016 , 209, 429-430	5.4	42
207	Variation and ethnic inequalities in treatment of common mental disorders before, during and after pregnancy: combined analysis of routine and research data in the Born in Bradford cohort. <i>BMC Psychiatry</i> , 2016 , 16, 99	4.2	17
206	Mirtazapine added to selective serotonin reuptake inhibitors for treatment-resistant depression in primary care (MIR trial): study protocol for a randomised controlled trial. <i>Trials</i> , 2016 , 17, 66	2.8	9
205	Exploring why people with SMI smoke and why they may want to quit: baseline data from the SCIMITAR RCT. <i>Journal of Psychiatric and Mental Health Nursing</i> , 2016 , 23, 282-9	2.4	10
204	Usefulness of PHQ-9 in primary care to determine meaningful symptoms of low mood: a qualitative study. <i>British Journal of General Practice</i> , 2016 , 66, e78-84	1.6	27
203	Clinical effectiveness and cost-effectiveness of collaborative care for depression in UK primary care (CADET): a cluster randomised controlled trial. <i>Health Technology Assessment</i> , 2016 , 20, 1-192	4.4	30
202	The second Randomised Evaluation of the Effectiveness, cost-effectiveness and Acceptability of Computerised Therapy (REEACT-2) trial: does the provision of telephone support enhance the effectiveness of computer-delivered cognitive behaviour therapy? A randomised controlled trial.	4.4	20
201	The Management of Depression and Anxiety in Older People: Evidence-Based Psychological Interventions, Stepped Care and Collaborative Care 2016 , 79-92		1
200	Making the journey with me: a qualitative study of experiences of a bespoke mental health smoking cessation intervention for service users with serious mental illness. <i>BMC Psychiatry</i> , 2016 , 16, 193	4.2	13

199	Identifying depression with the PHQ-2: A diagnostic meta-analysis. <i>Journal of Affective Disorders</i> , 2016 , 203, 382-395	6.6	83
198	The diagnostic accuracy of brief versions of the Geriatric Depression Scale: a systematic review and meta-analysis. <i>International Journal of Geriatric Psychiatry</i> , 2016 , 31, 837-57	3.9	111
197	Identification of depression in women during pregnancy and the early postnatal period using the Whooley questions and the Edinburgh Postnatal Depression Scale: protocol for the Born and Bred in Yorkshire: PeriNatal Depression Diagnostic Accuracy (BaBY PaNDA) study. <i>BMJ Open</i> , 2016 , 6, e0112	3 223	11
196	Varenicline for smoking cessation and reduction in people with severe mental illnesses: systematic review and meta-analysis. <i>Addiction</i> , 2016 , 111, 1554-67	4.6	26
195	Loneliness and social isolation as risk factors for coronary heart disease and stroke: systematic review and meta-analysis of longitudinal observational studies. <i>Heart</i> , 2016 , 102, 1009-16	5.1	650
194	Loneliness, social isolation and social relationships: what are we measuring? A novel framework for classifying and comparing tools. <i>BMJ Open</i> , 2016 , 6, e010799	3	122
193	Disentangling psychopathology, substance use and dependence: a factor analysis. <i>BMC Psychiatry</i> , 2016 , 16, 281	4.2	4
192	Diagnostic accuracy of the Depression subscale of the Hospital Anxiety and Depression Scale (HADS-D) for detecting major depression: protocol for a systematic review and individual patient data meta-analyses. <i>BMJ Open</i> , 2016 , 6, e011913	3	19
191	Infection in people with severe mental illness - Authors' reply. Lancet Psychiatry, the, 2016, 3, 204-5	23.3	
190	Evaluation of ethnic disparities in detection of depression and anxiety in primary care during the maternal period: combined analysis of routine and cohort data. <i>British Journal of Psychiatry</i> , 2016 , 208, 453-61	5.4	30
189	Life adversity in depressed and non-depressed older adults: A cross-sectional comparison of the brief LTE-Q questionnaire and life events and difficulties interview as part of the CASPER study. Journal of Affective Disorders, 2016, 193, 31-8	6.6	12
188	Routine use of patient reported outcome measures (PROMs) for improving treatment of common mental health disorders in adults. <i>The Cochrane Library</i> , 2016 , 7, CD011119	5.2	83
187	Cost and Outcome of Behavioural Activation versus Cognitive Behavioural Therapy for Depression (COBRA): a randomised, controlled, non-inferiority trial. <i>Lancet, The</i> , 2016 , 388, 871-80	40	282
186	Diagnostic accuracy of the Whooley depression tool in older adults in UK primary care. <i>Journal of Affective Disorders</i> , 2015 , 182, 39-43	6.6	7
185	Feasibility randomized controlled trial of cognitive and behavioral interventions for depression symptoms in patients accessing drug and alcohol treatment. <i>Journal of Substance Abuse Treatment</i> , 2015 , 55, 6-14	4.2	22
184	Bespoke smoking cessation for people with severe mental ill health (SCIMITAR): a pilot randomised controlled trial. <i>Lancet Psychiatry,the</i> , 2015 , 2, 395-402	23.3	59
183	The effectiveness of self-management support interventions for men with long-term conditions: a systematic review and meta-analysis. <i>BMJ Open</i> , 2015 , 5, e006620	3	25
182	Diagnostic accuracy of the Whooley questions for the identification of depression: a diagnostic meta-analysis. <i>BMJ Open</i> , 2015 , 5, e008913	3	47

181	Determinants of hospital length of stay for people with serious mental illness in England and implications for payment systems: a regression analysis. <i>BMC Health Services Research</i> , 2015 , 15, 439	2.9	29
180	Revealing hidden depression in older people: a qualitative study within a randomised controlled trial. <i>BMC Family Practice</i> , 2015 , 16, 142	2.6	17
179	Does the quality and outcomes framework reduce psychiatric admissions in people with serious mental illness? A regression analysis. <i>BMJ Open</i> , 2015 , 5, e007342	3	14
178	M9 Lung Health of OPiate Users (LHOp): A pilot study to assess the respiratory health of opiate misusers attending a community substance misuse clinic. <i>Thorax</i> , 2015 , 70, A230.2-A231	7.3	
177	Post-it notes to improve questionnaire response rates in RCTs - findings from a randomised sub-study. <i>Trials</i> , 2015 , 16,	2.8	78
176	The acceptability of case-finding questions to identify perinatal depression. <i>British Journal of Midwifery</i> , 2015 , 23, 630-638	0.3	4
175	Evidence-based guidelines for treating depressive disorders with antidepressants: A revision of the 2008 British Association for Psychopharmacology guidelines. <i>Journal of Psychopharmacology</i> , 2015 , 29, 459-525	4.6	381
174	Systematic review and meta-analysis of transdiagnostic psychological treatments for anxiety and depressive disorders in adulthood. <i>Clinical Psychology Review</i> , 2015 , 40, 91-110	10.8	297
173	Patient experience of computerised therapy for depression in primary care. <i>BMJ Open</i> , 2015 , 5, e00858	313	43
172	Computerised cognitive behaviour therapy (cCBT) as treatment for depression in primary care (REEACT trial): large scale pragmatic randomised controlled trial. <i>BMJ, The</i> , 2015 , 351, h5627	5.9	235
171	Screening and case finding for major depressive disorder using the Patient Health Questionnaire (PHQ-9): a meta-analysis. <i>General Hospital Psychiatry</i> , 2015 , 37, 567-76	5.6	156
170	Diagnostic accuracy of the Edinburgh Postnatal Depression Scale (EPDS) for detecting major depression in pregnant and postnatal women: protocol for a systematic review and individual patient data meta-analyses. <i>BMJ Open</i> , 2015 , 5, e009742	3	35
169	A diagnostic meta-analysis of the Patient Health Questionnaire-9 (PHQ-9) algorithm scoring method as a screen for depression. <i>General Hospital Psychiatry</i> , 2015 , 37, 67-75	5.6	311
168	Do higher primary care practice performance scores predict lower rates of emergency admissions for persons with serious mental illness? An analysis of secondary panel data. <i>Health Services and Delivery Research</i> , 2015 , 3, 1-68	1.5	4
167	A systematic review and metaethnography to identify how effective, cost-effective, accessible and acceptable self-management support interventions are for men with long-term conditions (SELF-MAN). <i>Health Services and Delivery Research</i> , 2015 , 3, 1-302	1.5	5
166	Screening for psychological and mental health difficulties in young people who offend: a systematic review and decision model. <i>Health Technology Assessment</i> , 2015 , 19, 1-128	4.4	5
165	A randomised controlled trial of computerised cognitive behaviour therapy for the treatment of depression in primary care: the Randomised Evaluation of the Effectiveness and Acceptability of Computerised Therapy (REEACT) trial. <i>Health Technology Assessment</i> , 2015 , 19, viii, xxi-171	4.4	40
164	Smoking Cessation Intervention for severe Mental Ill Health Trial (SCIMITAR): a pilot randomised control trial of the clinical effectiveness and cost-effectiveness of a bespoke smoking cessation service. Health Technology Assessment 2015, 19, 1-148, v-vi	4.4	22

163	Clinical effectiveness and cost-effectiveness of parenting interventions for children with severe attachment problems: a systematic review and meta-analysis. <i>Health Technology Assessment</i> , 2015 , 19, vii-xxviii, 1-347	4.4	22
162	Depression screening and patient outcomes in pregnancy or postpartum: a systematic review. Journal of Psychosomatic Research, 2014 , 76, 433-46	4.1	69
161	Cost and outcome of behavioural activation versus cognitive behaviour therapy for depression (COBRA): study protocol for a randomised controlled trial. <i>Trials</i> , 2014 , 15, 29	2.8	18
160	Early changes, attrition, and dose-response in low intensity psychological interventions. <i>British Journal of Clinical Psychology</i> , 2014 , 53, 114-30	3.6	61
159	Computerised Cognitive Behaviour Therapy for Depression Management: A Cost-Effectiveness Analysis. <i>Value in Health</i> , 2014 , 17, A459	3.3	
158	Sudden gains in behavioural activation for depression. <i>Behaviour Research and Therapy</i> , 2014 , 60, 34-8	5.2	18
157	Obsessive Compulsive Treatment Efficacy Trial (OCTET) comparing the clinical and cost effectiveness of self-managed therapies: study protocol for a randomised controlled trial. <i>Trials</i> , 2014 , 15, 278	2.8	9
156	Heterogeneity in patient-reported outcomes following low-intensity mental health interventions: a multilevel analysis. <i>PLoS ONE</i> , 2014 , 9, e99658	3.7	15
155	Characteristics of effective collaborative care for treatment of depression: a systematic review and meta-regression of 74 randomised controlled trials. <i>PLoS ONE</i> , 2014 , 9, e108114	3.7	115
154	Routine use of patient reported outcome measures (PROMs) for improving treatment of common mental health disorders in adults 2014 ,		3
153	The accessibility and acceptability of self-management support interventions for men with long term conditions: a systematic review and meta-synthesis of qualitative studies. <i>BMC Public Health</i> , 2014 , 14, 1230	4.1	45
152	The diagnostic accuracy of the Patient Health Questionnaire-2 (PHQ-2), Patient Health Questionnaire-8 (PHQ-8), and Patient Health Questionnaire-9 (PHQ-9) for detecting major depression: protocol for a systematic review and individual patient data meta-analyses. <i>Systematic</i>	3	53
151	Computerised cognitive behaviour therapy for depression in adolescents: study protocol for a feasibility randomised controlled trial. <i>BMJ Open</i> , 2014 , 4, e006488	3	13
150	Promoting HIV and sexual safety behaviour in people with severe mental illness: a systematic review of behavioural interventions. <i>International Journal of Mental Health Nursing</i> , 2014 , 23, 344-54	3.8	14
149	CASPER plus (CollAborative care in Screen-Positive EldeRs with major depressive disorder): study protocol for a randomised controlled trial. <i>Trials</i> , 2014 , 15, 451	2.8	12
148	Benchmarking routine psychological services: a discussion of challenges and methods. <i>Behavioural and Cognitive Psychotherapy</i> , 2014 , 42, 16-30	2.1	16
147	Adapting manualized Behavioural Activation treatment for older adults with depression. <i>The Cognitive Behaviour Therapist</i> , 2014 , 7,	2.9	15
146	Cost-effectiveness of combining systematic identification and treatment of co-morbid major depression for people with chronic diseases: the example of cancer. <i>Psychological Medicine</i> , 2014 , 44, 1451-60	6.9	26

145	Qualitative meta-synthesis of user experience of computerised therapy for depression and anxiety. <i>PLoS ONE</i> , 2014 , 9, e84323	3.7	107
144	Behavioural activation for depression; an update of meta-analysis of effectiveness and sub group analysis. <i>PLoS ONE</i> , 2014 , 9, e100100	3.7	259
143	Cost-effectiveness of collaborative care for depression in UK primary care: economic evaluation of a randomised controlled trial (CADET). <i>PLoS ONE</i> , 2014 , 9, e104225	3.7	33
142	A systematic review, psychometric analysis and qualitative assessment of generic preference-based measures of health in mental health populations and the estimation of mapping functions from widely used specific measures. <i>Health Technology Assessment</i> , 2014 , 18, vii-viii, xiii-xxv, 1-188	4.4	141
141	Assessing the risk of self-harm in an adult offender population: an incidence cohort study. <i>Health Technology Assessment</i> , 2014 , 18, 1-151, vii-viii	4.4	9
140	The psychometric properties of the subscales of the GHQ-28 in a multi-ethnic maternal sample: results from the Born in Bradford cohort. <i>BMC Psychiatry</i> , 2013 , 13, 55	4.2	18
139	Nurse-delivered collaborative care for depression and long-term physical conditions: a systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2013 , 149, 14-22	6.6	69
138	Psychometric properties of the Treatment Outcomes Profile (TOP) psychological health scale. <i>Mental Health and Substance Use: Dual Diagnosis</i> , 2013 , 6, 140-149		4
137	Decision curve analysis for assessing the usefulness of tests for making decisions to treat: an application to tests for prodromal psychosis. <i>Psychological Assessment</i> , 2013 , 25, 730-7	5.3	5
136	Clinical effectiveness of collaborative care for depression in UK primary care (CADET): cluster randomised controlled trial. <i>BMJ, The</i> , 2013 , 347, f4913	5.9	136
135	Influence of initial severity of depression on effectiveness of low intensity interventions: meta-analysis of individual patient data. <i>BMJ, The</i> , 2013 , 346, f540	5.9	182
134	Depression, anxiety and comorbid substance use: association patterns in outpatient addictions treatment. <i>Mental Health and Substance Use: Dual Diagnosis</i> , 2013 , 6, 59-75		6
133	The CORE-10: A short measure of psychological distress for routine use in the psychological therapies. <i>Counselling and Psychotherapy Research</i> , 2013 , 13, 3-13	1.3	125
132	Psychological distress during pregnancy in a multi-ethnic community: findings from the born in Bradford cohort study. <i>PLoS ONE</i> , 2013 , 8, e60693	3.7	30
131	Should methodological filters for diagnostic test accuracy studies be used in systematic reviews of psychometric instruments? A case study involving screening for postnatal depression. <i>Systematic Reviews</i> , 2012 , 1, 9	3	8
130	Brief case finding tools for anxiety disorders: validation of GAD-7 and GAD-2 in addictions treatment. <i>Drug and Alcohol Dependence</i> , 2012 , 125, 37-42	4.9	50
129	Delivering stepped care: an analysis of implementation in routine practice. <i>Implementation Science</i> , 2012 , 7, 3	8.4	90
128	Acceptability of mental health screening in routine addictions treatment. <i>General Hospital Psychiatry</i> , 2012 , 34, 415-22	5.6	7

127	Rethinking recommendations for screening for depression in primary care. <i>Cmaj</i> , 2012 , 184, 413-8	3.5	142
126	Diagnostic accuracy of case-finding questions to identify perinatal depression. <i>Cmaj</i> , 2012 , 184, E424-3	03.5	25
125	Collaborative care for depression and anxiety problems. <i>The Cochrane Library</i> , 2012 , 10, CD006525	5.2	451
124	Re-rethinking the article by Thombs and colleagues: Figure 1:. <i>Cmaj</i> , 2012 , 184, 438.2-439	3.5	2
123	Optimal cut-off score for diagnosing depression with the Patient Health Questionnaire (PHQ-9): a meta-analysis. <i>Cmaj</i> , 2012 , 184, E191-6	3.5	892
122	How reliable is depression screening in alcohol and drug users? A validation of brief and ultra-brief questionnaires. <i>Journal of Affective Disorders</i> , 2011 , 134, 266-71	6.6	41
121	P1-294 Risk factors for poor mental health during pregnancy from the born in Bradford cohort: early results from a bi-ethnic sample. <i>Journal of Epidemiology and Community Health</i> , 2011 , 65, A148-A	14 ⁵ 8 ¹	
120	P1-496 An examination of the psychometric properties of the GHQ-28 subscales in a bi-ethnic antenatal sample in the UK: preliminary results from the born in Bradford cohort. <i>Journal of Epidemiology and Community Health</i> , 2011 , 65, A204-A204	5.1	
119	Behavioural activation delivered by the non-specialist: phase II randomised controlled trial. <i>British Journal of Psychiatry</i> , 2011 , 198, 66-72	5.4	141
118	Cost utility of behavioural activation delivered by the non-specialist. <i>British Journal of Psychiatry</i> , 2011 , 199, 510-1	5.4	30
117	Screening and case finding for depression in offender populations: a systematic review of diagnostic properties. <i>Journal of Affective Disorders</i> , 2011 , 128, 72-82	6.6	12
116	Validity of two case finding questions to detect postnatal depression: a review of diagnostic test accuracy. <i>Journal of Affective Disorders</i> , 2011 , 133, 388-97	6.6	15
115	A randomised evaluation of CollAborative care and active surveillance for Screen-Positive EldeRs with sub-threshold depression (CASPER): study protocol for a randomized controlled trial. <i>Trials</i> , 2011 , 12, 225	2.8	26
114	Can we improve recruitment to trials and informed consent by improving participant information sheets? - A nested RCT. <i>Trials</i> , 2011 , 12,	2.8	2
113	Recruiting ahead of target: What worked in the REEACT trial?. <i>Trials</i> , 2011 , 12,	2.8	2
112	The effectiveness of behavioural therapy for the treatment of depression in older adults: a meta-analysis. <i>International Journal of Geriatric Psychiatry</i> , 2011 , 26, 1211-20	3.9	35
111	Smoking cessation in severe mental illness: what works?. <i>Addiction</i> , 2010 , 105, 1176-89	4.6	174
110	Effectiveness of interventions designed to promote patient involvement to enhance safety: a systematic review. <i>BMJ Quality and Safety</i> , 2010 , 19, e10	5.4	51

(2009-2010)

109	validated, in what setting and in which language?. <i>International Journal of Psychiatry in Clinical Practice</i> , 2010 , 14, 72-6	2.4	44	
108	Screening for postnatal depression lis it acceptable to women and healthcare professionals? A systematic review and meta-synthesis. <i>Journal of Reproductive and Infant Psychology</i> , 2010 , 28, 328-344	2.9	54	
107	The Hospital Anxiety and Depression Scale: a diagnostic meta-analysis of case-finding ability. Journal of Psychosomatic Research, 2010 , 69, 371-8	4.1	223	
106	Prevalence and incidence of postnatal depression: what can systematic reviews tell us?. <i>Archives of Womenis Mental Health</i> , 2010 , 13, 295-305	5	38	
105	Improving health and productivity of depressed workers: a pilot randomized controlled trial of telephone cognitive behavioral therapy delivery in workplace settings. <i>General Hospital Psychiatry</i> , 2010 , 32, 337-40	5.6	24	
104	Defining successful treatment outcome in depression using the PHQ-9: a comparison of methods. <i>Journal of Affective Disorders</i> , 2010 , 127, 122-9	6.6	127	
103	What work has to be done to implement collaborative care for depression? Process evaluation of a trial utilizing the Normalization Process Model. <i>Implementation Science</i> , 2010 , 5, 15	8.4	44	
102	Translating research into practice in Leeds and Bradford (TRiPLaB): a protocol for a programme of research. <i>Implementation Science</i> , 2010 , 5, 37	8.4	18	
101	User-defined outcomes in mental health: A qualitative study and consensus development exercise. <i>Journal of Mental Health</i> , 2009 , 18, 415-423	2.7	5	
100	A stepped care intervention is effective for elderly people with subthreshold depression or anxiety. <i>Evidence-Based Mental Health</i> , 2009 , 12, 115	11.1		
99	Collaborative Depression Trial (CADET): multi-centre randomised controlled trial of collaborative care for depressionstudy protocol. <i>BMC Health Services Research</i> , 2009 , 9, 188	2.9	27	
98	Assessing the quality of diagnostic studies using psychometric instruments: applying QUADAS. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2009 , 44, 300-7	4.5	22	
97	Putting the 'Q' in depression QALYs: a comparison of utility measurement using EQ-5D and SF-6D health related quality of life measures. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2009 , 44, 569-78	4.5	28	
96	Detecting and predicting self-harm behaviour in prisoners: a prospective psychometric analysis of three instruments. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2009 , 44, 853-61	4.5	20	
95	Is it clinically and cost effective to screen for postnatal depression: a systematic review of controlled clinical trials and economic evidence. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2009 , 116, 1019-27	3.7	46	
94	Barriers to the uptake of computerized cognitive behavioural therapy: a systematic review of the quantitative and qualitative evidence. <i>Psychological Medicine</i> , 2009 , 39, 705-12	6.9	267	
93	Collaborative care for depression in UK primary care: a randomized controlled trial ©orrigendum. <i>Psychological Medicine</i> , 2009 , 39, 701-701	6.9	2	
92	Screening for postnatal depression in primary care: cost effectiveness analysis. <i>BMJ, The</i> , 2009 , 339, b5	2 9 3)	87	

91	Methods to identify postnatal depression in primary care: an integrated evidence synthesis and value of information analysis. <i>Health Technology Assessment</i> , 2009 , 13, 1-145, 147-230	4.4	139
90	Cluster randomized trials produced similar results to individually randomized trials in a meta-analysis of enhanced care for depression. <i>Journal of Clinical Epidemiology</i> , 2008 , 61, 160-168	5.7	22
89	Telephone treatment support improves outcomes for depressed employees. <i>Evidence-Based Mental Health</i> , 2008 , 11, 47	11.1	
88	Screening and case-finding instruments for depression: a meta-analysis. <i>Cmaj</i> , 2008 , 178, 997-1003	3.5	312
87	A meta-analysis of randomized trials of behavioural treatment of depression. <i>Psychological Medicine</i> , 2008 , 38, 611-23	6.9	279
86	Depression in older adults: collaborative care model seems effective. <i>Evidence-Based Mental Health</i> , 2008 , 11, 44	11.1	3
85	The European pharmaceutical industry: coming to a screen near you?. <i>Journal of the Royal Society of Medicine</i> , 2008 , 101, 485-8	2.3	1
84	Collaborative care for depression in UK primary care: a randomized controlled trial. <i>Psychological Medicine</i> , 2008 , 38, 279-87	6.9	120
83	Psychotherapy mediated by remote communication technologies: a meta-analytic review. <i>BMC Psychiatry</i> , 2008 , 8, 60	4.2	117
82	A smoking ban in psychiatric units: threat or opportunity?. <i>Mental Health in Family Medicine</i> , 2008 , 5, 129-33		
81	A review and critical appraisal of measures of therapist-patient interactions in mental health settings. <i>Health Technology Assessment</i> , 2008 , 12, iii, ix-47	4.4	31
80	Screening for depression in medical settings with the Patient Health Questionnaire (PHQ): a diagnostic meta-analysis. <i>Journal of General Internal Medicine</i> , 2007 , 22, 1596-602	4	767
79	Methylenetetrahydrofolate reductase (MTHFR) genetic polymorphisms and psychiatric disorders: a HuGE review. <i>American Journal of Epidemiology</i> , 2007 , 165, 1-13	3.8	247
78	Can we predict suicide and non-fatal self-harm with the Beck Hopelessness Scale? A meta-analysis. <i>Psychological Medicine</i> , 2007 , 37, 769-78	6.9	172
77	Is low folate a risk factor for depression? A meta-analysis and exploration of heterogeneity. <i>Journal of Epidemiology and Community Health</i> , 2007 , 61, 631-7	5.1	162
76	IMPACT collaborative care programme reduces suicide ideation in depressed older adults. <i>Evidence-Based Mental Health</i> , 2007 , 10, 51	11.1	5
75	What makes self-help interventions effective in the management of depressive symptoms? Meta-analysis and meta-regression. <i>Psychological Medicine</i> , 2007 , 37, 1217-28	6.9	293
74	Vitamin K and the Prevention of Fractures Reply. Archives of Internal Medicine, 2007, 167, 94		1

73	Collaborative care for depression and anxiety problems 2007,		6
72	Randomized trials with concurrent economic evaluations reported unrepresentatively large clinical effect sizes. <i>Journal of Clinical Epidemiology</i> , 2007 , 60, 781-6	5.7	10
71	Diagnosing depression in primary care using self-completed instruments: UK validation of PHQ-9 and CORE-OM. <i>British Journal of General Practice</i> , 2007 , 57, 650-2	1.6	125
70	A systematic review of quantitative and qualitative research on the role and effectiveness of written information available to patients about individual medicines. <i>Health Technology Assessment</i> , 2007, 11, iii, 1-160	4.4	176
69	A systematic review and economic model of the clinical effectiveness and cost-effectiveness of interventions for preventing relapse in people with bipolar disorder. <i>Health Technology Assessment</i> , 2007 , 11, iii-iv, ix-206	4.4	70
68	Developing a U.K. protocol for collaborative care: a qualitative study. <i>General Hospital Psychiatry</i> , 2006 , 28, 296-305	5.6	54
67	Should we screen for depression?. <i>BMJ, The</i> , 2006 , 332, 1027-30	5.9	138
66	Collaborative care for depression: a cumulative meta-analysis and review of longer-term outcomes. <i>Archives of Internal Medicine</i> , 2006 , 166, 2314-21		913
65	Vitamin K and the prevention of fractures: systematic review and meta-analysis of randomized controlled trials. <i>Archives of Internal Medicine</i> , 2006 , 166, 1256-61		262
64	Costs and consequences of enhanced primary care for depression: systematic review of randomised economic evaluations. <i>British Journal of Psychiatry</i> , 2006 , 189, 297-308	5.4	134
63	Can we improve the morale of staff working in psychiatric units? A systematic review. <i>Journal of Mental Health</i> , 2006 , 15, 7-17	2.7	43
62	Collaborative care for depression in primary care. Making sense of a complex intervention: systematic review and meta-regression. <i>British Journal of Psychiatry</i> , 2006 , 189, 484-93	5.4	293
61	The prevalence of nursing staff stress on adult acute psychiatric in-patient wards. A systematic review. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2006 , 41, 34-43	4.5	32
60	Screening and case finding instruments for depression. <i>The Cochrane Library</i> , 2005 , CD002792	5.2	102
59	Benefits and harms of direct to consumer advertising: a systematic review. <i>Quality and Safety in Health Care</i> , 2005 , 14, 246-50		138
58	Managing common mental health disorders in primary care: conceptual models and evidence base. <i>BMJ, The</i> , 2005 , 330, 839-42	5.9	143
57	Comparison of large versus smaller randomized trials for mental health-related interventions. <i>American Journal of Psychiatry</i> , 2005 , 162, 578-84	11.9	28
56	Stepped care in psychological therapies: access, effectiveness and efficiency. Narrative literature review. <i>British Journal of Psychiatry</i> , 2005 , 186, 11-7	5.4	609

55	Review: disease management programmes improve detection and care of people with depression. <i>Evidence-Based Mental Health</i> , 2004 , 7, 80	11.1	9
54	Educational and organizational interventions to improve the management of depression in primary care: a systematic review. <i>JAMA - Journal of the American Medical Association</i> , 2003 , 289, 3145-51	27.4	501
53	Outcome measures and needs assessment tools for schizophrenia and related disorders. <i>The Cochrane Library</i> , 2003 , CD003081	5.2	23
52	Improving the detection and management of depression in primary care. <i>Quality and Safety in Health Care</i> , 2003 , 12, 149-55		67
51	Improving the delivery and organisation of mental health services: beyond the conventional randomised controlled trial. <i>British Journal of Psychiatry</i> , 2002 , 180, 13-8	5.4	43
50	Randomized controlled trials in schizophrenia: a critical perspective on the literature. <i>Acta Psychiatrica Scandinavica</i> , 2002 , 105, 243-51	6.5	38
49	4 factors were predictive of extended duration in hospital in acute psychiatric admission. <i>Evidence-Based Mental Health</i> , 2002 , 5, 64	11.1	1
48	Routine administration of Health Related Quality of Life (HRQoL) and needs assessment instruments to improve psychological outcomea systematic review. <i>Psychological Medicine</i> , 2002 , 32, 1345-56	6.9	77
47	Outcomes research in mental health. Systematic review. British Journal of Psychiatry, 2002, 181, 8-16	5.4	74
46	Newer atypical antipsychotic medication in comparison to clozapine: a systematic review of randomized trials. <i>Schizophrenia Research</i> , 2002 , 56, 1-10	3.6	81
45	Routinely administered questionnaires for depression and anxiety: systematic review. <i>BMJ: British Medical Journal</i> , 2001 , 322, 406-9		227
44	Nobody ever expects the Spanish Inquisition (Python, 1991). <i>Psychiatric Bulletin</i> , 2001 , 25, 291-292		6
43	Risperidone versus other atypical antipsychotic medication for schizophrenia. <i>Cochrane Database of Systematic Reviews</i> , 2000 , CD002306		32
42	The causes, consequences and detection of publication bias in psychiatry. <i>Acta Psychiatrica Scandinavica</i> , 2000 , 102, 241-9	6.5	66
41	Influence of methodology on outcomes of randomised clozapine trials. <i>Pharmacopsychiatry</i> , 2000 , 33, 54-9	2	10
40	Drug treatments for schizophrenia. <i>Quality in Health Care: QHC</i> , 2000 , 9, 73-9		4
39	Efficacy of beta-blocker supplementation for schizophrenia: a systematic review of randomized trials. <i>Schizophrenia Research</i> , 2000 , 41, 341-7	3.6	12
38	Modelling publication bias in meta-analysis: a review. <i>Statistical Methods in Medical Research</i> , 2000 , 9, 421-445	2.3	63

37	Publication and related biases Health Technology Assessment, 2000, 4,	4.4	261
36	Lost information? The fate of papers presented at the 40th society for Social Medicine Conference. <i>Journal of Epidemiology and Community Health</i> , 1999 , 53, 442-3	5.1	11
35	The role of electronic journals in reducing publication bias. <i>Informatics for Health and Social Care</i> , 1999 , 24, 223-9		12
34	Vegetarianism in young women: another means of weight control?. <i>International Journal of Eating Disorders</i> , 1999 , 26, 87-90	6.3	42
33	Relation between hostility and coronary heart disease. Evidence does not support link. <i>BMJ: British Medical Journal</i> , 1999 , 319, 917-8		17
32	NICE work?: Potential and problems for psychiatry in the new National Health Service. <i>Psychiatric Bulletin</i> , 1999 , 23, 513-516		2
31	Mental health promotion in high risk groups. Quality and Safety in Health Care, 1997, 6, 219-25		4
30	Rational decision-making in psychiatry: evidence-based psychiatry is just the start. <i>British Journal of Psychiatry</i> , 1997 , 171, 226-7	5.4	2
29	Prescribing antidepressants in general practice 1997 , 314, 826-826		6
28	'Number needed to treat'. British Journal of Psychiatry, 1996 , 169, 113	5.4	1
27	Deliberate self harm. <i>Emergency Medicine Journal</i> , 1996 , 13, 432	1.5	1
26	Differences in use of abbreviated mental test score by geriatricians and psychiatrists. <i>BMJ: British Medical Journal</i> , 1996 , 313, 465		13
25	Evidence-based medicine: An improved format for journal clubs. <i>Psychiatric Bulletin</i> , 1996 , 20, 673-675		18
24	Publication bias and meta-analysis. <i>British Journal of Psychiatry</i> , 1995 , 167, 266	5.4	5
23	Misleading meta-analysis. Subject to many potential biases. <i>BMJ: British Medical Journal</i> , 1995 , 311, 130	3-4	3
22	Linking evidence to practice31-54		
21	Planning and Conducting Systematic Reviews150-179		3
20	Rapid systematic review of systematic reviews: what befriending, social support and low intensity psychosocial interventions, delivered remotely, are effective in reducing social isolation and loneliness among older adults? How do they work?. <i>F1000Research</i> ,9, 1368	3.6	2

19	What factors indicate prognosis for adults with depression in primary care? A protocol for meta-analyses of individual patient data using the Dep-GP database. <i>Wellcome Open Research</i> ,4, 69	4.8	1
18	Depression in primary care1-16		1
17	United Kingdom perspective127-136		1
16	Rapid systematic review of systematic reviews: what befriending, social support and low intensity psychosocial interventions, delivered remotely, may reduce social isolation and loneliness among older adults and how?. <i>F1000Research</i> ,9, 1368	3.6	5
15	Home hazard assessment and environmental modification to prevent falls in older people: the OTIS trial. <i>F1000Research</i> ,10, 500	3.6	2
14	Use of the Internet and digital devices among people with severe mental ill health during the COVID-19 pandemic restrictions		1
13	Improving Access to Care386-400		
12	Collaborative Care and Stepped Care401-419		
11	Loneliness among people with severe mental ill health during the COVID-19 pandemic: results from a linked UK population cohort study		1
10	Pharmacy-based management for depression in adults. <i>The Cochrane Library</i> ,	5.2	3
9	The Current View of Evidence and Evidence-Based Practice1-20		9
8	Depressive Disorders in Primary Care: A Review271-318		1
7	Models of care for depression17-30		
6	Anatomy of a review55-72		
5	Collaborative care87-102		
4	Referral103-110		
3	Summary of the evidence111-116		
2	Making it happen117-126		

LIST OF PUBLICATIONS

1 United States perspective 137-144