

Zhijian Huang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6339088/publications.pdf>

Version: 2024-02-01

16
papers

590
citations

687220

13
h-index

940416

16
g-index

16
all docs

16
docs citations

16
times ranked

781
citing authors

#	ARTICLE	IF	CITATIONS
1	The long non-coding RNA Neat1 is an important mediator of the therapeutic effect of bexarotene on traumatic brain injury in mice. <i>Brain, Behavior, and Immunity</i> , 2017, 65, 183-194.	2.0	86
2	Altered expression of long non-coding RNA and mRNA in mouse cortex after traumatic brain injury. <i>Brain Research</i> , 2016, 1646, 589-600.	1.1	73
3	Apolipoprotein E Deficiency Aggravates Neuronal Injury by Enhancing Neuroinflammation via the JNK/c-Jun Pathway in the Early Phase of Experimental Subarachnoid Hemorrhage in Mice. <i>Oxidative Medicine and Cellular Longevity</i> , 2019, 2019, 1-15.	1.9	57
4	Fecal Microbiota Transplantation Is a Promising Method to Restore Gut Microbiota Dysbiosis and Relieve Neurological Deficits after Traumatic Brain Injury. <i>Oxidative Medicine and Cellular Longevity</i> , 2021, 2021, 1-21.	1.9	54
5	ApoE Influences the Blood-Brain Barrier Through the NF- κ B/MMP-9 Pathway After Traumatic Brain Injury. <i>Scientific Reports</i> , 2017, 7, 6649.	1.6	47
6	Bexarotene protects against traumatic brain injury in mice partially through apolipoprotein E. <i>Neuroscience</i> , 2017, 343, 434-448.	1.1	46
7	Bexarotene protects against neurotoxicity partially through a PPAR δ -dependent mechanism in mice following traumatic brain injury. <i>Neurobiology of Disease</i> , 2018, 117, 114-124.	2.1	38
8	Significant changes in circular RNA in the mouse cerebral cortex around an injury site after traumatic brain injury. <i>Experimental Neurology</i> , 2019, 313, 37-48.	2.0	33
9	<p>The association of neutrophil-to-lymphocyte ratio and delayed cerebral ischemia in patients with aneurysmal subarachnoid hemorrhage: possible involvement of cerebral blood perfusion</p>. <i>Neuropsychiatric Disease and Treatment</i> , 2019, Volume 15, 1001-1007.	1.0	30
10	Intraventricular apolipoprotein ApoJ infusion acts protectively in Traumatic Brain Injury. <i>Journal of Neurochemistry</i> , 2016, 136, 1017-1025.	2.1	26
11	Peroxisome Proliferator-Activated Receptor δ Alleviates Early Brain Injury After Subarachnoid Hemorrhage in Rats. <i>Stroke</i> , 2016, 47, 196-205.	1.0	25
12	Effect of APOE Gene Polymorphism on Early Cerebral Perfusion After Aneurysmal Subarachnoid Hemorrhage. <i>Translational Stroke Research</i> , 2015, 6, 446-450.	2.3	24
13	Bexarotene promotes microglia/macrophages - Specific brain - Derived Neurotrophic factor expression and axon sprouting after traumatic brain injury. <i>Experimental Neurology</i> , 2020, 334, 113462.	2.0	16
14	Clinical and Basic Evaluation of the Prognostic Value of Uric Acid in Traumatic Brain Injury. <i>International Journal of Medical Sciences</i> , 2018, 15, 1072-1082.	1.1	15
15	The lncRNA-AK046375 Upregulates Metallothionein-2 by Sequestering miR-491-5p to Relieve the Brain Oxidative Stress Burden after Traumatic Brain Injury. <i>Oxidative Medicine and Cellular Longevity</i> , 2022, 2022, 1-26.	1.9	12
16	Endothelial Regulation by Exogenous Annexin A1 in Inflammatory Response and BBB Integrity Following Traumatic Brain Injury. <i>Frontiers in Neuroscience</i> , 2021, 15, 627110.	1.4	8