

Flávio Augusto Cardoso de Faria

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/634127/publications.pdf>

Version: 2024-02-01

23
papers

480
citations

686830

13
h-index

713013

21
g-index

23
all docs

23
docs citations

23
times ranked

567
citing authors

#	ARTICLE	IF	CITATIONS
1	Pharmacogenetic and Pharmacokinetic Assays from Saliva Samples Can Guarantee Personalized Drug Prescription. Brazilian Dental Journal, 2021, 32, 3-8.	0.5	6
2	Multifocal Analysis of Acute Pain After Third Molar Removal. Frontiers in Pharmacology, 2021, 12, 643874.	1.6	3
3	Simultaneous separation of naproxen and 6-O-desmethylnaproxen metabolite in saliva samples by liquid chromatography-tandem mass spectrometry: Pharmacokinetic study of naproxen alone and associated with esomeprazole. PLoS ONE, 2020, 15, e0236297.	1.1	10
4	Losartan and isoproterenol promote alterations in the local renin-angiotensin system of rat salivary glands. PLoS ONE, 2019, 14, e0217030.	1.1	17
5	Impacto do Tabagismo Passivo na Resposta Pressórica À Epinefrina e Felipressina em Ratos Hipertensos 1K1C Tratados ou Não com Atenolol. Arquivos Brasileiros De Cardiologia, 2019, 114, 295-303.	0.3	0
6	Efficacy of Ketoprofen With or Without Omeprazole for Pain And Inflammation Control After Third Molar Removal. Brazilian Dental Journal, 2018, 29, 140-149.	0.5	10
7	Efficacy of oral diclofenac with or without codeine for pain control after invasive bilateral third molar extractions. International Journal of Oral and Maxillofacial Surgery, 2017, 46, 621-627.	0.7	16
8	Temporomandibular disorders and painful comorbidities: clinical association and underlying mechanisms. Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology, 2017, 123, 288-297.	0.2	51
9	Efficacy of piroxicam for postoperative pain after lower third molar surgery associated with CYP2C8 and CYP2C9. Journal of Pain Research, 2017, Volume 10, 1581-1589.	0.8	19
10	Effective method for the detection of piroxicam in human plasma using HPLC. Brazilian Oral Research, 2016, 30, .	0.6	12
11	Quantification of piroxicam and 5-hydroxy-piroxicam in human plasma and saliva using liquid chromatography-tandem mass spectrometry following oral administration. Journal of Pharmaceutical and Biomedical Analysis, 2016, 120, 212-220.	1.4	21
12	Efficacy of naproxen with or without esomeprazole for pain and inflammation in patients after bilateral third molar extractions: A double blinded crossover study. Medicina Oral, Patologia Oral Y Cirugia Bucal, 2016, 22, 0-0.	0.7	8
13	Functional Local Renin-Angiotensin System in Human and Rat Periodontal Tissue. PLoS ONE, 2015, 10, e0134601.	1.1	47
14	Comparison of epinephrine and felypressin pressure effects in 1K1C hypertensive rats treated or not with atenolol. Journal of Anesthesia, 2015, 29, 56-64.	0.7	5
15	Efficacy and Safety of 2% and 4% Articaine for Lower Third Molar Surgery. Journal of Dental Research, 2015, 94, 166S-173S.	2.5	30
16	Sublingual ketorolac and sublingual piroxicam are equally effective for postoperative pain, trismus, and swelling management in lower third molar removal. Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology, 2012, 114, 27-34.	0.2	17
17	Comparison of oral versus sublingual piroxicam during postoperative pain management after lower third molar extraction. International Journal of Oral and Maxillofacial Surgery, 2011, 40, 292-297.	0.7	18
18	Comparative study between homeopathy and nimesulide on the prevention of bone resorption in experimental periodontitis in rats.. Faculdade De Odontologia De Porto Alegre Revista, 2010, 51, 5-8.	0.1	0

#	ARTICLE	IF	CITATIONS
19	A comparison of the clinical anesthetic efficacy of 4% articaine and 0.5% bupivacaine (both with) Tj ETQq1 1 0.784314 rgBT /Overlock 1 Oral Radiology and Endodontics, 2008, 106, 19-28.	1.6	55
20	The selective and non-selective cyclooxygenase inhibitors valdecoxib and piroxicam induce the same postoperative analgesia and control of trismus and swelling after lower third molar removal. Brazilian Journal of Medical and Biological Research, 2007, 40, 1133-1140.	0.7	29
21	Analgesic and anti-inflammatory dose-response relationship of 7.5 and 15mg meloxicam after lower third molar removal: a double-blind, randomized, crossover study. International Journal of Oral and Maxillofacial Surgery, 2007, 36, 26-31.	0.7	35
22	Epinephrine Concentration (1:100,000 or 1:200,000) Does Not Affect the Clinical Efficacy of 4% Articaine for Lower Third Molar Removal: A Double-Blind, Randomized, Crossover Study. Journal of Oral and Maxillofacial Surgery, 2007, 65, 2445-2452.	0.5	59
23	Efeitos da inalação passiva da fumaça de cigarro sobre as pregas vocais de ratos. Revista Brasileira De Otorrinolaringologia, 2006, 72, 210-216.	0.2	12