## Jean A Quinn

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6351444/publications.pdf Version: 2024-02-01



Ιέλνι Δ Οιμνινι

#	Article	IF	CITATIONS
1	Spatial expression of IKK-alpha is associated with a differential mutational landscape and survival in primary colorectal cancer. British Journal of Cancer, 2022, , .	6.4	2
2	Molecular mechanisms of tumour budding and its association with microenvironment in colorectal cancer. Clinical Science, 2022, 136, 521-535.	4.3	4
3	The Relationship Between the Tumor Cell Expression of Hypoxic Markers and Survival in Patients With ER-positive Invasive Ductal Breast Cancer. Journal of Histochemistry and Cytochemistry, 2022, 70, 479-494.	2.5	4
4	The stress-responsive kinase DYRK2 activates heat shock factor 1 promoting resistance to proteotoxic stress. Cell Death and Differentiation, 2021, 28, 1563-1578.	11.2	19
5	Relationship between immune checkpoint proteins, tumour microenvironment characteristics, and prognosis in primary operable colorectal cancer. Journal of Pathology: Clinical Research, 2021, 7, 121-134.	3.0	17
6	The relationship between β-catenin and patient survival in colorectal cancer systematic review and meta-analysis. Critical Reviews in Oncology/Hematology, 2021, 163, 103337.	4.4	8
7	Systematic review of tumour budding and association with common mutations in patients with colorectal cancer. Critical Reviews in Oncology/Hematology, 2021, 167, 103490.	4.4	3
8	High NRF2 Levels Correlate with Poor Prognosis in Colorectal Cancer Patients and with Sensitivity to the Kinase Inhibitor AT9283 In Vitro. Biomolecules, 2020, 10, 1365.	4.0	22
9	The relationship between members of the canonical NF-kB pathway, tumour microenvironment and cancer specific survival in colorectal cancer patients. Histology and Histopathology, 2020, 35, 569-578.	0.7	1
10	A novel tumorâ€based epithelialâ€ŧoâ€mesenchymal transition score that associates with prognosis and metastasis in patients with Stage II/III colorectal cancer. International Journal of Cancer, 2019, 144, 150-159.	5.1	28
11	The association between markers of tumour cell metabolism, the tumour microenvironment and outcomes in patients with colorectal cancer. International Journal of Cancer, 2019, 144, 2320-2329.	5.1	10
12	Tumour infiltrating lymphocyte expression of PD-1 as a favourable prognostic factor in patients with mismatch repair competent colorectal cancer Journal of Clinical Oncology, 2018, 36, 631-631.	1.6	4
13	High IKKα expression is associated with reduced time to recurrence and cancer specific survival in oestrogen receptor (ER)â€positive breast cancer. International Journal of Cancer, 2017, 140, 1633-1644.	5.1	22
14	Signal Transduction and Activator of Transcription-3 (STAT3) in Patients with Colorectal Cancer: Associations with the Phenotypic Features of the Tumor and Host. Clinical Cancer Research, 2017, 23, 1698-1709.	7.0	38
15	The relationship between the non-canonical NF-κB pathway, tumour microenvironment, systemic inflammation and survival in patients undergoing surgery for colorectal caner Journal of Clinical Oncology, 2017, 35, 631-631.	1.6	0
16	Abstract B125: The relationship between members of the canonical NF-κB pathway, components of the microenvironment and survival in patients with colorectal cancer. , 2016, , .		0
17	PTEN ablation in RasHa/Fos skin carcinogenesis invokes p53-dependent p21 to delay conversion while p53-independent p21 limits progression via cyclin D1/E2 inhibition. Oncogene, 2014, 33, 4132-4143.	5.9	11
18	Intravital FLIM-FRET Imaging Reveals Dasatinib-Induced Spatial Control of Src in Pancreatic Cancer. Cancer Research, 2013, 73, 4674-4686.	0.9	111

Jean A Quinn

#	Article	IF	CITATIONS
19	Organotypic Collagen I Assay: A Malleable Platform to Assess Cell Behaviour in a 3-Dimensional Context. Journal of Visualized Experiments, 2011, , e3089.	0.3	66
20	Keratinocytes stimulate fibroblast hyaluronan synthesis through the release of stratifin: A possible role in the suppression of scar tissue formation. Wound Repair and Regeneration, 2011, 19, 379-386.	3.0	9
21	Spatial Regulation of RhoA Activity during Pancreatic Cancer Cell Invasion Driven by Mutant p53. Cancer Research, 2011, 71, 747-757.	0.9	127
22	4-Methylumbelliferone inhibits tumour cell growth and the activation of stromal hyaluronan synthesis by melanoma cell-derived factors. British Journal of Dermatology, 2010, 162, 1224-1232.	1.5	43
23	A gene on the HER2 amplicon, C35, is an oncogene in breast cancer whose actions are prevented by inhibition of Syk. British Journal of Cancer, 2010, 103, 401-410.	6.4	65
24	Effects of human papillomavirus type 16 E5 deletion mutants on epithelial morphology: functional characterization of each transmembrane domain. Journal of General Virology, 2010, 91, 521-530.	2.9	37
25	Effect of Different Classes of Gadolinium-based Contrast Agents on Control and Nephrogenic Systemic Fibrosis–derived Fibroblast Proliferation. Radiology, 2010, 256, 735-743.	7.3	53
26	Gadodiamide contrast agent â€~activates' fibroblasts: a possible cause of nephrogenic systemic fibrosis. Journal of Pathology, 2008, 214, 584-593.	4.5	124
27	Fos cooperation with PTEN loss elicits keratoacanthoma not carcinoma, owing to p53/p21WAF-induced differentiation triggered by GSK3β inactivation and reduced AKT activity. Journal of Cell Science, 2008, 121, 1758-1769.	2.0	15
28	PTEN Loss Promotes rasHa-Mediated Papillomatogenesis via Dual Up-Regulation of AKT Activity and Cell Cycle Deregulation but Malignant Conversion Proceeds via PTEN-Associated Pathways. Cancer Research, 2006, 66, 1302-1312.	0.9	29
29	Retinoic acid-induced inhibition of metastatic melanoma cell lung colonization and adhesion to endothelium and subendothelial extracellular matrix. Clinical and Experimental Metastasis, 1992, 10, 61-67.	3.3	14
30	Different susceptibilities of melanoma cells to retinoic acid-induced changes in melanotic expression. Biochemical and Biophysical Research Communications, 1988, 155, 773-778.	2.1	52