

Masabumi Shibuya

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6356668/masabumi-shibuya-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

158
papers

16,076
citations

63
h-index

126
g-index

168
ext. papers

17,402
ext. citations

8.2
avg, IF

6.93
L-index

#	Paper	IF	Citations
158	A simple detection method for the serum sFLT1 protein in preeclampsia. <i>Scientific Reports</i> , 2021 , 11, 20613	4.9	2
157	Dysregulation of Amphiregulin stimulates the pathogenesis of cystic lymphangioma. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021 , 118,	11.5	4
156	Lymphangiogenesis induced by vascular endothelial growth factor receptor 1 signaling contributes to the progression of endometriosis in mice. <i>Journal of Pharmacological Sciences</i> , 2020 , 143, 255-263	3.7	11
155	Production of an anti-angiogenic factor sFLT1 is suppressed via promoter hypermethylation of FLT1 gene in choriocarcinoma cells. <i>BMC Cancer</i> , 2020 , 20, 112	4.8	4
154	Absence of VEGFR-1/Flt-1 signaling pathway in mice results in insensitivity to discogenic low back pain in an established disc injury mouse model. <i>Journal of Cellular Physiology</i> , 2020 , 235, 5305-5317	7	5
153	Phosphoethanolamine Accumulation Protects Cancer Cells under Glutamine Starvation through Downregulation of PCYT2. <i>Cell Reports</i> , 2019 , 29, 89-103.e7	10.6	12
152	The role of vascular endothelial growth factor receptor 1 tyrosine kinase signaling in bleomycin-induced pulmonary fibrosis. <i>Biomedicine and Pharmacotherapy</i> , 2019 , 117, 109067	7.5	8
151	VEGF Receptor 1-Expressing Macrophages Recruited from Bone Marrow Enhances Angiogenesis in Endometrial Tissues. <i>Scientific Reports</i> , 2019 , 9, 7037	4.9	19
150	Myc-dependent endothelial proliferation is controlled by phosphotyrosine 1212 in VEGFR receptor-2. <i>EMBO Reports</i> , 2019 , 20, e47845	6.5	14
149	Vascular endothelial growth factor receptor 1 tyrosine kinase signaling facilitates healing of DSS-induced colitis by accumulation of Tregs in ulcer area. <i>Biomedicine and Pharmacotherapy</i> , 2019 , 111, 131-141	7.5	8
148	Vascular endothelial growth factor receptor 1 (VEGFR1) tyrosine kinase signaling facilitates granulation tissue formation with recruitment of VEGFR1 cells from bone marrow. <i>Anatomical Science International</i> , 2018 , 93, 372-383	2	6
147	VEGF-A selectively inhibits FLT1 ectodomain shedding independent of receptor activation and receptor endocytosis. <i>American Journal of Physiology - Cell Physiology</i> , 2018 , 315, C214-C224	5.4	1
146	The Novel Pathogenesis of Retinopathy Mediated by Multiple RTK Signals is Uncovered in Newly Developed Mouse Model. <i>EBioMedicine</i> , 2018 , 31, 190-201	8.8	15
145	VEGFR1-TK signaling protects exacerbation of dextran sulfate sodium-induced colitis in mice. <i>Proceedings for Annual Meeting of the Japanese Pharmacological Society</i> , 2018 , WCP2018, PO3-5-31	0	
144	Effect of Age on the Prognosis of Molecular Abnormalities in Pediatric Acute Myeloid Leukemia. <i>Blood</i> , 2018 , 132, 1506-1506	2.2	
143	HIF-2 but not HIF-1 mediates hypoxia-induced up-regulation of Flt-1 gene expression in placental trophoblasts. <i>Scientific Reports</i> , 2018 , 8, 17375	4.9	21
142	A subset of cerebrovascular pericytes originates from mature macrophages in the very early phase of vascular development in CNS. <i>Scientific Reports</i> , 2017 , 7, 3855	4.9	52

141	Sustained inflammation after pericyte depletion induces irreversible blood-retina barrier breakdown. <i>JCI Insight</i> , 2017 , 2, e90905	9.9	81
140	Flt1/VEGFR1 heterozygosity causes transient embryonic edema. <i>Scientific Reports</i> , 2016 , 6, 27186	4.9	4
139	New anti-cancer chemicals Ertredin and its derivatives, regulate oxidative phosphorylation and glycolysis and suppress sphere formation in vitro and tumor growth in EGFRvIII-transformed cells. <i>BMC Cancer</i> , 2016 , 16, 496	4.8	2
138	Establishment and characterization of a novel VEGF-producing HHV-8-unrelated PEL-like lymphoma cell line, OGU1. <i>European Journal of Haematology</i> , 2016 , 96, 144-51	3.8	1
137	PlGF/VEGFR-1 Signaling Promotes Macrophage Polarization and Accelerated Tumor Progression in Obesity. <i>Clinical Cancer Research</i> , 2016 , 22, 2993-3004	12.9	81
136	Vascular endothelial growth factor receptor-1 (VEGFR-1) signaling enhances angiogenesis in a surgical sponge model. <i>Biomedicine and Pharmacotherapy</i> , 2016 , 78, 140-149	7.5	9
135	Vascular Endothelial Growth Factor Receptor Type1 Signaling Prevents Delayed Wound Healing in Diabetes by Attenuating the Production of IL-1 β by Recruited Macrophages. <i>American Journal of Pathology</i> , 2016 , 186, 1481-98	5.8	40
134	The guanine nucleotide exchange factor Vav3 regulates differentiation of progenitor cells in the developing mouse retina. <i>Cell and Tissue Research</i> , 2015 , 359, 423-440	4.2	12
133	Endothelial Gab1 deficiency aggravates splenomegaly in portal hypertension independent of angiogenesis. <i>American Journal of Physiology - Renal Physiology</i> , 2015 , 308, G416-26	5.1	3
132	Thromboxane A2 induces blood flow recovery via platelet adhesion to ischaemic regions. <i>Cardiovascular Research</i> , 2015 , 107, 509-21	9.9	15
131	Identification and characterization of VEGF-A-responsive neutrophils expressing CD49d, VEGFR1, and CXCR4 in mice and humans. <i>Blood</i> , 2015 , 126, 2016-26	2.2	126
130	The Role of Vascular Endothelial Growth Factor Receptor-1 Signaling in the Recovery from Ischemia. <i>PLoS ONE</i> , 2015 , 10, e0131445	3.7	23
129	VEGF-VEGFR System as a Target for Suppressing Inflammation and other Diseases. <i>Endocrine, Metabolic and Immune Disorders - Drug Targets</i> , 2015 , 15, 135-44	2.2	63
128	VEGFR1-positive macrophages facilitate liver repair and sinusoidal reconstruction after hepatic ischemia/reperfusion injury. <i>PLoS ONE</i> , 2014 , 9, e105533	3.7	28
127	VEGF-VEGFR Signals in Health and Disease. <i>Biomolecules and Therapeutics</i> , 2014 , 22, 1-9	4.2	113
126	Vascular endothelial growth factor and its receptor system: physiological functions in angiogenesis and pathological roles in various diseases. <i>Journal of Biochemistry</i> , 2013 , 153, 13-9	3.1	457
125	The RNA aptamer inhibiting human vesicular endothelial growth factor receptor 1 without affecting cytokine binding. <i>Biochemistry</i> , 2013 , 52, 2274-9	3.2	7
124	Leukotriene B4 type-1 receptor signaling promotes liver repair after hepatic ischemia/reperfusion injury through the enhancement of macrophage recruitment. <i>FASEB Journal</i> , 2013 , 27, 3132-43	0.9	18

123	Inhibition of histone demethylase JMJD1A improves anti-angiogenic therapy and reduces tumor-associated macrophages. <i>Cancer Research</i> , 2013 , 73, 3019-28	10.1	71
122	Vascular Endothelial Growth Factor Receptor Family in Ascidians, <i>Halocynthia roretzi</i> (Sea Squirt). Its High Expression in Circulatory System-Containing Tissues. <i>International Journal of Molecular Sciences</i> , 2013 , 14, 4841-53	6.3	3
121	VEGFR and type-V RTK activation and signaling. <i>Cold Spring Harbor Perspectives in Biology</i> , 2013 , 5, a009092	6.2	67
120	Photoreceptor avascular privilege is shielded by soluble VEGF receptor-1. <i>ELife</i> , 2013 , 2, e00324	8.9	57
119	Soluble FLT1 binds lipid microdomains in podocytes to control cell morphology and glomerular barrier function. <i>Cell</i> , 2012 , 151, 384-99	56.2	115
118	Adventitial gene transfer of VEGFR-2 specific VEGF-E chimera induces MCP-1 expression in vascular smooth muscle cells and enhances neointimal formation. <i>Atherosclerosis</i> , 2011 , 219, 84-91	3.1	24
117	Involvement of Flt-1 (VEGF receptor-1) in cancer and preeclampsia. <i>Proceedings of the Japan Academy Series B: Physical and Biological Sciences</i> , 2011 , 87, 167-78	4	40
116	Downregulation of receptor for activated C-kinase 1 (RACK1) suppresses tumor growth by inhibiting tumor cell proliferation and tumor-associated angiogenesis. <i>Cancer Science</i> , 2011 , 102, 2007-13	6.9	23
115	Increased expression of histone demethylase JHDM1D under nutrient starvation suppresses tumor growth via down-regulating angiogenesis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 20725-9	11.5	53
114	Vascular Endothelial Growth Factor (VEGF) and Its Receptor (VEGFR) Signaling in Angiogenesis: A Crucial Target for Anti- and Pro-Angiogenic Therapies. <i>Genes and Cancer</i> , 2011 , 2, 1097-105	2.9	699
113	Neuronal FLT1 receptor and its selective ligand VEGF-B protect against retrograde degeneration of sensory neurons. <i>FASEB Journal</i> , 2011 , 25, 1461-73	0.9	40
112	RACK1 regulates VEGF/Flt1-mediated cell migration via activation of a PI3K/Akt pathway. <i>Journal of Biological Chemistry</i> , 2011 , 286, 9097-106	5.4	85
111	Grb-2-associated binder 1 (Gab1) regulates postnatal ischemic and VEGF-induced angiogenesis through the protein kinase A-endothelial NOS pathway. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 2957-62	11.5	55
110	Vascular endothelial growth factor receptor-1 signaling promotes liver repair through restoration of liver microvasculature after acetaminophen hepatotoxicity. <i>Toxicological Sciences</i> , 2011 , 120, 218-29	4.4	42
109	The effects of VEGF-R1 and VEGF-R2 ligands on angiogenic responses and left ventricular function in mice. <i>Cardiovascular Research</i> , 2010 , 86, 122-30	9.9	40
108	Tyrosine Kinase Receptor Flt/VEGFR Family: Its Characterization Related to Angiogenesis and Cancer. <i>Genes and Cancer</i> , 2010 , 1, 1119-23	2.9	29
107	Vascular endothelial growth factor receptor-1 signaling promotes mobilization of macrophage lineage cells from bone marrow and stimulates solid tumor growth. <i>Cancer Research</i> , 2010 , 70, 8211-21	10.1	78
106	PLGF blockade does not inhibit angiogenesis during primary tumor growth. <i>Cell</i> , 2010 , 141, 166-77	56.2	137

105	Vascular endothelial growth factor (VEGF) receptor-2 tyrosine 1175 signaling controls VEGF-induced von Willebrand factor release from endothelial cells via phospholipase C-gamma 1- and protein kinase A-dependent pathways. <i>Journal of Biological Chemistry</i> , 2009 , 284, 23217-24	5.4	46
104	Professor Hidesaburo Hanafusa: a 50-year quest for the molecular basis of cancer. <i>Journal of Biochemistry</i> , 2009 , 146, 3-5	3.1	
103	Hypoxia and low-nutrition double stress induces aggressiveness in a murine model of melanoma. <i>Cancer Science</i> , 2009 , 100, 844-51	6.9	27
102	Brain angiogenesis in developmental and pathological processes: therapeutic aspects of vascular endothelial growth factor. <i>FEBS Journal</i> , 2009 , 276, 4636-43	5.7	63
101	Growth inhibition of AML cells with specific chromosome abnormalities by monoclonal antibodies to receptors for vascular endothelial growth factor. <i>Leukemia Research</i> , 2009 , 33, 1650-7	2.7	12
100	Unique signal transduction of the VEGF family members VEGF-A and VEGF-E. <i>Biochemical Society Transactions</i> , 2009 , 37, 1161-6	5.1	6
99	Blocking VEGFR-3 suppresses angiogenic sprouting and vascular network formation. <i>Nature</i> , 2008 , 454, 656-60	50.4	649
98	Novel role for vascular endothelial growth factor (VEGF) receptor-1 and its ligand VEGF-B in motor neuron degeneration. <i>Journal of Neuroscience</i> , 2008 , 28, 10451-9	6.6	104
97	Vascular endothelial growth factor receptor-1 regulates postnatal angiogenesis through inhibition of the excessive activation of Akt. <i>Circulation Research</i> , 2008 , 103, 261-8	15.7	44
96	Flt-1 signaling in macrophages promotes glioma growth in vivo. <i>Cancer Research</i> , 2008 , 68, 7342-51	10.1	139
95	VEGF receptor signal transduction. <i>Methods in Enzymology</i> , 2008 , 443, 261-84	1.7	28
94	VEGFR1 tyrosine kinase signaling promotes lymphangiogenesis as well as angiogenesis indirectly via macrophage recruitment. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2008 , 28, 658-64	9.4	110
93	Inhibition of choroidal neovascularization by blocking vascular endothelial growth factor receptor tyrosine kinase. <i>Japanese Journal of Ophthalmology</i> , 2008 , 52, 91-98	2.6	28
92	Involvement of vascular endothelial growth factor receptor-1 in rheumatoid arthritis. <i>Inflammation and Regeneration</i> , 2008 , 28, 78-85	10.9	1
91	Vascular endothelial growth factor-dependent and -independent regulation of angiogenesis. <i>BMB Reports</i> , 2008 , 41, 278-86	5.5	191
90	Vascular Permeability/Vascular Endothelial Growth Factor 2008 , 89-98		4
89	Ligand-independent activation of vascular endothelial growth factor receptor 1 by low-density lipoprotein. <i>EMBO Reports</i> , 2007 , 8, 1155-61	6.5	29
88	Novel antiangiogenic pathway of thrombospondin-1 mediated by suppression of the cell cycle. <i>Cancer Science</i> , 2007 , 98, 1491-7	6.9	46

87	The lysine 831 of vascular endothelial growth factor receptor 1 is a novel target of methylation by SMYD3. <i>Cancer Research</i> , 2007 , 67, 10759-65	10.1	125
86	Therapeutic angiogenesis using novel vascular endothelial growth factor-E/human placental growth factor chimera genes. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2007 , 27, 99-105	9.4	22
85	Chimeric VEGF-ENZ7/PlGF specifically binding to VEGFR-2 accelerates skin wound healing via enhancement of neovascularization. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2007 , 27, 503-11	9.4	35
84	Distinct vascular endothelial growth factor signals for lymphatic vessel enlargement and sprouting. <i>Journal of Experimental Medicine</i> , 2007 , 204, 1431-40	16.6	137
83	Chimeric VEGF-E(NZ7)/PlGF promotes angiogenesis via VEGFR-2 without significant enhancement of vascular permeability and inflammation. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2006 , 26, 2019-26	9.4	38
82	Vascular endothelial growth factor (VEGF)-Receptor2: its biological functions, major signaling pathway, and specific ligand VEGF-E. <i>Endothelium: Journal of Endothelial Cell Research</i> , 2006 , 13, 63-9		104
81	Signal transduction by VEGF receptors in regulation of angiogenesis and lymphangiogenesis. <i>Experimental Cell Research</i> , 2006 , 312, 549-60	4.2	782
80	KRN951, a highly potent inhibitor of vascular endothelial growth factor receptor tyrosine kinases, has antitumor activities and affects functional vascular properties. <i>Cancer Research</i> , 2006 , 66, 9134-42	10.1	158
79	Signaling of vascular endothelial growth factor receptor-1 tyrosine kinase promotes rheumatoid arthritis through activation of monocytes/macrophages. <i>Blood</i> , 2006 , 108, 1849-56	2.2	142
78	Corneal avascularity is due to soluble VEGF receptor-1. <i>Nature</i> , 2006 , 443, 993-7	50.4	528
77	Differential roles of vascular endothelial growth factor receptor-1 and receptor-2 in angiogenesis. <i>BMB Reports</i> , 2006 , 39, 469-78	5.5	375
76	α-Integrin+ Endothelium Derived from Primate Embryonic Stem Cells Generates Both Primitive and Definitive Hematopoietic Cells.. <i>Blood</i> , 2006 , 108, 683-683	2.2	
75	Different Kinetics and Function of Vascular Endothelial Growth Factor Receptor-1 and α during Hemangioblast Development from Primate Embryonic Stem Cells.. <i>Blood</i> , 2006 , 108, 3920-3920	2.2	
74	Tumor necrosis factor and vascular endothelial growth factor induce endothelial integrin repertoires, regulating endovascular differentiation and apoptosis in a human extravillous trophoblast cell line. <i>Biology of Reproduction</i> , 2005 , 73, 172-9	3.9	41
73	Essential role of Flk-1 (VEGF receptor 2) tyrosine residue 1173 in vasculogenesis in mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005 , 102, 1076-81	11.5	250
72	VEGF receptor 1 signaling is essential for osteoclast development and bone marrow formation in colony-stimulating factor 1-deficient mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005 , 102, 14016-21	11.5	97
71	Vascular endothelial growth factor A (VEGF-A) is involved in guidance of VEGF receptor-positive cells to the anterior portion of early embryos. <i>Molecular and Cellular Biology</i> , 2005 , 25, 355-63	4.8	62
70	The vascular endothelial growth factor (VEGF)/VEGF receptor system and its role under physiological and pathological conditions. <i>Clinical Science</i> , 2005 , 109, 227-41	6.5	628

69	A hypoxia-driven vascular endothelial growth factor/Flt1 autocrine loop interacts with hypoxia-inducible factor-1alpha through mitogen-activated protein kinase/extracellular signal-regulated kinase 1/2 pathway in neuroblastoma. <i>Cancer Research</i> , 2005 , 65, 7267-75	10.1	111
68	Rationale for antiangiogenic cancer therapy with vaccination using epitope peptides derived from human vascular endothelial growth factor receptor 2. <i>Cancer Research</i> , 2005 , 65, 4939-46	10.1	83
67	Membrane fixation of vascular endothelial growth factor receptor 1 ligand-binding domain is important for vasculogenesis and angiogenesis in mice. <i>Molecular and Cellular Biology</i> , 2005 , 25, 346-54	4.8	64
66	Sequential Analysis of the β and ϵ Globin Gene Expressions during Erythropoietic Differentiation from Primate ES Cells.. <i>Blood</i> , 2005 , 106, 1744-1744	2.2	
65	Molecular Basis of Angiogenesis. <i>Ensho Saisei</i> , 2004 , 24, 144-153		
64	A novel snake venom vascular endothelial growth factor (VEGF) predominantly induces vascular permeability through preferential signaling via VEGF receptor-1. <i>Journal of Biological Chemistry</i> , 2004 , 279, 46304-14	5.4	76
63	Blockade of vascular endothelial growth factor suppresses experimental restenosis after intraluminal injury by inhibiting recruitment of monocyte lineage cells. <i>Circulation</i> , 2004 , 110, 2444-52	16.7	117
62	Essential role of vascular endothelial growth factor and Flt-1 signals in neointimal formation after periadventitial injury. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2004 , 24, 2284-9	9.4	75
61	Dynamic regulation of gene expression by the Flt-1 kinase and Matrigel in endothelial tubulogenesis. <i>Genomics</i> , 2004 , 84, 185-92	4.3	12
60	Nox1 regulates apoptosis and potentially stimulates branching morphogenesis in sinusoidal endothelial cells. <i>Experimental Cell Research</i> , 2004 , 300, 455-62	4.2	54
59	Identification and Characterization of Hemoangiogenic Progenitors during Cynomolgus Monkey ES Cell Differentiation.. <i>Blood</i> , 2004 , 104, 3222-3222	2.2	
58	A set of loop-1 and -3 structures in the novel vascular endothelial growth factor (VEGF) family member, VEGF-ENZ-7, is essential for the activation of VEGFR-2 signaling. <i>Journal of Biological Chemistry</i> , 2003 , 278, 13453-61	5.4	39
57	Vascular endothelial growth factor receptor-2: its unique signaling and specific ligand, VEGF-E. <i>Cancer Science</i> , 2003 , 94, 751-6	6.9	105
56	Induction of tube formation by angiopoietin-1 in endothelial cell/fibroblast co-culture is dependent on endogenous VEGF. <i>Cancer Science</i> , 2003 , 94, 782-90	6.9	60
55	Mammalian Sprouty4 suppresses Ras-independent ERK activation by binding to Raf1. <i>Nature Cell Biology</i> , 2003 , 5, 427-32	23.4	214
54	Role of PlGF in the intra- and intermolecular cross talk between the VEGF receptors Flt1 and Flk1. <i>Nature Medicine</i> , 2003 , 9, 936-43	50.5	631
53	VEGFR-2-specific ligand VEGF-E induces non-edematous hyper-vascularization in mice. <i>Biochemical and Biophysical Research Communications</i> , 2003 , 301, 371-7	3.4	78
52	Bone morphogenetic protein 4 mediates apoptosis of capillary endothelial cells during rat pupillary membrane regression. <i>Molecular and Cellular Biology</i> , 2003 , 23, 4627-36	4.8	51

51	MMP9 induction by vascular endothelial growth factor receptor-1 is involved in lung-specific metastasis. <i>Cancer Cell</i> , 2002 , 2, 289-300	24.3	724
50	Vascular endothelial growth factor receptor family genes: when did the three genes phylogenetically segregate?. <i>Biological Chemistry</i> , 2002 , 383, 1573-9	4.5	32
49	Vascular endothelial growth factor is necessary in the development of arteriosclerosis by recruiting/activating monocytes in a rat model of long-term inhibition of nitric oxide synthesis. <i>Circulation</i> , 2002 , 105, 1110-5	16.7	129
48	Soluble Flt-1 (soluble VEGFR-1), a potent natural antiangiogenic molecule in mammals, is phylogenetically conserved in avians. <i>Biochemical and Biophysical Research Communications</i> , 2002 , 291, 554-9	3.4	48
47	Soluble FLT-1 expression suppresses carcinomatous ascites in nude mice bearing ovarian cancer. <i>Cancer Research</i> , 2002 , 62, 2019-23	10.1	51
46	Expression of vascular endothelial growth factor receptors in smooth muscle cells. <i>Journal of Cellular Physiology</i> , 2001 , 188, 359-68	7	178
45	A variant of nuclear localization signal of bipartite-type is required for the nuclear translocation of hypoxia inducible factors (1alpha, 2alpha and 3alpha). <i>Oncogene</i> , 2001 , 20, 1435-44	9.2	65
44	VEGF protects against oxidized LDL toxicity to endothelial cells by an intracellular glutathione-dependent mechanism through the KDR receptor. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2001 , 21, 765-70	9.4	27
43	Flt-1, vascular endothelial growth factor receptor 1, is a novel cell surface marker for the lineage of monocyte-macrophages in humans. <i>Blood</i> , 2001 , 97, 785-91	2.2	404
42	The overexpression of PKCdelta is involved in vascular endothelial growth factor-resistant apoptosis in cultured primary sinusoidal endothelial cells. <i>Biochemical and Biophysical Research Communications</i> , 2001 , 280, 415-20	3.4	14
41	Structure and dual function of vascular endothelial growth factor receptor-1 (Flt-1). <i>International Journal of Biochemistry and Cell Biology</i> , 2001 , 33, 409-20	5.6	200
40	Structure and function of VEGF/VEGF-receptor system involved in angiogenesis. <i>Cell Structure and Function</i> , 2001 , 26, 25-35	2.2	395
39	Properties of two VEGF receptors, Flt-1 and KDR, in signal transduction. <i>Annals of the New York Academy of Sciences</i> , 2000 , 902, 201-5; discussion 205-7	6.5	90
38	Roles of two VEGF receptors, Flt-1 and KDR, in the signal transduction of VEGF effects in human vascular endothelial cells. <i>Oncogene</i> , 2000 , 19, 2138-46	9.2	241
37	VEGF activates protein kinase C-dependent, but Ras-independent Raf-MEK-MAP kinase pathway for DNA synthesis in primary endothelial cells. <i>Oncogene</i> , 1999 , 18, 2221-30	9.2	472
36	Involvement of VEGF and its receptors in ascites tumor formation. <i>Cancer Chemotherapy and Pharmacology</i> , 1999 , 43 Suppl, S72-7	3.5	34
35	Tumorigenicity depends on angiogenic potential of tumor cells: dominant role of vascular endothelial growth factor and/or fibroblast growth factors produced by tumor cells. <i>Angiogenesis</i> , 1998 , 2, 57-66	10.6	10
34	Clotrimazole, an imidazole antimycotic, is a potent inhibitor of angiogenesis. <i>Japanese Journal of Cancer Research</i> , 1998 , 89, 445-51		25

33	Flt-1, a receptor for vascular endothelial growth factor, has transforming and morphogenic potentials. <i>Oncogene</i> , 1998 , 16, 2585-95	9.2	46
32	Mammary carcinoma cells over-expressing tissue inhibitor of metalloproteinases-1 show enhanced vascular endothelial growth factor expression. <i>International Journal of Cancer</i> , 1998 , 75, 81-7	7.5	97
31	Virally activated Ras cooperates with integrin to induce tubulogenesis in sinusoidal endothelial cell lines. <i>Journal of Cellular Physiology</i> , 1998 , 176, 223-34	7	28
30	Genomic organization of the flt-1 gene encoding for vascular endothelial growth factor (VEGF) receptor-1 suggests an intimate evolutionary relationship between the 7-Ig and the 5-Ig tyrosine kinase receptors. <i>Gene</i> , 1998 , 208, 297-305	3.8	76
29	Mapping of the sites involved in ligand association and dissociation at the extracellular domain of the kinase insert domain-containing receptor for vascular endothelial growth factor. <i>Journal of Biological Chemistry</i> , 1998 , 273, 31283-8	5.4	123
28	Involvement of MAP kinase-independent protein kinase C signaling pathway in the EGF-induced p21(WAF1/Cip1) expression and growth inhibition of A431 cells. <i>Biochemical and Biophysical Research Communications</i> , 1998 , 250, 430-5	3.4	25
27	Angiotensin II type 1 receptor-induced extracellular signal-regulated protein kinase activation is mediated by Ca ²⁺ /calmodulin-dependent transactivation of epidermal growth factor receptor. <i>Circulation Research</i> , 1998 , 82, 1338-48	15.7	172
26	A novel type of vascular endothelial growth factor, VEGF-E (NZ-7 VEGF), preferentially utilizes KDR/Flk-1 receptor and carries a potent mitotic activity without heparin-binding domain. <i>Journal of Biological Chemistry</i> , 1998 , 273, 31273-82	5.4	283
25	The mechanisms of hepatic sinusoidal endothelial cell regeneration: A possible communication system associated with vascular endothelial growth factor in liver cells. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 1998 , 13, S1-S5	4	21
24	Vascular Endothelial Growth Factor Expression in the Rat Uterus and Placenta throughout Pregnancy.. <i>Acta Histochemica Et Cytochemica</i> , 1998 , 31, 419-426	1.9	2
23	Flt-1 lacking the tyrosine kinase domain is sufficient for normal development and angiogenesis in mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998 , 95, 9349-54	11.5	856
22	How do endothelial cells form vascular structure in vitro?. <i>The Journal of Japan Atherosclerosis Society</i> , 1998 , 25, 383-387		
21	Mammary carcinoma cells over-expressing tissue inhibitor of metalloproteinases-1 show vascular endothelial growth factor expression 1998 , 75, 81		2
20	The phosphorylated 1169-tyrosine containing region of flt-1 kinase (VEGFR-1) is a major binding site for PLCgamma. <i>Biochemical and Biophysical Research Communications</i> , 1997 , 238, 487-91	3.4	107
19	Characterization of the extracellular domain in vascular endothelial growth factor receptor-1 (Flt-1 tyrosine kinase). <i>Japanese Journal of Cancer Research</i> , 1997 , 88, 867-76		54
18	The 230 kDa mature form of KDR/Flk-1 (VEGF receptor-2) activates the PLC-gamma pathway and partially induces mitotic signals in NIH3T3 fibroblasts. <i>Oncogene</i> , 1997 , 14, 2079-89	9.2	264
17	Deletion of the ABL SH3 domain reactivates de-oligomerized BCR-ABL for growth factor independence. <i>FEBS Letters</i> , 1996 , 379, 244-6	3.8	18
16	Characterization of the promoter region for flt-1 tyrosine kinase gene, a receptor for vascular endothelial growth factor. <i>Growth Factors</i> , 1996 , 13, 151-62	1.6	20

15	A cAMP response element and an Ets motif are involved in the transcriptional regulation of flt-1 tyrosine kinase (vascular endothelial growth factor receptor 1) gene. <i>Journal of Biological Chemistry</i> , 1996 , 271, 30823-8	5.4	118
14	A Novel Endothelial Growth Factor VEGF and Its Receptor. <i>Japanese Journal of Thrombosis and Hemostasis</i> , 1996 , 7, 96-101	0	
13	Role of VEGF-flt receptor system in normal and tumor angiogenesis. <i>Advances in Cancer Research</i> , 1995 , 67, 281-316	5.9	238
12	Germ-line and somatic mutations of the APC gene in patients with Turcot syndrome and analysis of APC mutations in brain tumors. <i>Genes Chromosomes and Cancer</i> , 1994 , 9, 168-72	5	88
11	Establishment of a human small cell lung carcinoma cell line carrying amplification of c-myc gene and chromosomal translocation of t(3p;6p) and t(12q;17p). <i>Japanese Journal of Cancer Research</i> , 1993 , 84, 355-9		1
10	Inhibition of epidermal growth factor receptor functions by tyrosine kinase inhibitors in NIH3T3 cells. <i>FEBS Letters</i> , 1992 , 314, 289-92	3.8	13
9	In situ localization of male germ cell-associated kinase (mak) mRNA in adult mouse testis: specific expression in germ cells at stages around meiotic cell division. <i>Cell Biochemistry and Function</i> , 1992 , 10, 273-9	4.2	15
8	Alterations of mouse proto-oncogenes in sarcomas induced after transplantation of human tumors in athymic nude mice. <i>Japanese Journal of Cancer Research</i> , 1990 , 81, 333-9		1
7	A deletion mutation within the ligand binding domain is responsible for activation of epidermal growth factor receptor gene in human brain tumors. <i>Japanese Journal of Cancer Research</i> , 1990 , 81, 773-9		120
6	Co-amplification of c-myc and c-erbB-2 oncogenes in a poorly differentiated human gastric cancer. <i>Japanese Journal of Cancer Research</i> , 1989 , 80, 920-3		10
5	Undetectable bcr-abl rearrangements in some CML patients are due to a deletion mutation in the bcr gene. <i>American Journal of Hematology</i> , 1988 , 28, 33-6	7.1	19
4	Structural abnormality and over-expression of the myc gene in feline leukemias. <i>International Journal of Cancer</i> , 1987 , 40, 564-9	7.5	37
3	Nucleotide sequence of Fujinami sarcoma virus: evolutionary relationship of its transforming gene with transforming genes of other sarcoma viruses. <i>Cell</i> , 1982 , 30, 787-95	56.2	238
2	Transcription of the E. coli tufB gene: cotranscription with four tRNA genes and inhibition by guanosine-5Rdiphosphate-3Rdiphosphate. <i>Molecular Genetics and Genomics</i> , 1981 , 183, 13-9		31
1	Construction and characterization of the two hybrid Co1E1 plasmids carrying Escherichia coli tufB gene. <i>FEBS Letters</i> , 1979 , 102, 207-10	3.8	34