List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6359204/publications.pdf Version: 2024-02-01



ALICIA E TOPANZO

#	Article	IF	CITATIONS
1	Evidence for the facultative intracellular behaviour of the fish pathogen <i>Vibrio ordalii</i> . Journal of Fish Diseases, 2019, 42, 1447-1455.	1.9	4
2	Biofilm development and cell viability: An undervalued mechanism in the persistence of the fish pathogen Tenacibaculum maritimum. Aquaculture, 2019, 511, 734267.	3.5	18
3	The Fish Pathogen Vibrio ordalii Under Iron Deprivation Produces the Siderophore Piscibactin. Microorganisms, 2019, 7, 313.	3.6	15
4	Genetic studies to re-affiliate Edwardsiella tarda fish isolates to Edwardsiella piscicida and Edwardsiella anguillarum species. Systematic and Applied Microbiology, 2018, 41, 30-37.	2.8	58
5	Population genetic and evolution analysis of controversial genus Edwardsiella by multilocus sequence typing. Molecular Phylogenetics and Evolution, 2018, 127, 513-521.	2.7	11
6	Edwardsiella piscicida: a significant bacterial pathogen of cultured fish. Diseases of Aquatic Organisms, 2018, 131, 59-71.	1.0	71
7	Draft Genome Sequence of <i>Edwardsiella piscicida</i> Strain ACC35.1 Isolated from Diseased Turbot (Scophthalmus maximus) in Europe. Genome Announcements, 2017, 5, .	0.8	4
8	Insights into the virulenceâ€related genes of <i>Edwardsiella tarda</i> isolated from turbot in Europe: genetic homogeneity and evidence for vibrioferrin production. Journal of Fish Diseases, 2016, 39, 565-576.	1.9	11
9	Iron assimilation and siderophore production by Vibrio ordalii strains isolated from diseased Atlantic salmon Salmo salar in Chile. Diseases of Aquatic Organisms, 2016, 118, 217-226.	1.0	9
10	Survival behaviour and virulence of the fish pathogen Vibrio ordalii in seawater microcosms. Diseases of Aquatic Organisms, 2016, 120, 27-38.	1.0	6
11	Cell-surface properties of Vibrio ordalii strains isolated from Atlantic salmon Salmo salar in Chilean farms. Diseases of Aquatic Organisms, 2015, 113, 9-23.	1.0	16
12	First Isolation and Characterization of Tenacibaculum soleae from Sea Bass Dicentrarchus labrax. Fish Pathology, 2014, 49, 16-22.	0.7	5
13	Vibrio ostreicida sp. nov., a new pathogen of bivalve larvae. International Journal of Systematic and Evolutionary Microbiology, 2014, 64, 1641-1646.	1.7	26
14	Multilocus Sequence Analysis of the Marine Bacterial Genus Tenacibaculum Suggests Parallel Evolution of Fish Pathogenicity and Endemic Colonization of Aquaculture Systems. Applied and Environmental Microbiology, 2014, 80, 5503-5514.	3.1	69
15	Comparative polyphasic characterization of Streptococcus phocae strains with different host origin and description of the subspecies Streptococcus phocae subsp. salmonis subsp. nov International Journal of Systematic and Evolutionary Microbiology, 2014, 64, 1775-1781.	1.7	19
16	A multiplex PCR for the simultaneous detection of Tenacibaculum maritimum and Edwardsiella tarda in aquaculture. International Microbiology, 2014, 17, 111-7.	2.4	10
17	Vibrio ordalii antimicrobial susceptibility testing—Modified culture conditions required and laboratory-specific epidemiological cut-off values. Veterinary Microbiology, 2013, 165, 434-442.	1.9	11
18	Pseudomonas baetica sp. nov., a fish pathogen isolated from wedge sole, Dicologlossa cuneata (Moreau). International Journal of Systematic and Evolutionary Microbiology, 2012, 62, 874-882.	1.7	56

#	Article	IF	CITATIONS
19	Iron utilization and siderophore production by Streptococcus phocae isolated from diseased Atlantic salmon (Salmo salar). Aquaculture, 2012, 364-365, 305-311.	3.5	8
20	First description of <i>Edwardsiella tarda</i> in Senegalese sole, <i>Solea senegalensis</i> (Kaup). Journal of Fish Diseases, 2012, 35, 79-82.	1.9	12
21	Estimation of epidemiological cut-off values for disk diffusion susceptibility test data for Streptococcus phocae. Aquaculture, 2011, 314, 44-48.	3.5	13
22	Characterization of Vibrio tapetis strains isolated from diseased cultured Wedge sole (Dicologoglossa cuneata Moreau). Research in Veterinary Science, 2011, 90, 189-195.	1.9	15
23	Intraspecific genetic variability of Edwardsiella tarda strains from cultured turbot. Diseases of Aquatic Organisms, 2011, 95, 253-258.	1.0	19
24	Pathogenic Potential of Edwardsiella tarda Strains Isolated from Turbot. Fish Pathology, 2011, 46, 27-30.	0.7	5
25	Serological and molecular heterogeneity among Yersinia ruckeri strains isolated from farmed Atlantic salmon Salmo salar in Chile. Diseases of Aquatic Organisms, 2011, 93, 207-214.	1.0	33
26	Surface properties of Streptococcus phocae strains isolated from diseased Atlantic salmon, Salmo salar L. Journal of Fish Diseases, 2011, 34, 203-215.	1.9	30
27	Pseudo-membranes on internal organs associated with Rhodococcus qingshengii infection in Atlantic salmon (Salmo salar). Veterinary Microbiology, 2011, 147, 200-204.	1.9	8
28	Multiplex PCR for the detection of Piscirickettsia salmonis, Vibrio anguillarum, Aeromonas salmonicida and Streptococcus phocae in Chilean marine farms. Diseases of Aquatic Organisms, 2011, 97, 135-142.	1.0	22
29	Contact Lens Hydrophobicity and Roughness Effects on Bacterial Adhesion. Optometry and Vision Science, 2010, 87, E426-E431.	1.2	64
30	Evaluation of four polymerase chain reaction primer pairs for the detection of Edwardsiella tarda in turbot. Diseases of Aquatic Organisms, 2010, 90, 55-61.	1.0	29
31	Vibrio gallaecicus sp. nov. isolated from cultured clams in north-western Spain. Systematic and Applied Microbiology, 2009, 32, 111-117.	2.8	33
32	Genetic characterization of <i>Streptococcus phocae</i> strains isolated from Atlantic salmon, <i>Salmo salar</i> L., in Chile. Journal of Fish Diseases, 2009, 32, 351-358.	1.9	18
33	First isolation of <i>Tenacibaculum maritimum</i> from wedge sole, <i>Dicologoglossa cuneata</i> (Moreau). Journal of Fish Diseases, 2009, 32, 603-610.	1.9	15
34	Identification of Vibrio harveyi isolated from diseased cultured wedge sole Dicologoglossa cuneata. Diseases of Aquatic Organisms, 2009, 84, 209-217.	1.0	19
35	Vaccination Strategies to Prevent Streptococcal Infections in Cultured Fish. , 2009, , 111-149.		1
36	Evolution of drug resistance and minimum inhibitory concentration to enrofloxacin in Tenacibaculum maritimum strains isolated in fish farms. Aquaculture International, 2008, 16, 1-11.	2.2	23

#	Article	IF	CITATIONS
37	First description of serotype O3 in <i>Vibrio anguillarum</i> strains isolated from salmonids in Chile. Journal of Fish Diseases, 2008, 31, 235-239.	1.9	30
38	Streptococcus phocae, an emerging pathogen for salmonid culture. Veterinary Microbiology, 2008, 130, 198-207.	1.9	56
39	Antigenic and molecular characterization of Vibrio ordalii strains isolated from Atlantic salmon Salmo salar in Chile. Diseases of Aquatic Organisms, 2008, 79, 27-35.	1.0	21
40	Development of an effective Edwardsiella tarda vaccine for cultured turbot (Scophthalmus maximus). Fish and Shellfish Immunology, 2008, 25, 208-212.	3.6	59
41	Polymerase Chain Reaction Amplification of Repetitive Intergenic Consensus and Repetitive Extragenic Palindromic Sequences for Molecular Typing of <i>Pseudomonas anguilliseptica</i> and <i>Aeromonas salmonicida</i> . Journal of Aquatic Animal Health, 2008, 20, 75-85.	1.4	14
42	Evaluation of different DNA-based fingerprinting methods for typing Photobacterium damselae ssp. piscicida. Biological Research, 2007, 40, 85-92.	3.4	13
43	Use of adjuvanted vaccines to lengthen the protection against lactococcosis in rainbow trout (Oncorhynchus mykiss). Aquaculture, 2006, 251, 153-158.	3.5	37
44	Use of hydrogen peroxide against the fish pathogen Tenacibaculum maritimum and its effect on infected turbot (Scophthalmus maximus). Aquaculture, 2006, 257, 104-110.	3.5	60
45	Tenacibaculosis infection in marine fish caused by Tenacibaculum maritimum: a review. Diseases of Aquatic Organisms, 2006, 71, 255-266.	1.0	215
46	Use of microcosms to determine the survival of the fish pathogen Tenacibaculum maritimum in seawater. Environmental Microbiology, 2006, 8, 921-928.	3.8	26
47	A challenge model for Tenacibaculum maritimum infection in turbot, Scophthalmus maximus (L.). Journal of Fish Diseases, 2006, 29, 371-374.	1.9	43
48	Characterization of Edwardsiella tarda strains isolated from turbot, Psetta maxima (L.). Journal of Fish Diseases, 2006, 29, 541-547.	1.9	103
49	Effectiveness of a divalent vaccine for sole, Solea senegalensis (Kaup), against Vibrio harveyi and Photobacterium damselae subsp. piscicida. Journal of Fish Diseases, 2005, 28, 33-38.	1.9	60
50	Recommendation of an Appropriate Medium for In Vitro Drug Susceptibility Testing of the Fish Pathogen Tenacibaculum maritimum. Antimicrobial Agents and Chemotherapy, 2005, 49, 82-87.	3.2	15
51	Iron Uptake Mechanisms in the Fish Pathogen Tenacibaculum maritimum. Applied and Environmental Microbiology, 2005, 71, 6947-6953.	3.1	34
52	Variation in 16S-23S rRNA Intergenic Spacer Regions in Photobacterium damselae : a Mosaic-Like Structure. Applied and Environmental Microbiology, 2005, 71, 636-645.	3.1	41
53	A review of the main bacterial fish diseases in mariculture systems. Aquaculture, 2005, 246, 37-61.	3.5	671
54	Development and Validation of a PCR-based Protocol for the Detection of Pseudomonas anguilliseptica. Fish Pathology, 2004, 39, 33-41.	0.7	13

ALICIA E TORANZO

#	Article	IF	CITATIONS
55	Intraspecific diversity of the marine fish pathogen Tenacibaculum maritimum as determined by randomly amplified polymorphic DNA-PCR. Journal of Applied Microbiology, 2004, 96, 871-877.	3.1	29
56	Oral immunization using alginate microparticles as a useful strategy for booster vaccination against fish lactoccocosis. Aquaculture, 2004, 236, 119-129.	3.5	86
57	Species-specific polymerase chain reaction primer sets for the diagnosis of Tenacibaculum maritimum infection. Diseases of Aquatic Organisms, 2004, 62, 75-83.	1.0	45
58	Phenotypic characterization and description of two major O-serotypes in Tenacibaculum maritimum strains from marine fishes. Diseases of Aquatic Organisms, 2004, 58, 1-8.	1.0	39
59	Characterization of the 23S and 5S rRNA genes and 23S-5S intergenic spacer region (ITS-2) of Photobacterium damselae. Diseases of Aquatic Organisms, 2004, 61, 33-39.	1.0	5
60	Isolation of mutants of Vibrio anguillarum defective in haeme utilisation and cloning of huvA, a gene coding for an outer membrane protein involved in the use of haeme as iron source. Archives of Microbiology, 2003, 179, 329-338.	2.2	26
61	Existence of two O-serotypes in the fish pathogen Pseudomonas anguilliseptica. Veterinary Microbiology, 2003, 94, 325-333.	1.9	16
62	lsolation of a highly pathogenic Vibrio pelagius strain associated with mass mortalities of turbot, Scophthalmus maximus (L.), larvae. Journal of Fish Diseases, 2003, 26, 293-303.	1.9	44
63	Phenotypic and Genetic Characterization ofPseudomonas anguillisepticaStrains Isolated from Fish. Journal of Aquatic Animal Health, 2003, 15, 39-47.	1.4	26
64	Molecular Fingerprinting of Fish-Pathogenic Lactococcus garvieae Strains by Random Amplified Polymorphic DNA Analysis. Journal of Clinical Microbiology, 2003, 41, 751-756.	3.9	62
65	Lactococcus garvieae in wild Red Sea wrasse Coris aygula (Labridae). Diseases of Aquatic Organisms, 2003, 56, 275-278.	1.0	30
66	Dormancy as a survival strategy of the fish pathogen Streptococcus parauberis in the marine environment. Diseases of Aquatic Organisms, 2002, 52, 129-136.	1.0	28
67	Molecular Approaches for the Study and Diagnosis of Salmonid Streptococcosis. Reviews: Methods and Technologies in Fish Biology and Fisheries, 2002, , 211-233.	0.6	15
68	Binding of haemin by the fish pathogen Photobacterium damselae subsp. piscicida. Diseases of Aquatic Organisms, 2002, 48, 109-115.	1.0	23
69	Effect of temperature on the development of pasteurellosis in carrier gilthead seabream (Sparus) Tj ETQq1 1	0.784314 rgB	T /Overlock
70	Molecular characterization of Portuguese strains of Yersinia ruckeri isolated from fish culture systems. Journal of Fish Diseases, 2001, 24, 151-159.	1.9	14
71	Multiplex PCR assay for ureC and 16S rRNA genes clearly discriminates between both subspecies of Photobacterium damselae. Diseases of Aquatic Organisms, 2000, 40, 177-183.	1.0	86
72	Existence of two geographically-linked clonal lineages in the bacterial fish pathogen Photobacterium damselae subsp. piscicida evidenced by random amplified polymorphic DNA analysis. Epidemiology and Infection, 2000, 125, 213-219.	2.1	35

#	Article	IF	CITATIONS
73	Evidence that water transmits the disease caused by the fish pathogen Photobacterium damselae subsp. damselae. Journal of Applied Microbiology, 2000, 88, 531-535.	3.1	90
74	Presence of phospholipase-D (dly) gene coding for damselysin production is not a pre-requisite for pathogenicity in Photobacterium damselae subsp. damselae. Microbial Pathogenesis, 2000, 28, 119-126.	2.9	25
75	Electrotransformation of Photobacterium damselae subsp. piscicida. , 2000, , 175-181.		1
76	Electrotransformation of Yersinia ruckeri. , 2000, , 127-133.		0
77	16S rRNA Gene Sequence Analysis of <i>Photobacterium damselae</i> and Nested PCR Method for Rapid Detection of the Causative Agent of Fish Pasteurellosis. Applied and Environmental Microbiology, 1999, 65, 2942-2946.	3.1	94
78	Genetic analysis of turbot pathogenicStreptococcus parauberisstrains by ribotyping and random amplified polymorphic DNA. FEMS Microbiology Letters, 1999, 179, 297-304.	1.8	38
79	Assessment of a magnetic bead-EIA based kit for rapid diagnosis of fish pasteurellosis. Journal of Microbiological Methods, 1999, 38, 147-154.	1.6	12
80	Sanitary control in Spanish aquaculture. Journal of Applied Ichthyology, 1998, 14, 275-277.	0.7	1
81	EDWARDSIELLOSIS IN WILD STRIPED BASS FROM THE CHESAPEAKE BAY. Journal of Wildlife Diseases, 1997, 33, 517-525.	0.8	30
82	Applicability of Ribotyping for Intraspecific Classification and Epidemiological Studies of Photobacterium damsela subsp. piscicida. Systematic and Applied Microbiology, 1997, 20, 634-639.	2.8	22
83	Viability of starved Pasteurella piscicida in seawater monitored by flow cytometry and the effect of antibiotics on its resuscitation. Letters in Applied Microbiology, 1997, 24, 122-126.	2.2	25
84	Phenotypic, serological and molecular characterization of Portuguese isolates of Aeromonas salmonicida subsp. salmonicida Fish Pathology, 1997, 32, 177-178.	0.7	2
85	Antigenic characterization of Vibrio anguillarum-related organisms isolated from turbot and cod. Diseases of Aquatic Organisms, 1997, 28, 45-50.	1.0	10
86	Association ofAeromonas hydrophilaandVibrio alginolyticuswith Larval Mortalities of Scallop (Argopecten purpuratus). Journal of Invertebrate Pathology, 1996, 67, 213-218.	3.2	67
87	Influence of the capsular layer on the virulence ofPasteurella piscicidafor fish. Microbial Pathogenesis, 1996, 21, 289-297.	2.9	62
88	Adherence and invasive capacities of the fish pathogenPasteurella piscicida. FEMS Microbiology Letters, 1996, 138, 29-34.	1.8	57
89	Phenotypic and pathobiological characteristics of Pasteurella piscicida. Annual Review of Fish Diseases, 1996, 6, 41-64.	1.0	66
90	Health status of two salmonid aquaculture facilities in North Portugal: characterization of the bacterial and viral pathogens causing notifiable diseases. Journal of Fish Diseases, 1996, 19, 83-89.	1.9	13

#	Article	IF	CITATIONS
91	Evaluation of media for the successful culture of Flexibacter maritimus. Journal of Fish Diseases, 1996, 19, 193-197.	1.9	103
92	Immunological analysis of extracellular products and cell surface components of motile Aeromonas isolated from fish. Journal of Applied Bacteriology, 1996, 81, 585-593.	1.1	6
93	Phenotypic and pathobiological characteristics of Pasteurella piscicida. Annual Review of Fish Diseases, 1996, 6, 41-64.	1.0	52
94	Occurrence of an erythrocytic virus infection in cultured turbot Scophthalmus maximus. Diseases of Aquatic Organisms, 1996, 24, 159-167.	1.0	7
95	Biochemical and serological analysis of Vibrio anguillarum related organisms. Diseases of Aquatic Organisms, 1996, 26, 67-73.	1.0	6
96	Pathogenicity studies on a Vibrio anguillarum-related (VAR) strain causing an epizootic in Argopecten purpuratus larvae cultured in Chile. Diseases of Aquatic Organisms, 1995, 22, 135-141.	1.0	82
97	In vitro killing of Pasteurella piscicida by fish macrophages. Diseases of Aquatic Organisms, 1995, 23, 51-57.	1.0	38
98	Pathogenicity of live bacteria and extracellular products of motile <i>Aeromonas</i> isolated from eels. Journal of Applied Bacteriology, 1995, 78, 555-562.	1.1	34
99	Pathology associated with a viral erythrocytic infection in turbot, Scophthalmus maximus (L.). Journal of Fish Diseases, 1995, 18, 425-433.	1.9	3
100	Pathology of Enterococcus sp. infection in farmed turbot, Scophthalmus maximus L Journal of Fish Diseases, 1995, 18, 21-30.	1.9	28
101	Histopathological lesions caused by Vibrio damsela in cultured turbot, Scophthalmus maximus (L.): inoculations with live cells and extracellular products. Journal of Fish Diseases, 1995, 18, 357-364.	1.9	23
102	Genetic transformation ofVibrio anguillarumandPasteurella piscicidaby electroporation. FEMS Microbiology Letters, 1995, 128, 75-80.	1.8	8
103	Pathogenicity of birnaviruses isolated from turbot (Scophthalmus maximus): comparison with reference serotypes of IPNV. Aquaculture, 1995, 130, 7-14.	3.5	12
104	Efficacy of intraperitoneal and immersion vaccination against Enterococcus sp. infection in turbot. Aquaculture, 1995, 134, 17-27.	3.5	61
105	Fatty acid profiles of ?Pasteurella? piscicida: comparison with other fish pathogenic gram-negative bacteria. Archives of Microbiology, 1995, 163, 211-216.	2.2	11
106	Response of Pasteurella piscicida and Flexibacter maritimus to skin mucus of marine fish. Diseases of Aquatic Organisms, 1995, 21, 103-108.	1.0	85
107	Fatty acid profiles of "Pasteurella" piscicida: comparison with other fish pathogenic gram-negative bacteria. Archives of Microbiology, 1995, 163, 211-216.	2.2	12
108	Analysis of antigens present in the extracellular products and cell surface of Vibrio anguillarum serotypes O1, O2, and O3. Applied and Environmental Microbiology, 1995, 61, 2493-2498.	3.1	33

#	Article	IF	CITATIONS
109	Evaluation of BIONOR Mono-kits for rapid detection of bacterial fish pathogens. Diseases of Aquatic Organisms, 1995, 21, 25-34.	1.0	31
110	Sequential pathology of experimental pasteurellosis in gilthead seabream Sparus aurata. A light- and electron-microscopic study. Diseases of Aquatic Organisms, 1995, 21, 177-186.	1.0	47
111	Antigenic characterization of Enterococcus strains pathogenic for turbot and their relationship with other Gram-positive bacteria. Diseases of Aquatic Organisms, 1995, 21, 187-191.	1.0	22
112	A Comparison of Pathological Changes Caused by Vibrio anguillarum and Its Extracellular Products in Rainbow Trout(Oncorhynchus mykiss) Fish Pathology, 1994, 29, 79-89.	0.7	29
113	O-Serogrouping and surface components ofAeromonas hydrophilaandAeromonas jandaeipathogenic for eels. FEMS Microbiology Letters, 1994, 117, 85-90.	1.8	39
114	Capsular polysaccharide expressed byPasteurella piscicidagrown in vitro. FEMS Microbiology Letters, 1994, 124, 285-289.	1.8	32
115	Non-specific cellular responses of rainbow trout to Vibrio anguillarum and its extracellular products (ECPs). Journal of Fish Biology, 1994, 45, 839-854.	1.6	45
116	Incidence of Yersinia ruckeri in two farms in Galicia (NW Spain) during a one-year period. Journal of Fish Diseases, 1994, 17, 533-539.	1.9	7
117	Starvation-Survival Processes of the Bacterial Fish Pathogen Yersinia ruckeri. Systematic and Applied Microbiology, 1994, 17, 161-168.	2.8	29
118	Kinetics of antibody production against Vibrio anguillarum antigens in turbot. Aquaculture, 1994, 123, 191-196.	3.5	13
119	Role of serum antibodies in protection of vaccinated turbot (Scophthalmus maximus) against vibriosis. Aquaculture, 1994, 123, 197-204.	3.5	13
120	Vaccination trials on gilthead seabream (Sparus aurata) against Pasteurella piscicida. Aquaculture, 1994, 120, 201-208.	3.5	50
121	Role of iron, capsule, and toxins in the pathogenicity of Vibrio vulnificus biotype 2 for mice. Infection and Immunity, 1994, 62, 759-763.	2.2	102
122	Histopathology of rainbow trout, Oncorhynchus mykiss (Walbaum), and striped bass, Morone saxatilis (Walbaum), experimentally infected with Carnobacterium piscicola. Journal of Fish Diseases, 1993, 16, 261-267.	1.9	12
123	Pathological activities ofYersinia ruckeri, the Enteric Redmouth (ERM) bacterium. FEMS Microbiology Letters, 1993, 112, 291-300.	1.8	62
124	Toxicity of the extracellular products ofVibrio damsela isolated from diseased fish. Current Microbiology, 1993, 27, 341-347.	2.2	48
125	Usefulness of the API-20E system for the identification of bacterial fish pathogens. Aquaculture, 1993, 116, 111-120.	3.5	52
126	Microflora associated with healthy and diseased turbot (Scophthalmus maximus) from three farms in northwest Spain. Aquaculture, 1993, 114, 189-202.	3.5	55

#	Article	IF	CITATIONS
127	Virulence factors of bacteria pathogenic for coldwater fish. Annual Review of Fish Diseases, 1993, 3, 5-36.	1.0	73
128	Antigenic and Molecular Characterization of Yersinia ruckeri Proposal for a New Intraspecies Classification. Systematic and Applied Microbiology, 1993, 16, 411-419.	2.8	70
129	Detection of a Common Antigen amongRenibacterium salmoninarum,Corynebacterium aquaticum, andCarnobacterium piscicolaby the Western Blot Technique. Journal of Aquatic Animal Health, 1993, 5, 172-176.	1.4	8
130	Pathological activities of Yersinia ruckeri, the Enteric Redmouth (ERM) bacterium. FEMS Microbiology Letters, 1993, 112, 291-299.	1.8	2
131	Phenotypic Characteristics and Virulence of <i>Vibrio anguillarum</i> -Related Organisms. Applied and Environmental Microbiology, 1993, 59, 2969-2976.	3.1	38
132	Vibrio mimicus and Vibrio cholerae non-01 Isolated from Wild and Hatchery-Reared Fish Fish Pathology, 1993, 28, 15-26.	0.7	4
133	Characterization of a birnavirus isolated from diseased turbot cultured in Spain. Diseases of Aquatic Organisms, 1993, 15, 163-163.	1.0	25
134	Isolation of IPN virus serotype VR-299 from turbot in Europe. Diseases of Aquatic Organisms, 1993, 17, 61-65.	1.0	11
135	An epizootic in farmed, market-sized rainbow trout in Spain caused by a strain of Carnobactenum piscicola of unusual virulence. Diseases of Aquatic Organisms, 1993, 17, 87-99.	1.0	37
136	Detection of a vascular permeability factor in the extracellular products of Renibacterium salmoninarum. Microbial Pathogenesis, 1992, 13, 237-241.	2.9	2
137	Epizootic study in a turbot farm: bacteriology, virology, parasitology and histology. Aquaculture, 1992, 107, 253-258.	3.5	14
138	Comparison of the extracellular biological activities of Vibrio anguillarum and Aeromonas hydrophila. Aquaculture, 1992, 107, 259-270.	3.5	22
139	The detection of two antigenic groups amongRenibacterium salmoninarumisolates. FEMS Microbiology Letters, 1992, 94, 105-110.	1.8	2
140	Serratia marcescens: a potential pathogen for fish. Journal of Fish Diseases, 1992, 15, 15-26.	1.9	42
141	Antigenic relationships among rotaviruses isolated from fish. Journal of Fish Diseases, 1992, 15, 27-36.	1.9	16
142	Presence of skin permeability factors in the extracellular products ofYersinia ruckeri. Current Microbiology, 1992, 24, 263-267.	2.2	8
143	Comparative study of phenotypic and virulence properties in Vibrio vulnificus biotypes 1 and 2 obtained from a European eel farm experiencing mortalities. Diseases of Aquatic Organisms, 1992, 13, 29-35.	1.0	78
144	Phenotypic and pathobiological properties of Corynebacterium aquaticum isolated from diseased striped bass. Diseases of Aquatic Organisms, 1992, 14, 115-126.	1.0	26

ALICIA E TORANZO

#	Article	IF	CITATIONS
145	Tetramicra brevifilum, a potential threat to farmed turbot Scophthalmus maximus. Diseases of Aquatic Organisms, 1992, 14, 127-135.	1.0	30
146	Characterization of strains related to brown ring disease outbreaks in southwestern Spain. Diseases of Aquatic Organisms, 1992, 14, 229-236.	1.0	35
147	MICs and MBCs of chemotherapeutic agents against Renibacterium salmoninarum. Antimicrobial Agents and Chemotherapy, 1991, 35, 1011-1013.	3.2	25
148	Lack of Biological Activities in the Extracellular Products of <i>Renibacterium salmoninarum</i> . Canadian Journal of Fisheries and Aquatic Sciences, 1991, 48, 421-425.	1.4	23
149	Pasteurellosis in cultured gilthead seabream (Sparus aurata): first report in Spain. Aquaculture, 1991, 99, 1-15.	3.5	143
150	Biochemical and Serological Characteristics, Drug Resistance and Plasmid Profiles of Spanish Isolates of Aeromonas salmonicida Fish Pathology, 1991, 26, 55-60.	0.7	23
151	Cell-Surface-Associated Properties of Fish Pathogenic Bacteria. Journal of Aquatic Animal Health, 1991, 3, 297-301.	1.4	30
152	Susceptibility of turbot (Scophthalmus maximus), coho salmon (Oncorhynchus kisutch, and rainbow) Tj ETQqO Ichthyology, 1991, 7, 160-167.	0 0 rgBT / 0.7	Overlock 10 T 26
153	Protection of turbot, Scophthalmus maximus (L.), and rainbow trout, Oncorhynchus mykiss (Richardson), against vibriosis using two different vaccines. Journal of Fish Diseases, 1991, 14, 407-411.	1.9	29
154	Evidence thatYersinia ruckeripossesses a high affinity iron uptake system. FEMS Microbiology Letters, 1991, 80, 121-126.	1.8	29
155	Competitive dominance of antibioticâ€producing marine bacteria in mixed cultures. Journal of Applied Bacteriology, 1991, 71, 228-232.	1.1	34
156	Plasmid associated virulence properties of environmental isolates of Aeromonas hydrophila. Journal of Medical Microbiology, 1991, 35, 264-269.	1.8	21
157	Distribution of plasmid- and chromosome-mediated iron uptake systems in Vibrio anguillarum strains of different origins. Applied and Environmental Microbiology, 1991, 57, 2956-2962.	3.1	44
158	Biochemical and serological characterization of Carnobacterium spp. isolated from farmed and natural populations of striped bass and catfish. Applied and Environmental Microbiology, 1991, 57, 3114-3120.	3.1	69
159	Influence of the Temperature and Cell Line on the Adsorption and Replication of Five Fish Rotaviruses Fish Pathology, 1991, 26, 169-175.	0.7	4
160	Isolation of Serratia plymuthica as an opportunistic pathogen in rainbow trout, Salmo gairdneri Richardson. Journal of Fish Diseases, 1990, 13, 175-177.	1.9	15
161	Changes in bacterial populations during red tides caused by <i>Mesodinium rubrum</i> and <i>Cymnodinium catenatum</i> in North West Coast of Spain. Journal of Applied Bacteriology, 1990, 68, 123-132.	1.1	28
162	Plasmids and factors associated with virulence in environmental isolates of Vibrio cholerae non-O 1 in Bangladesh. Journal of Medical Microbiology, 1990, 33, 107-114.	1.8	18

#	Article	IF	CITATIONS
163	Serological Relationships amongVibrio anguillarumStrains. Journal of Aquatic Animal Health, 1990, 2, 21-29.	1.4	27
164	RELATIONSHIP AMONG PATHOGENIC VIBRIO ANGUILLARUM AND VIBRIO TUBIASHII WITH ENVIRONMENTAL VIBRIOS11In memory of J. Bolinches, who died as a result of a car accident during the elaboration of the present work , 1990, , 77-89.		11
165	ASSOCIATION OF A MORAXELLA SP. AND A REO-LIKE VIRUS WITH MORTALITIES OF STRIPED BASS, MORONE SAXATILIS. , 1990, , 91-99.		18
166	COMPARISON OF THE CELL SURFACE HYDROPHOBICITY OF BACTERIAL FISH PATHOGENS BY DIFFERENT PROCEDURES. , 1990, , 101-115.		13
167	ADHESIVE PROPERTIES AND OTHER VIRULENCE FACTORS IN YERSINIA RUCKERI. , 1990, , 123-139.		8
168	Surface and virulence properties of environmental Vibrio cholerae non-O1 from Albufera Lake (Valencia, Spain). Applied and Environmental Microbiology, 1990, 56, 1140-1147.	3.1	18
169	Visceral neoplasia and epidermal papillomas in cultured turbot Scophthalmus maximus. Diseases of Aquatic Organisms, 1990, 8, 179-187.	1.0	20
170	A review of the taxonomy and seroepizootiology of Vibrio anguillarum, with special reference to aquaculture in the northwest of Spain. Diseases of Aquatic Organisms, 1990, 9, 73-82.	1.0	131
171	Purification and characterization of an antibacterial substance produced by a marine Alteromonas species. Antimicrobial Agents and Chemotherapy, 1989, 33, 1674-1679.	3.2	30
172	Influence of the growth conditions on the hydrophobicity ofRenibacterium salmoninarumevaluated by different methods. FEMS Microbiology Letters, 1989, 60, 71-78.	1.8	15
173	Ulcerative skin and fin lesions in reared turbot, Scophthalmus maximus (L.). Journal of Fish Diseases, 1989, 12, 323-333.	1.9	39
174	Association of Aeromonas sobria with mortalities of adult gizzard shad, Dorosoma cepedianum Lesueur. Journal of Fish Diseases, 1989, 12, 439-448.	1.9	32
175	Inhibitory activity of antibioticâ€producing marine bacteria against fish pathogens. Journal of Applied Bacteriology, 1988, 65, 97-101.	1.1	146
176	Evaluation of selective media for isolation and enumeration of vibrios from estuarine waters. Journal of Microbiological Methods, 1988, 8, 151-160.	1.6	56
177	Chromosome-mediated iron uptake system in pathogenic strains of Vibrio anguillarum. Journal of Bacteriology, 1988, 170, 1920-1925.	2.2	120
178	Virulence properties and enterotoxin production of Aeromonas strains isolated from fish. Infection and Immunity, 1988, 56, 3285-3293.	2.2	137
179	Relationships among virulence for fish, enterotoxigenicity, and phenotypic characteristics of motile Aeromonas. Aquaculture, 1987, 67, 29-39.	3.5	29
180	Homology of Vibrio anguillarum strains causing epizootics in turbot, salmon and trout reared on the Atlantic coast of Spain. Aquaculture, 1987, 67, 41-52.	3.5	77

ALICIA E TORANZO

#	Article	IF	CITATIONS
181	Bacillary necrosis in hatcheries of Ostrea edulis in Spain. Aquaculture, 1987, 65, 15-29.	3.5	56
182	Specificity of slide agglutination test for detecting bacterial fish pathogens. Aquaculture, 1987, 61, 81-97.	3.5	90
183	Evaluation of Different Assay Systems for Identification of Environmental <i>Aeromonas</i> Strains. Applied and Environmental Microbiology, 1986, 51, 652-656.	3.1	32
184	Antibiotic activity of epiphytic bacteria isolated from intertidal seaweeds. Microbial Ecology, 1985, 11, 149-163.	2.8	194
185	Modified Medium for the Oxidation-Fermentation Test in the Identification of Marine Bacteria. Applied and Environmental Microbiology, 1985, 49, 1541-1543.	3.1	117
186	Relation of water temperature to infection of Salmo gairdneri with motile Aeromonas Fish Pathology, 1985, 20, 99-105.	0.7	17
187	Comparison between the bacterial flora associated with fingerling rainbow trout cultured in two different hatcheries in the North-West of Spain. Aquaculture, 1984, 42, 193-206.	3.5	72
188	Plasmid coding for transferable drug resistance in bacteria isolated from cultured rainbow trout. Applied and Environmental Microbiology, 1984, 48, 872-877.	3.1	53
189	Haemagglutinating, haemolytic and cytotoxic activities ofVibrio anguillarumand related vibrios isolated from striped bass on the Atlantic Coast. FEMS Microbiology Letters, 1983, 18, 257-262.	1.8	71
190	Mechanism of poliovirus inactivation by cell-free filtrates of marine bacteria. Canadian Journal of Microbiology, 1983, 29, 1481-1486.	1.7	27
191	Antiviral activity of antibiotic-producing marine bacteria. Canadian Journal of Microbiology, 1982, 28, 231-238.	1.7	36
192	Comparative stability of two salmonid viruses and poliovirus in fresh, estuarine and marine waters. Journal of Fish Diseases, 1982, 5, 223-231.	1.9	58