

# Alicia E Toranzo

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6359204/publications.pdf>

Version: 2024-02-01

191  
papers

7,701  
citations

47409

49  
h-index

87275

74  
g-index

192  
all docs

192  
docs citations

192  
times ranked

4336  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Evidence for the facultative intracellular behaviour of the fish pathogen <i>Vibrio ordalii</i> . Journal of Fish Diseases, 2019, 42, 1447-1455.  | 0.9 | 4         |
| 2  | Biofilm development and cell viability: An undervalued mechanism in the persistence of the fish pathogen <i>Tenacibaculum maritimum</i> . Aquaculture, 2019, 511, 734267.   | 1.7 | 18        |
| 3  | The Fish Pathogen <i>Vibrio ordalii</i> Under Iron Deprivation Produces the Siderophore Piscibactin. Microorganisms, 2019, 7, 313.  | 1.6 | 15        |
| 4  | Genetic studies to re-affiliate <i>Edwardsiella tarda</i> fish isolates to <i>Edwardsiella piscicida</i> and <i>Edwardsiella anguillarum</i> species. Systematic and Applied Microbiology, 2018, 41, 30-37.   | 1.2 | 58        |
| 5  | Population genetic and evolution analysis of controversial genus <i>Edwardsiella</i> by multilocus sequence typing. Molecular Phylogenetics and Evolution, 2018, 127, 513-521.  | 1.2 | 11        |
| 6  | <i>Edwardsiella piscicida</i> : a significant bacterial pathogen of cultured fish. Diseases of Aquatic Organisms, 2018, 131, 59-71.   | 0.5 | 71        |
| 7  | Draft Genome Sequence of <i>Edwardsiella piscicida</i> Strain ACC35.1 Isolated from Diseased Turbot ( <i>Scophthalmus maximus</i> ) in Europe. Genome Announcements, 2017, 5, .   | 0.8 | 4         |
| 8  | Insights into the virulence-related genes of <i>Edwardsiella tarda</i> isolated from turbot in Europe: genetic homogeneity and evidence for vibrioferrin production. Journal of Fish Diseases, 2016, 39, 565-576.   | 0.9 | 11        |
| 9  | Iron assimilation and siderophore production by <i>Vibrio ordalii</i> strains isolated from diseased Atlantic salmon <i>Salmo salar</i> in Chile. Diseases of Aquatic Organisms, 2016, 118, 217-226.  | 0.5 | 9         |
| 10 | Survival behaviour and virulence of the fish pathogen <i>Vibrio ordalii</i> in seawater microcosms. Diseases of Aquatic Organisms, 2016, 120, 27-38.  | 0.5 | 6         |
| 11 | Cell-surface properties of <i>Vibrio ordalii</i> strains isolated from Atlantic salmon <i>Salmo salar</i> in Chilean farms. Diseases of Aquatic Organisms, 2015, 113, 9-23.   | 0.5 | 16        |
| 12 | First Isolation and Characterization of <i>Tenacibaculum soleae</i> from Sea Bass <i>Dicentrarchus labrax</i> . Fish Pathology, 2014, 49, 16-22.  | 0.4 | 5         |
| 13 | <i>Vibrio ostreicida</i> sp. nov., a new pathogen of bivalve larvae. International Journal of Systematic and Evolutionary Microbiology, 2014, 64, 1641-1646.  | 0.8 | 26        |
| 14 | Multilocus Sequence Analysis of the Marine Bacterial Genus <i>Tenacibaculum</i> Suggests Parallel Evolution of Fish Pathogenicity and Endemic Colonization of Aquaculture Systems. Applied and Environmental Microbiology, 2014, 80, 5503-5514.   | 1.4 | 69        |
| 15 | Comparative polyphasic characterization of <i>Streptococcus phocae</i> strains with different host origin and description of the subspecies <i>Streptococcus phocae</i> subsp. <i>salmonis</i> subsp. nov.. International Journal of Systematic and Evolutionary Microbiology, 2014, 64, 1775-1781. | 0.8 | 19        |
| 16 | A multiplex PCR for the simultaneous detection of <i>Tenacibaculum maritimum</i> and <i>Edwardsiella tarda</i> in aquaculture. International Microbiology, 2014, 17, 111-7.   | 1.1 | 10        |
| 17 | <i>Vibrio ordalii</i> antimicrobial susceptibility testing—Modified culture conditions required and laboratory-specific epidemiological cut-off values. Veterinary Microbiology, 2013, 165, 434-442.  | 0.8 | 11        |
| 18 | <i>Pseudomonas baetica</i> sp. nov., a fish pathogen isolated from wedge sole, <i>Dicologlossa cuneata</i> (Moreau). International Journal of Systematic and Evolutionary Microbiology, 2012, 62, 874-882.  | 0.8 | 56        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | Iron utilization and siderophore production by <i>Streptococcus phocae</i> isolated from diseased Atlantic salmon ( <i>Salmo salar</i> ). <i>Aquaculture</i> , 2012, 364-365, 305-311.   | 1.7 | 8         |
| 20 | First description of <i>Edwardsiella tarda</i> in Senegalese sole, <i>Solea senegalensis</i> (Kaup). <i>Journal of Fish Diseases</i> , 2012, 35, 79-82.  | 0.9 | 12        |
| 21 | Estimation of epidemiological cut-off values for disk diffusion susceptibility test data for <i>Streptococcus phocae</i> . <i>Aquaculture</i> , 2011, 314, 44-48.  | 1.7 | 13        |
| 22 | Characterization of <i>Vibrio tapetis</i> strains isolated from diseased cultured Wedge sole ( <i>Dicologlossa cuneata</i> Moreau). <i>Research in Veterinary Science</i> , 2011, 90, 189-195.   | 0.9 | 15        |
| 23 | Intraspecific genetic variability of <i>Edwardsiella tarda</i> strains from cultured turbot. <i>Diseases of Aquatic Organisms</i> , 2011, 95, 253-258.   | 0.5 | 19        |
| 24 | Pathogenic Potential of <i>Edwardsiella tarda</i> Strains Isolated from Turbot. <i>Fish Pathology</i> , 2011, 46, 27-30.   | 0.4 | 5         |
| 25 | Serological and molecular heterogeneity among <i>Yersinia ruckeri</i> strains isolated from farmed Atlantic salmon <i>Salmo salar</i> in Chile. <i>Diseases of Aquatic Organisms</i> , 2011, 93, 207-214.  | 0.5 | 33        |
| 26 | Surface properties of <i>Streptococcus phocae</i> strains isolated from diseased Atlantic salmon, <i>Salmo salar</i> L.. <i>Journal of Fish Diseases</i> , 2011, 34, 203-215.  | 0.9 | 30        |
| 27 | Pseudo-membranes on internal organs associated with <i>Rhodococcus qingshengii</i> infection in Atlantic salmon ( <i>Salmo salar</i> ). <i>Veterinary Microbiology</i> , 2011, 147, 200-204.   | 0.8 | 8         |
| 28 | Multiplex PCR for the detection of <i>Piscirickettsia salmonis</i> , <i>Vibrio anguillarum</i> , <i>Aeromonas salmonicida</i> and <i>Streptococcus phocae</i> in Chilean marine farms. <i>Diseases of Aquatic Organisms</i> , 2011, 97, 135-142. | 0.5 | 22        |
| 29 | Contact Lens Hydrophobicity and Roughness Effects on Bacterial Adhesion. <i>Optometry and Vision Science</i> , 2010, 87, E426-E431.  | 0.6 | 64        |
| 30 | Evaluation of four polymerase chain reaction primer pairs for the detection of <i>Edwardsiella tarda</i> in turbot. <i>Diseases of Aquatic Organisms</i> , 2010, 90, 55-61.  | 0.5 | 29        |
| 31 | <i>Vibrio gallaecicus</i> sp. nov. isolated from cultured clams in north-western Spain. <i>Systematic and Applied Microbiology</i> , 2009, 32, 111-117.  | 1.2 | 33        |
| 32 | Genetic characterization of <i>Streptococcus phocae</i> strains isolated from Atlantic salmon, <i>Salmo salar</i> L., in Chile. <i>Journal of Fish Diseases</i> , 2009, 32, 351-358.   | 0.9 | 18        |
| 33 | First isolation of <i>Tenacibaculum maritimum</i> from wedge sole, <i>Dicologlossa cuneata</i> (Moreau). <i>Journal of Fish Diseases</i> , 2009, 32, 603-610.  | 0.9 | 15        |
| 34 | Identification of <i>Vibrio harveyi</i> isolated from diseased cultured wedge sole <i>Dicologlossa cuneata</i> . <i>Diseases of Aquatic Organisms</i> , 2009, 84, 209-217.   | 0.5 | 19        |
| 35 | Vaccination Strategies to Prevent Streptococcal Infections in Cultured Fish., 2009, , 111-149.   |     | 1         |
| 36 | Evolution of drug resistance and minimum inhibitory concentration to enrofloxacin in <i>Tenacibaculum maritimum</i> strains isolated in fish farms. <i>Aquaculture International</i> , 2008, 16, 1-11.   | 1.1 | 23        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 37 | First description of serotype O3 in <i>Vibrio anguillarum</i> strains isolated from salmonids in Chile. <i>Journal of Fish Diseases</i> , 2008, 31, 235-239.  | 0.9 | 30        |
| 38 | <i>Streptococcus phocae</i> , an emerging pathogen for salmonid culture. <i>Veterinary Microbiology</i> , 2008, 130, 198-207.   | 0.8 | 56        |
| 39 | Antigenic and molecular characterization of <i>Vibrio ordalii</i> strains isolated from Atlantic salmon <i>Salmo salar</i> in Chile. <i>Diseases of Aquatic Organisms</i> , 2008, 79, 27-35.  | 0.5 | 21        |
| 40 | Development of an effective <i>Edwardsiella tarda</i> vaccine for cultured turbot ( <i>Scophthalmus maximus</i> ). <i>Fish and Shellfish Immunology</i> , 2008, 25, 208-212.  | 1.6 | 59        |
| 41 | Polymerase Chain Reaction Amplification of Repetitive Intergenic Consensus and Repetitive Extragenic Palindromic Sequences for Molecular Typing of <i>Pseudomonas anguilliseptica</i> and <i>Aeromonas salmonicida</i> . <i>Journal of Aquatic Animal Health</i> , 2008, 20, 75-85. | 0.6 | 14        |
| 42 | Evaluation of different DNA-based fingerprinting methods for typing <i>Photobacterium damsela</i> ssp. <i>piscicida</i> . <i>Biological Research</i> , 2007, 40, 85-92.   | 1.5 | 13        |
| 43 | Use of adjuvanted vaccines to lengthen the protection against lactococcosis in rainbow trout ( <i>Oncorhynchus mykiss</i> ). <i>Aquaculture</i> , 2006, 251, 153-158.   | 1.7 | 37        |
| 44 | Use of hydrogen peroxide against the fish pathogen <i>Tenacibaculum maritimum</i> and its effect on infected turbot ( <i>Scophthalmus maximus</i> ). <i>Aquaculture</i> , 2006, 257, 104-110.   | 1.7 | 60        |
| 45 | <i>Tenacibaculosis</i> infection in marine fish caused by <i>Tenacibaculum maritimum</i> : a review. <i>Diseases of Aquatic Organisms</i> , 2006, 71, 255-266.  | 0.5 | 215       |
| 46 | Use of microcosms to determine the survival of the fish pathogen <i>Tenacibaculum maritimum</i> in seawater. <i>Environmental Microbiology</i> , 2006, 8, 921-928.  | 1.8 | 26        |
| 47 | A challenge model for <i>Tenacibaculum maritimum</i> infection in turbot, <i>Scophthalmus maximus</i> (L.). <i>Journal of Fish Diseases</i> , 2006, 29, 371-374.  | 0.9 | 43        |
| 48 | Characterization of <i>Edwardsiella tarda</i> strains isolated from turbot, <i>Psetta maxima</i> (L.). <i>Journal of Fish Diseases</i> , 2006, 29, 541-547.   | 0.9 | 103       |
| 49 | Effectiveness of a divalent vaccine for sole, <i>Solea senegalensis</i> (Kaup), against <i>Vibrio harveyi</i> and <i>Photobacterium damsela</i> subsp. <i>piscicida</i> . <i>Journal of Fish Diseases</i> , 2005, 28, 33-38.  | 0.9 | 60        |
| 50 | Recommendation of an Appropriate Medium for In Vitro Drug Susceptibility Testing of the Fish Pathogen <i>Tenacibaculum maritimum</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2005, 49, 82-87.  | 1.4 | 15        |
| 51 | Iron Uptake Mechanisms in the Fish Pathogen <i>Tenacibaculum maritimum</i> . <i>Applied and Environmental Microbiology</i> , 2005, 71, 6947-6953.   | 1.4 | 34        |
| 52 | Variation in 16S-23S rRNA Intergenic Spacer Regions in <i>Photobacterium damsela</i> : a Mosaic-Like Structure. <i>Applied and Environmental Microbiology</i> , 2005, 71, 636-645.  | 1.4 | 41        |
| 53 | A review of the main bacterial fish diseases in mariculture systems. <i>Aquaculture</i> , 2005, 246, 37-61.   | 1.7 | 671       |
| 54 | Development and Validation of a PCR-based Protocol for the Detection of <i>Pseudomonas anguilliseptica</i> . <i>Fish Pathology</i> , 2004, 39, 33-41.   | 0.4 | 13        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 55 | Intraspecific diversity of the marine fish pathogen <i>Tenacibaculum maritimum</i> as determined by randomly amplified polymorphic DNA-PCR. <i>Journal of Applied Microbiology</i> , 2004, 96, 871-877.  | 1.4 | 29        |
| 56 | Oral immunization using alginate microparticles as a useful strategy for booster vaccination against fish lactococcosis. <i>Aquaculture</i> , 2004, 236, 119-129.  | 1.7 | 86        |
| 57 | Species-specific polymerase chain reaction primer sets for the diagnosis of <i>Tenacibaculum maritimum</i> infection. <i>Diseases of Aquatic Organisms</i> , 2004, 62, 75-83.  | 0.5 | 45        |
| 58 | Phenotypic characterization and description of two major O-serotypes in <i>Tenacibaculum maritimum</i> strains from marine fishes. <i>Diseases of Aquatic Organisms</i> , 2004, 58, 1-8.   | 0.5 | 39        |
| 59 | Characterization of the 23S and 5S rRNA genes and 23S-5S intergenic spacer region (ITS-2) of <i>Photobacterium damsela</i> . <i>Diseases of Aquatic Organisms</i> , 2004, 61, 33-39.   | 0.5 | 5         |
| 60 | Isolation of mutants of <i>Vibrio anguillarum</i> defective in haeme utilisation and cloning of <i>huvA</i> , a gene coding for an outer membrane protein involved in the use of haeme as iron source. <i>Archives of Microbiology</i> , 2003, 179, 329-338. | 1.0 | 26        |
| 61 | Existence of two O-serotypes in the fish pathogen <i>Pseudomonas anguilliseptica</i> . <i>Veterinary Microbiology</i> , 2003, 94, 325-333.   | 0.8 | 16        |
| 62 | Isolation of a highly pathogenic <i>Vibrio pelagius</i> strain associated with mass mortalities of turbot, <i>Scophthalmus maximus</i> (L.), larvae. <i>Journal of Fish Diseases</i> , 2003, 26, 293-303.  | 0.9 | 44        |
| 63 | Phenotypic and Genetic Characterization of <i>Pseudomonas anguilliseptica</i> Strains Isolated from Fish. <i>Journal of Aquatic Animal Health</i> , 2003, 15, 39-47.   | 0.6 | 26        |
| 64 | Molecular Fingerprinting of Fish-Pathogenic <i>Lactococcus garvieae</i> Strains by Random Amplified Polymorphic DNA Analysis. <i>Journal of Clinical Microbiology</i> , 2003, 41, 751-756.   | 1.8 | 62        |
| 65 | <i>Lactococcus garvieae</i> in wild Red Sea wrasse <i>Coris aygula</i> (Labridae). <i>Diseases of Aquatic Organisms</i> , 2003, 56, 275-278.   | 0.5 | 30        |
| 66 | Dormancy as a survival strategy of the fish pathogen <i>Streptococcus parauberis</i> in the marine environment. <i>Diseases of Aquatic Organisms</i> , 2002, 52, 129-136.  | 0.5 | 28        |
| 67 | Molecular Approaches for the Study and Diagnosis of Salmonid Streptococcosis. <i>Reviews: Methods and Technologies in Fish Biology and Fisheries</i> , 2002, , 211-233.  | 0.6 | 15        |
| 68 | Binding of haemin by the fish pathogen <i>Photobacterium damsela</i> subsp. <i>piscicida</i> . <i>Diseases of Aquatic Organisms</i> , 2002, 48, 109-115.   | 0.5 | 23        |
| 69 | Effect of temperature on the development of pasteurellosis in carrier gilthead seabream ( <i>Sparus</i> ) Tj ETQq1 1 0.784314 rgBT /Overlock 1   | 1.7 | 34        |
| 70 | Molecular characterization of Portuguese strains of <i>Yersinia ruckeri</i> isolated from fish culture systems. <i>Journal of Fish Diseases</i> , 2001, 24, 151-159.   | 0.9 | 14        |
| 71 | Multiplex PCR assay for <i>ureC</i> and 16S rRNA genes clearly discriminates between both subspecies of <i>Photobacterium damsela</i> . <i>Diseases of Aquatic Organisms</i> , 2000, 40, 177-183.  | 0.5 | 86        |
| 72 | Existence of two geographically-linked clonal lineages in the bacterial fish pathogen <i>Photobacterium damsela</i> subsp. <i>piscicida</i> evidenced by random amplified polymorphic DNA analysis. <i>Epidemiology and Infection</i> , 2000, 125, 213-219.  | 1.0 | 35        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 73 | Evidence that water transmits the disease caused by the fish pathogen <i>Photobacterium damsela</i> subsp. <i>damsela</i> . <i>Journal of Applied Microbiology</i> , 2000, 88, 531-535.  | 1.4 | 90        |
| 74 | Presence of phospholipase-D ( <i>dly</i> ) gene coding for <i>damselysin</i> production is not a pre-requisite for pathogenicity in <i>Photobacterium damsela</i> subsp. <i>damsela</i> . <i>Microbial Pathogenesis</i> , 2000, 28, 119-126. | 1.3 | 25        |
| 75 | Electrotransformation of <i>Photobacterium damsela</i> subsp. <i>piscicida</i> . , 2000, , 175-181.  |     | 1         |
| 76 | Electrotransformation of <i>Yersinia ruckeri</i> . , 2000, , 127-133.  |     | 0         |
| 77 | 16S rRNA Gene Sequence Analysis of <i>Photobacterium damsela</i> and Nested PCR Method for Rapid Detection of the Causative Agent of Fish Pasteurellosis. <i>Applied and Environmental Microbiology</i> , 1999, 65, 2942-2946.               | 1.4 | 94        |
| 78 | Genetic analysis of turbot pathogenic <i>Streptococcus parauberis</i> strains by ribotyping and random amplified polymorphic DNA. <i>FEMS Microbiology Letters</i> , 1999, 179, 297-304.   | 0.7 | 38        |
| 79 | Assessment of a magnetic bead-EIA based kit for rapid diagnosis of fish pasteurellosis. <i>Journal of Microbiological Methods</i> , 1999, 38, 147-154.   | 0.7 | 12        |
| 80 | Sanitary control in Spanish aquaculture. <i>Journal of Applied Ichthyology</i> , 1998, 14, 275-277.  | 0.3 | 1         |
| 81 | EDWARDSIELLOSIS IN WILD STRIPED BASS FROM THE CHESAPEAKE BAY. <i>Journal of Wildlife Diseases</i> , 1997, 33, 517-525.   | 0.3 | 30        |
| 82 | Applicability of Ribotyping for Intraspecific Classification and Epidemiological Studies of <i>Photobacterium damsela</i> subsp. <i>piscicida</i> . <i>Systematic and Applied Microbiology</i> , 1997, 20, 634-639.                          | 1.2 | 22        |
| 83 | Viability of starved <i>Pasteurella piscicida</i> in seawater monitored by flow cytometry and the effect of antibiotics on its resuscitation. <i>Letters in Applied Microbiology</i> , 1997, 24, 122-126.                                    | 1.0 | 25        |
| 84 | Phenotypic, serological and molecular characterization of Portuguese isolates of <i>Aeromonas salmonicida</i> subsp. <i>salmonicida</i> . <i>Fish Pathology</i> , 1997, 32, 177-178.   | 0.4 | 2         |
| 85 | Antigenic characterization of <i>Vibrio anguillarum</i> -related organisms isolated from turbot and cod. <i>Diseases of Aquatic Organisms</i> , 1997, 28, 45-50.   | 0.5 | 10        |
| 86 | Association of <i>Aeromonas hydrophila</i> and <i>Vibrio alginolyticus</i> with Larval Mortalities of Scallop ( <i>Argopecten purpuratus</i> ). <i>Journal of Invertebrate Pathology</i> , 1996, 67, 213-218.                                | 1.5 | 67        |
| 87 | Influence of the capsular layer on the virulence of <i>Pasteurella piscicida</i> for fish. <i>Microbial Pathogenesis</i> , 1996, 21, 289-297.  | 1.3 | 62        |
| 88 | Adherence and invasive capacities of the fish pathogen <i>Pasteurella piscicida</i> . <i>FEMS Microbiology Letters</i> , 1996, 138, 29-34.   | 0.7 | 57        |
| 89 | Phenotypic and pathobiological characteristics of <i>Pasteurella piscicida</i> . <i>Annual Review of Fish Diseases</i> , 1996, 6, 41-64.   | 1.1 | 66        |
| 90 | Health status of two salmonid aquaculture facilities in North Portugal: characterization of the bacterial and viral pathogens causing notifiable diseases. <i>Journal of Fish Diseases</i> , 1996, 19, 83-89.                                | 0.9 | 13        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 91  | Evaluation of media for the successful culture of <i>Flexibacter maritimus</i> . <i>Journal of Fish Diseases</i> , 1996, 19, 193-197.   | 0.9 | 103       |
| 92  | Immunological analysis of extracellular products and cell surface components of motile <i>Aeromonas</i> isolated from fish. <i>Journal of Applied Bacteriology</i> , 1996, 81, 585-593.                                       | 1.1 | 6         |
| 93  | Phenotypic and pathobiological characteristics of <i>Pasteurella piscicida</i> . <i>Annual Review of Fish Diseases</i> , 1996, 6, 41-64.  | 1.1 | 52        |
| 94  | Occurrence of an erythrocytic virus infection in cultured turbot <i>Scophthalmus maximus</i> . <i>Diseases of Aquatic Organisms</i> , 1996, 24, 159-167.  | 0.5 | 7         |
| 95  | Biochemical and serological analysis of <i>Vibrio anguillarum</i> related organisms. <i>Diseases of Aquatic Organisms</i> , 1996, 26, 67-73.  | 0.5 | 6         |
| 96  | Pathogenicity studies on a <i>Vibrio anguillarum</i> -related (VAR) strain causing an epizootic in <i>Argopecten purpuratus</i> larvae cultured in Chile. <i>Diseases of Aquatic Organisms</i> , 1995, 22, 135-141.           | 0.5 | 82        |
| 97  | In vitro killing of <i>Pasteurella piscicida</i> by fish macrophages. <i>Diseases of Aquatic Organisms</i> , 1995, 23, 51-57.   | 0.5 | 38        |
| 98  | Pathogenicity of live bacteria and extracellular products of motile <i>Aeromonas</i> isolated from eels. <i>Journal of Applied Bacteriology</i> , 1995, 78, 555-562.  | 1.1 | 34        |
| 99  | Pathology associated with a viral erythrocytic infection in turbot, <i>Scophthalmus maximus</i> (L.). <i>Journal of Fish Diseases</i> , 1995, 18, 425-433.  | 0.9 | 3         |
| 100 | Pathology of <i>Enterococcus</i> sp. infection in farmed turbot, <i>Scophthalmus maximus</i> L.. <i>Journal of Fish Diseases</i> , 1995, 18, 21-30.   | 0.9 | 28        |
| 101 | Histopathological lesions caused by <i>Vibrio damsela</i> in cultured turbot, <i>Scophthalmus maximus</i> (L.): inoculations with live cells and extracellular products. <i>Journal of Fish Diseases</i> , 1995, 18, 357-364. | 0.9 | 23        |
| 102 | Genetic transformation of <i>Vibrio anguillarum</i> and <i>Pasteurella piscicida</i> by electroporation. <i>FEMS Microbiology Letters</i> , 1995, 128, 75-80.   | 0.7 | 8         |
| 103 | Pathogenicity of birnaviruses isolated from turbot ( <i>Scophthalmus maximus</i> ): comparison with reference serotypes of IPNV. <i>Aquaculture</i> , 1995, 130, 7-14.  | 1.7 | 12        |
| 104 | Efficacy of intraperitoneal and immersion vaccination against <i>Enterococcus</i> sp. infection in turbot. <i>Aquaculture</i> , 1995, 134, 17-27.   | 1.7 | 61        |
| 105 | Fatty acid profiles of <i>Pasteurella</i> <i>piscicida</i> : comparison with other fish pathogenic gram-negative bacteria. <i>Archives of Microbiology</i> , 1995, 163, 211-216.  | 1.0 | 11        |
| 106 | Response of <i>Pasteurella piscicida</i> and <i>Flexibacter maritimus</i> to skin mucus of marine fish. <i>Diseases of Aquatic Organisms</i> , 1995, 21, 103-108.   | 0.5 | 85        |
| 107 | Fatty acid profiles of "Pasteurella" <i>piscicida</i> : comparison with other fish pathogenic gram-negative bacteria. <i>Archives of Microbiology</i> , 1995, 163, 211-216.   | 1.0 | 12        |
| 108 | Analysis of antigens present in the extracellular products and cell surface of <i>Vibrio anguillarum</i> serotypes O1, O2, and O3. <i>Applied and Environmental Microbiology</i> , 1995, 61, 2493-2498.                       | 1.4 | 33        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 109 | Evaluation of BIONOR Mono-kits for rapid detection of bacterial fish pathogens. <i>Diseases of Aquatic Organisms</i> , 1995, 21, 25-34.   | 0.5 | 31        |
| 110 | Sequential pathology of experimental pasteurellosis in gilthead seabream <i>Sparus aurata</i> . A light- and electron-microscopic study. <i>Diseases of Aquatic Organisms</i> , 1995, 21, 177-186.  | 0.5 | 47        |
| 111 | Antigenic characterization of <i>Enterococcus</i> strains pathogenic for turbot and their relationship with other Gram-positive bacteria. <i>Diseases of Aquatic Organisms</i> , 1995, 21, 187-191.   | 0.5 | 22        |
| 112 | A Comparison of Pathological Changes Caused by <i>Vibrio anguillarum</i> and Its Extracellular Products in Rainbow Trout ( <i>Oncorhynchus mykiss</i> ).. <i>Fish Pathology</i> , 1994, 29, 79-89.  | 0.4 | 29        |
| 113 | O-Serogrouping and surface components of <i>Aeromonas hydrophila</i> and <i>Aeromonas jandaei</i> pathogenic for eels. <i>FEMS Microbiology Letters</i> , 1994, 117, 85-90.   | 0.7 | 39        |
| 114 | Capsular polysaccharide expressed by <i>Pasteurella piscicida</i> grown in vitro. <i>FEMS Microbiology Letters</i> , 1994, 124, 285-289.  | 0.7 | 32        |
| 115 | Non-specific cellular responses of rainbow trout to <i>Vibrio anguillarum</i> and its extracellular products (ECPs). <i>Journal of Fish Biology</i> , 1994, 45, 839-854.  | 0.7 | 45        |
| 116 | Incidence of <i>Yersinia ruckeri</i> in two farms in Galicia (NW Spain) during a one-year period. <i>Journal of Fish Diseases</i> , 1994, 17, 533-539.  | 0.9 | 7         |
| 117 | Starvation-Survival Processes of the Bacterial Fish Pathogen <i>Yersinia ruckeri</i> . <i>Systematic and Applied Microbiology</i> , 1994, 17, 161-168.  | 1.2 | 29        |
| 118 | Kinetics of antibody production against <i>Vibrio anguillarum</i> antigens in turbot. <i>Aquaculture</i> , 1994, 123, 191-196.  | 1.7 | 13        |
| 119 | Role of serum antibodies in protection of vaccinated turbot ( <i>Scophthalmus maximus</i> ) against vibriosis. <i>Aquaculture</i> , 1994, 123, 197-204.   | 1.7 | 13        |
| 120 | Vaccination trials on gilthead seabream ( <i>Sparus aurata</i> ) against <i>Pasteurella piscicida</i> . <i>Aquaculture</i> , 1994, 120, 201-208.  | 1.7 | 50        |
| 121 | Role of iron, capsule, and toxins in the pathogenicity of <i>Vibrio vulnificus</i> biotype 2 for mice. <i>Infection and Immunity</i> , 1994, 62, 759-763.   | 1.0 | 102       |
| 122 | Histopathology of rainbow trout, <i>Oncorhynchus mykiss</i> (Walbaum), and striped bass, <i>Morone saxatilis</i> (Walbaum), experimentally infected with <i>Carnobacterium piscicola</i> . <i>Journal of Fish Diseases</i> , 1993, 16, 261-267. | 0.9 | 12        |
| 123 | Pathological activities of <i>Yersinia ruckeri</i> , the Enteric Redmouth (ERM) bacterium. <i>FEMS Microbiology Letters</i> , 1993, 112, 291-300.   | 0.7 | 62        |
| 124 | Toxicity of the extracellular products of <i>Vibrio damsela</i> isolated from diseased fish. <i>Current Microbiology</i> , 1993, 27, 341-347.   | 1.0 | 48        |
| 125 | Usefulness of the API-20E system for the identification of bacterial fish pathogens. <i>Aquaculture</i> , 1993, 116, 111-120.   | 1.7 | 52        |
| 126 | Microflora associated with healthy and diseased turbot ( <i>Scophthalmus maximus</i> ) from three farms in northwest Spain. <i>Aquaculture</i> , 1993, 114, 189-202.  | 1.7 | 55        |



| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 127 | Virulence factors of bacteria pathogenic for coldwater fish. Annual Review of Fish Diseases, 1993, 3, 5-36.   | 1.1 | 73        |
| 128 | Antigenic and Molecular Characterization of <i>Yersinia ruckeri</i> Proposal for a New Intraspecies Classification. Systematic and Applied Microbiology, 1993, 16, 411-419.   | 1.2 | 70        |
| 129 | Detection of a Common Antigen among <i>Renibacterium salmoninarum</i> , <i>Corynebacterium aquaticum</i> , and <i>Carnobacterium piscicola</i> by the Western Blot Technique. Journal of Aquatic Animal Health, 1993, 5, 172-176. | 0.6 | 8         |
| 130 | Phenotypic Characteristics and Virulence of <i>Vibrio anguillarum</i> -Related Organisms. Applied and Environmental Microbiology, 1993, 59, 2969-2976.  | 1.4 | 38        |
| 131 | <i>Vibrio mimicus</i> and <i>Vibrio cholerae</i> non-O1 Isolated from Wild and Hatchery-Reared Fish.. Fish Pathology, 1993, 28, 15-26.  | 0.4 | 4         |
| 132 | Characterization of a birnavirus isolated from diseased turbot cultured in Spain. Diseases of Aquatic Organisms, 1993, 15, 163-163.   | 0.5 | 25        |
| 133 | Isolation of IPN virus serotype VR-299 from turbot in Europe. Diseases of Aquatic Organisms, 1993, 17, 61-65.   | 0.5 | 11        |
| 134 | An epizootic in farmed, market-sized rainbow trout in Spain caused by a strain of <i>Carnobacterium piscicola</i> of unusual virulence. Diseases of Aquatic Organisms, 1993, 17, 87-99.   | 0.5 | 37        |
| 135 | Detection of a vascular permeability factor in the extracellular products of <i>Renibacterium salmoninarum</i> . Microbial Pathogenesis, 1992, 13, 237-241.   | 1.3 | 2         |
| 136 | Epizootic study in a turbot farm: bacteriology, virology, parasitology and histology. Aquaculture, 1992, 107, 253-258.  | 1.7 | 14        |
| 137 | Comparison of the extracellular biological activities of <i>Vibrio anguillarum</i> and <i>Aeromonas hydrophila</i> . Aquaculture, 1992, 107, 259-270.   | 1.7 | 22        |
| 138 | The detection of two antigenic groups among <i>Renibacterium salmoninarum</i> isolates. FEMS Microbiology Letters, 1992, 94, 105-110.   | 0.7 | 2         |
| 139 | <i>Serratia marcescens</i> : a potential pathogen for fish. Journal of Fish Diseases, 1992, 15, 15-26.  | 0.9 | 42        |
| 140 | Antigenic relationships among rotaviruses isolated from fish. Journal of Fish Diseases, 1992, 15, 27-36.  | 0.9 | 16        |
| 141 | Presence of skin permeability factors in the extracellular products of <i>Yersinia ruckeri</i> . Current Microbiology, 1992, 24, 263-267.   | 1.0 | 8         |
| 142 | Comparative study of phenotypic and virulence properties in <i>Vibrio vulnificus</i> biotypes 1 and 2 obtained from a European eel farm experiencing mortalities. Diseases of Aquatic Organisms, 1992, 13, 29-35.                 | 0.5 | 78        |
| 143 | Phenotypic and pathobiological properties of <i>Corynebacterium aquaticum</i> isolated from diseased striped bass. Diseases of Aquatic Organisms, 1992, 14, 115-126.  | 0.5 | 26        |
| 144 | <i>Tetramicra brevifilum</i> , a potential threat to farmed turbot <i>Scophthalmus maximus</i> . Diseases of Aquatic Organisms, 1992, 14, 127-135.  | 0.5 | 30        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 145 | Characterization of strains related to brown ring disease outbreaks in southwestern Spain. <i>Diseases of Aquatic Organisms</i> , 1992, 14, 229-236.  | 0.5 | 35        |
| 146 | MICs and MBCs of chemotherapeutic agents against <i>Renibacterium salmoninarum</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 1991, 35, 1011-1013.  | 1.4 | 25        |
| 147 | Lack of Biological Activities in the Extracellular Products of <i>Renibacterium salmoninarum</i> . <i>Canadian Journal of Fisheries and Aquatic Sciences</i> , 1991, 48, 421-425.   | 0.7 | 23        |
| 148 | Pasteurellosis in cultured gilthead seabream ( <i>Sparus aurata</i> ): first report in Spain. <i>Aquaculture</i> , 1991, 99, 1-15.  | 1.7 | 143       |
| 149 | Biochemical and Serological Characteristics, Drug Resistance and Plasmid Profiles of Spanish Isolates of <i>Aeromonas salmonicida</i> . <i>Fish Pathology</i> , 1991, 26, 55-60.  | 0.4 | 23        |
| 150 | Cell-Surface-Associated Properties of Fish Pathogenic Bacteria. <i>Journal of Aquatic Animal Health</i> , 1991, 3, 297-301.   | 0.6 | 30        |
| 151 | Susceptibility of turbot ( <i>Scophthalmus maximus</i> ), coho salmon ( <i>Oncorhynchus kisutch</i> , and rainbow) Tj ETQq1 1 0.784314 rgBT /Ove<br>lchthyology, 1991, 7, 160-167.  | 0.3 | 26        |
| 152 | Protection of turbot, <i>Scophthalmus maximus</i> (L.), and rainbow trout, <i>Oncorhynchus mykiss</i> (Richardson), against vibriosis using two different vaccines. <i>Journal of Fish Diseases</i> , 1991, 14, 407-411.  | 0.9 | 29        |
| 153 | Evidence that <i>Yersinia ruckeri</i> possesses a high affinity iron uptake system. <i>FEMS Microbiology Letters</i> , 1991, 80, 121-126.   | 0.7 | 29        |
| 154 | Competitive dominance of antibiotic-producing marine bacteria in mixed cultures. <i>Journal of Applied Bacteriology</i> , 1991, 71, 228-232.  | 1.1 | 34        |
| 155 | Plasmid associated virulence properties of environmental isolates of <i>Aeromonas hydrophila</i> . <i>Journal of Medical Microbiology</i> , 1991, 35, 264-269.  | 0.7 | 21        |
| 156 | Distribution of plasmid- and chromosome-mediated iron uptake systems in <i>Vibrio anguillarum</i> strains of different origins. <i>Applied and Environmental Microbiology</i> , 1991, 57, 2956-2962.                      | 1.4 | 44        |
| 157 | Biochemical and serological characterization of <i>Carnobacterium</i> spp. isolated from farmed and natural populations of striped bass and catfish. <i>Applied and Environmental Microbiology</i> , 1991, 57, 3114-3120. | 1.4 | 69        |
| 158 | Influence of the Temperature and Cell Line on the Adsorption and Replication of Five Fish Rotaviruses.. <i>Fish Pathology</i> , 1991, 26, 169-175.  | 0.4 | 4         |
| 159 | Isolation of <i>Serratia plymuthica</i> as an opportunistic pathogen in rainbow trout, <i>Salmo gairdneri</i> Richardson. <i>Journal of Fish Diseases</i> , 1990, 13, 175-177.  | 0.9 | 15        |
| 160 | Changes in bacterial populations during red tides caused by <i>Mesodinium rubrum</i> and <i>Gymnodinium catenatum</i> in North West Coast of Spain. <i>Journal of Applied Bacteriology</i> , 1990, 68, 123-132.           | 1.1 | 28        |
| 161 | Plasmids and factors associated with virulence in environmental isolates of <i>Vibrio cholerae</i> non-O 1 in Bangladesh. <i>Journal of Medical Microbiology</i> , 1990, 33, 107-114.                                     | 0.7 | 18        |
| 162 | Serological Relationships among <i>Vibrio anguillarum</i> Strains. <i>Journal of Aquatic Animal Health</i> , 1990, 2, 21-29.  | 0.6 | 27        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 163 | RELATIONSHIP AMONG PATHOGENIC VIBRIO ANGUILLARUM AND VIBRIO TUBIASHII WITH ENVIRONMENTAL VIBRIOS11In memory of J. Bolinches, who died as a result of a car accident during the elaboration of the present work.. , 1990, , 77-89. |     | 11        |
| 164 | ASSOCIATION OF A MORAXELLA SP. AND A REO-LIKE VIRUS WITH MORTALITIES OF STRIPED BASS, MORONE SAXATILIS. , 1990, , 91-99.  |     | 18        |
| 165 | COMPARISON OF THE CELL SURFACE HYDROPHOBICITY OF BACTERIAL FISH PATHOGENS BY DIFFERENT PROCEDURES. , 1990, , 101-115.   |     | 13        |
| 166 | ADHESIVE PROPERTIES AND OTHER VIRULENCE FACTORS IN YERSINIA RUCKERI. , 1990, , 123-139.   |     | 8         |
| 167 | Surface and virulence properties of environmental <i>Vibrio cholerae</i> non-O1 from Albufera Lake (Valencia, Spain). <i>Applied and Environmental Microbiology</i> , 1990, 56, 1140-1147.  | 1.4 | 18        |
| 168 | Visceral neoplasia and epidermal papillomas in cultured turbot <i>Scophthalmus maximus</i> . <i>Diseases of Aquatic Organisms</i> , 1990, 8, 179-187.   | 0.5 | 20        |
| 169 | A review of the taxonomy and seroepizootiology of <i>Vibrio anguillarum</i> , with special reference to aquaculture in the northwest of Spain. <i>Diseases of Aquatic Organisms</i> , 1990, 9, 73-82.                             | 0.5 | 131       |
| 170 | Purification and characterization of an antibacterial substance produced by a marine <i>Alteromonas</i> species. <i>Antimicrobial Agents and Chemotherapy</i> , 1989, 33, 1674-1679.  | 1.4 | 30        |
| 171 | Influence of the growth conditions on the hydrophobicity of <i>Renibacterium salmoninarum</i> evaluated by different methods. <i>FEMS Microbiology Letters</i> , 1989, 60, 71-78.   | 0.7 | 15        |
| 172 | Ulcerative skin and fin lesions in reared turbot, <i>Scophthalmus maximus</i> (L.). <i>Journal of Fish Diseases</i> , 1989, 12, 323-333.  | 0.9 | 39        |
| 173 | Association of <i>Aeromonas sobria</i> with mortalities of adult gizzard shad, <i>Dorosoma cepedianum</i> Lesueur. <i>Journal of Fish Diseases</i> , 1989, 12, 439-448.   | 0.9 | 32        |
| 174 | Inhibitory activity of antibiotic-producing marine bacteria against fish pathogens. <i>Journal of Applied Bacteriology</i> , 1988, 65, 97-101.  | 1.1 | 146       |
| 175 | Evaluation of selective media for isolation and enumeration of vibrios from estuarine waters. <i>Journal of Microbiological Methods</i> , 1988, 8, 151-160.   | 0.7 | 56        |
| 176 | Chromosome-mediated iron uptake system in pathogenic strains of <i>Vibrio anguillarum</i> . <i>Journal of Bacteriology</i> , 1988, 170, 1920-1925.  | 1.0 | 120       |
| 177 | Virulence properties and enterotoxin production of <i>Aeromonas</i> strains isolated from fish. <i>Infection and Immunity</i> , 1988, 56, 3285-3293.  | 1.0 | 137       |
| 178 | Relationships among virulence for fish, enterotoxigenicity, and phenotypic characteristics of motile <i>Aeromonas</i> . <i>Aquaculture</i> , 1987, 67, 29-39.   | 1.7 | 29        |
| 179 | Homology of <i>Vibrio anguillarum</i> strains causing epizootics in turbot, salmon and trout reared on the Atlantic coast of Spain. <i>Aquaculture</i> , 1987, 67, 41-52.   | 1.7 | 77        |
| 180 | Bacillary necrosis in hatcheries of <i>Ostrea edulis</i> in Spain. <i>Aquaculture</i> , 1987, 65, 15-29.  | 1.7 | 56        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 181 | Specificity of slide agglutination test for detecting bacterial fish pathogens. <i>Aquaculture</i> , 1987, 61, 81-97.   | 1.7 | 90        |
| 182 | Evaluation of Different Assay Systems for Identification of Environmental <i>Aeromonas</i> Strains. <i>Applied and Environmental Microbiology</i> , 1986, 51, 652-656.  | 1.4 | 32        |
| 183 | Antibiotic activity of epiphytic bacteria isolated from intertidal seaweeds. <i>Microbial Ecology</i> , 1985, 11, 149-163.  | 1.4 | 194       |
| 184 | Modified Medium for the Oxidation-Fermentation Test in the Identification of Marine Bacteria. <i>Applied and Environmental Microbiology</i> , 1985, 49, 1541-1543.  | 1.4 | 117       |
| 185 | Relation of water temperature to infection of <i>Salmo gairdneri</i> with motile <i>Aeromonas</i> . <i>Fish Pathology</i> , 1985, 20, 99-105.   | 0.4 | 17        |
| 186 | Comparison between the bacterial flora associated with fingerling rainbow trout cultured in two different hatcheries in the North-West of Spain. <i>Aquaculture</i> , 1984, 42, 193-206.                        | 1.7 | 72        |
| 187 | Plasmid coding for transferable drug resistance in bacteria isolated from cultured rainbow trout. <i>Applied and Environmental Microbiology</i> , 1984, 48, 872-877.  | 1.4 | 53        |
| 188 | Haemagglutinating, haemolytic and cytotoxic activities of <i>Vibrio anguillarum</i> and related vibrios isolated from striped bass on the Atlantic Coast. <i>FEMS Microbiology Letters</i> , 1983, 18, 257-262. | 0.7 | 71        |
| 189 | Mechanism of poliovirus inactivation by cell-free filtrates of marine bacteria. <i>Canadian Journal of Microbiology</i> , 1983, 29, 1481-1486.  | 0.8 | 27        |
| 190 | Antiviral activity of antibiotic-producing marine bacteria. <i>Canadian Journal of Microbiology</i> , 1982, 28, 231-238.  | 0.8 | 36        |
| 191 | Comparative stability of two salmonid viruses and poliovirus in fresh, estuarine and marine waters. <i>Journal of Fish Diseases</i> , 1982, 5, 223-231.   | 0.9 | 58        |