

Jian Zhou

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6362961/publications.pdf>

Version: 2024-02-01

138
papers

2,662
citations

249298

26
h-index

286692

43
g-index

146
all docs

146
docs citations

146
times ranked

1618
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Rare examples of hybrid chalcogenoarsenate(III) incorporating trivalent vanadium complexes. <i>Dalton Transactions</i> , 2022, 51, 6876-6883. | 1.6 | 1 |
| 2 | Two luminescent cuprous iodides with hitherto-unknown free imidazolate sites for efficiently sensing Fe^{3+} and Cr^{2+} . <i>Journal of Materials Chemistry C</i> , 2022, 10, 6365-6373. | 2.7 | 6 |
| 3 | Deep-Red Luminescent Cuprous-Lead Bromide as a Dual-Responsive Sensor for Fe^{3+} and Cr^{2+} . <i>Inorganic Chemistry</i> , 2022, 61, 5957-5964. | 1.9 | 10 |
| 4 | A series of new lanthanide benzoates: Syntheses, crystal structures, and luminescent properties. <i>Dyes and Pigments</i> , 2022, 201, 110182. | 2.0 | 7 |
| 5 | Two luminescent lanthanide coordination polymers incorporating free pyridyl sites as the multi-responsive sensors for hazardous ions. <i>Dyes and Pigments</i> , 2022, 203, 110384. | 2.0 | 8 |
| 6 | One Octasubstituted Trisalkoxotetradecavanadate Cluster. <i>Inorganic Chemistry</i> , 2021, 60, 14-18. | 1.9 | 3 |
| 7 | A novel 3-D lead-iodide polymer based on the linkage of rare binuclear $[\text{Pb}_2\text{I}]^{3+}$ cations and anionic bis(pyrazinyl)-triazole bridges. <i>Dalton Transactions</i> , 2021, 50, 4486-4489. | 1.6 | 4 |
| 8 | Vanadoborates: cluster-based architectures, preparation and properties. <i>Dalton Transactions</i> , 2021, 50, 1550-1568. | 1.6 | 17 |
| 9 | One-Dimensional Vanadium(III) Chalcogenidostannates Incorporating $[\text{V}(\text{tepa})]^{3+}$ Complexes as Bridging Groups. <i>Inorganic Chemistry</i> , 2021, 60, 2127-2132. | 1.9 | 5 |
| 10 | Two Organic Hybrid Iodoplumbates Directed by a Bifunctional Bis(pyrazinyl)triazole. <i>Inorganic Chemistry</i> , 2021, 60, 5362-5366. | 1.9 | 10 |
| 11 | A Copper(I)-Thioarsenate(III) Inorganic Framework Directed by $[\text{Ni}(\text{en})_3]^{2+}$. <i>Inorganic Chemistry</i> , 2021, 60, 6813-6819. | 1.9 | 7 |
| 12 | An Organic Hybrid Indium-Telluride Incorporating Binuclear Complexes $[\text{In}_2(\text{ea})_4]^{2+}$ with a Bridging Oxygen Donor. <i>Inorganic Chemistry</i> , 2021, 60, 12724-12729. | 1.9 | 1 |
| 13 | The first selenidostannate directed by low-valent vanadium(II) complex: Photocurrent response and magnetic properties. <i>Inorganic Chemistry Communication</i> , 2021, 133, 108862. | 1.8 | 6 |
| 14 | A series of organic hybrid polyoxovanadate clusters incorporating tris(hydroxymethyl)methane derivatives. <i>Dalton Transactions</i> , 2021, 50, 15224-15232. | 1.6 | 0 |
| 15 | Two Organic Hybrid Manganese Selenoarsenates: The Discovery of One-Dimensional Low-Valent Selenoarsenate(II). <i>Inorganic Chemistry</i> , 2021, 60, 19226-19232. | 1.9 | 5 |
| 16 | A series of new polynuclear lanthanide(III) clusters prepared in alkylol amine. <i>Inorganica Chimica Acta</i> , 2020, 499, 119201. | 1.2 | 5 |
| 17 | Thermochromic luminescent properties of a tetrazole-functionalized iodocuprate without cuphophilic interaction. <i>Dyes and Pigments</i> , 2020, 174, 108039. | 2.0 | 13 |
| 18 | Hydrothermal syntheses, structures and properties of a series of new hybrid iodometallates containing rare cross-shaped spirobifluorene derivatives. <i>Materials Science and Engineering B: Solid-State Materials for Advanced Technology</i> , 2020, 262, 114693. | 1.7 | 2 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Two Hybrid Polymeric Iodoargentates Incorporating Aromatic N-Heterocycle Derivatives as Electron Acceptors. <i>Inorganic Chemistry</i> , 2020, 59, 16814-16818. | 1.9 | 12 |
| 20 | Syntheses and luminescent properties of a series of new lanthanide azelates. <i>Dyes and Pigments</i> , 2020, 182, 108638. | 2.0 | 10 |
| 21 | Unique Two-Dimensional Indium Telluride Templated by a Rare Wheel-Shaped Heterobimetallic Mn/In Cluster. <i>Inorganic Chemistry</i> , 2020, 59, 5818-5822. | 1.9 | 3 |
| 22 | A series of new hybrid chalcogenogermanates: the rare examples of chalcogenogermanates combined with trivalent vanadium complexes. <i>Dalton Transactions</i> , 2019, 48, 10907-10914. | 1.6 | 5 |
| 23 | Three new metal coordination polymers of bifunctional imidazolate/tetrazolate bridges: the only example of a three-dimensional framework based on rare $[\text{Co}_{4}(\text{I}^{3/4}\text{-OH})_{2}(\text{I}^{1/4}\text{-Cl})_{2}]^{4+}$ mixed oxo-chloro-clusters. <i>RSC Advances</i> , 2019, 9, 13082-13087. | 1.7 | 2 |
| 24 | A series of new oxo-vanadium(IV) complexes with octahedral coordinated vanadium centers. <i>Journal of Coordination Chemistry</i> , 2019, 72, 1064-1074. | 0.8 | 7 |
| 25 | A series of new vanadium(III) chalcogenido-antimonates: an unusual seven-coordinate nitro-selenidovanadium(III) complex. <i>Dalton Transactions</i> , 2019, 48, 3090-3097. | 1.6 | 7 |
| 26 | A unique formyl iodoargentate exhibiting luminescent and photocurrent response properties. <i>Dalton Transactions</i> , 2019, 48, 15762-15766. | 1.6 | 8 |
| 27 | A new 3-D cuprous thiogermanate with rare 3-D $[\text{Cu-S-Cu}]_n$ network. <i>Materials Today Communications</i> , 2018, 15, 88-93. | 0.9 | 5 |
| 28 | Two new 3-D cadmium bromoplumbates: the only example of heterometallic bromoplumbate based on crown $[\text{Cd}(\text{Pb}_{4}\text{O}_{4}\text{Br})_{2}]$ clusters. <i>Dalton Transactions</i> , 2018, 47, 4833-4839. | 1.6 | 9 |
| 29 | A novel 3-D cuprous iodide polymer with a high Cu/I ratio. <i>Dalton Transactions</i> , 2018, 47, 3253-3257. | 1.6 | 13 |
| 30 | A Series of Lanthanide Selenidogermanates: The First Coexistence of Three Types of Selenidogermanate Units in the Same Architecture. <i>Inorganic Chemistry</i> , 2018, 57, 1242-1250. | 1.9 | 10 |
| 31 | Metal-Free Azidation of α -Hydroxy Esters and α -Hydroxy Ketones Using Azidotrimethylsilane. <i>Advanced Synthesis and Catalysis</i> , 2018, 360, 1116-1122. | 2.1 | 16 |
| 32 | Two new 1-D lanthanide selenidogermanates with the $[\text{Ge}_2\text{Se}_6]^{4-}$ anion as a bridging ligand to a lanthanide complex cation. <i>Journal of Coordination Chemistry</i> , 2018, 71, 1093-1101. | 0.8 | 1 |
| 33 | An unusual cuprous iodide polymer incorporating I^{I} , I^{II} and I^{III} structural units. <i>Dalton Transactions</i> , 2018, 47, 17216-17220. | 1.6 | 12 |
| 34 | A New Type of Three-Dimensional Hybrid Polymeric Haloplumbate Based on Rare High-Nuclear Heterometallic Clusters. <i>Inorganic Chemistry</i> , 2018, 57, 12860-12868. | 1.9 | 31 |
| 35 | A series of new hybrid selenidostannates with metal complexes prepared in alkylol amines. <i>Dalton Transactions</i> , 2018, 47, 14751-14759. | 1.6 | 9 |
| 36 | A 2-D dysprosium glutarate exhibiting slow magnetic relaxation and luminescent properties. <i>Journal of Coordination Chemistry</i> , 2018, 71, 2722-2731. | 0.8 | 5 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | A Series of Lanthanide Chalcogenidogermanates Displaying Two Types of 1-D Polymeric Chains. <i>Journal of Cluster Science</i> , 2018, 29, 777-783. | 1.7 | 4 |
| 38 | Two new oxyiodoplumbates: the unique 3-D hybrid oxyiodoplumbate based on neutral 2-D [Pb ₂ I ₄]n layers. <i>Dalton Transactions</i> , 2018, 47, 8442-8447. | 1.6 | 7 |
| 39 | The only examples of cationic lanthanide pimelate frameworks decorated by π -conjugated 1,10-phenanthrolines. <i>Inorganica Chimica Acta</i> , 2018, 471, 377-383. | 1.2 | 6 |
| 40 | Syntheses, structures and properties of a series of new lanthanide chalcogenates(III) containing crown-shaped [As ₃ Q ₆]3 ⁿ⁻ (Q=As, Se) clusters. <i>Journal of Alloys and Compounds</i> , 2017, 702, 594-600. | 2.8 | 9 |
| 41 | A novel 2-D heterometallic polymer containing two types of 1-D cuprous polymeric chains and circular [V ₄ O ₁₂]4 ⁿ⁻ clusters. <i>Journal of Alloys and Compounds</i> , 2017, 713, 46-50. | 2.8 | 3 |
| 42 | A unique dysprosium selenoarsenate (<scp>iii</scp>) exhibiting a photocurrent response and slow magnetic relaxation behavior. <i>Dalton Transactions</i> , 2017, 46, 342-346. | 1.6 | 10 |
| 43 | A series of lanthanide glutarates: lanthanide contraction effect on crystal frameworks of lanthanide glutarates. <i>RSC Advances</i> , 2017, 7, 17934-17940. | 1.7 | 17 |
| 44 | A novel 3-D photoluminescent cuprous chloride polymer based on bifunctional imidazolate/tetrazolate bridges. <i>Dalton Transactions</i> , 2017, 46, 1372-1376. | 1.6 | 15 |
| 45 | Few-layer arsenic trichalcogenides: Emerging two-dimensional semiconductors with tunable indirect-direct band-gaps. <i>Journal of Alloys and Compounds</i> , 2017, 699, 554-560. | 2.8 | 33 |
| 46 | A novel 2-D Mn selenidostannate(IV) incorporating high-nuclear Mn clusters with spin canting behavior. <i>Dalton Transactions</i> , 2017, 46, 16009-16013. | 1.6 | 6 |
| 47 | A Series of Lanthanide Germanate Oxo Clusters Decorated by 1,10-Phenanthroline Chromophores. <i>Inorganic Chemistry</i> , 2017, 56, 10361-10369. | 1.9 | 24 |
| 48 | Hydrothermal Syntheses and Crystal Structure of a New Organic Hybrid Holmium Germanate Oxo-Cluster. <i>Journal of Cluster Science</i> , 2017, 28, 3209-3215. | 1.7 | 5 |
| 49 | Two New 3-D Lead(II) Coordination Polymers of Glycolic Acid with Luminescent Properties. <i>Journal of Inorganic and Organometallic Polymers and Materials</i> , 2017, 27, 181-188. | 1.9 | 2 |
| 50 | A Series of Dimeric Selenidogermanates with Lanthanide Complexes of Multidentate Chelating Amines. <i>Journal of Cluster Science</i> , 2017, 28, 2589-2600. | 1.7 | 2 |
| 51 | Catalytic Enantioselective Construction of Sulfur-Containing Tetrasubstituted Carbon Stereocenters. <i>ACS Catalysis</i> , 2016, 6, 5319-5344. | 5.5 | 118 |
| 52 | A 3-D net based on weak metallophilic (Cu ⁺ -Cu) interactions. <i>Dalton Transactions</i> , 2016, 45, 11292-11296. | 1.6 | 4 |
| 53 | Incorporation of manganese complexes into thioarsenates(V), displaying a new structural motif. <i>Journal of Coordination Chemistry</i> , 2016, 69, 3726-3734. | 0.8 | 5 |
| 54 | Synthesis, Crystal Structures and Properties of a Series of Lanthanide Adipates [Ln ₂ (ad) ₃ (H ₂ O) ₄] (Ln=ÅY ³⁺ , Ho ³⁺ , Er ³⁺ , Tm ³⁺). <i>Journal of Cluster Science</i> , 2016, 27, 2025-2033. | 1.7 | 7 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Solvothermal Syntheses and Characterization of a Series of New Selenidostannates with Lanthanide Complexes as Counterions. <i>Journal of Cluster Science</i> , 2016, 27, 1475-1484. | 1.7 | 3 |
| 56 | Syntheses, structures and properties of two new 3-D vanadoborates based on V O B clusters. <i>Journal of Alloys and Compounds</i> , 2016, 684, 537-543. | 2.8 | 10 |
| 57 | Synthesis of heterometallic chalcogenides containing lanthanide and group 13-15 metal elements. <i>Coordination Chemistry Reviews</i> , 2016, 315, 112-134. | 9.5 | 81 |
| 58 | A series of new lanthanide fumarates displaying three types of 3-D frameworks. <i>Dalton Transactions</i> , 2016, 45, 5253-5261. | 1.6 | 15 |
| 59 | Two Quaternary Copper Thiostannates with Lanthanum(III) Complexes. <i>Journal of Cluster Science</i> , 2016, 27, 257-265. | 1.7 | 12 |
| 60 | The first examples of 1-D organic hybrid lanthanoid thioarsenates based on two [AsVS ₄] ³⁻ linkage modes. <i>Dalton Transactions</i> , 2016, 45, 6015-6022. | 1.6 | 7 |
| 61 | Solvothermal Syntheses and Characterization of Three New Thioantimonates Combined with Lanthanide Complexes. <i>Journal of Cluster Science</i> , 2015, 26, 1333-1341. | 1.7 | 1 |
| 62 | A series of lanthanoid selenidoantimonates(v): rare examples of lanthanoid selenidoantimonates based on dinuclear lanthanide complexes. <i>Dalton Transactions</i> , 2015, 44, 6032-6039. | 1.6 | 19 |
| 63 | Two types of lanthanide selenidostannates(iv) first prepared under the same solvothermal conditions. <i>Dalton Transactions</i> , 2015, 44, 1350-1357. | 1.6 | 19 |
| 64 | Two New Lanthanide Complexes with Cluster [Ln ₂ O ₂] Units. <i>Journal of Cluster Science</i> , 2015, 26, 1503-1510. | 1.7 | 2 |
| 65 | One-Step Preparation of 1-D Copper(I) Polymer of Pyridyl-Ester with Fluorescence Properties. <i>Journal of Cluster Science</i> , 2015, 26, 1735-1742. | 1.7 | 0 |
| 66 | A series of new lanthanoid thioarsenates: insights into the influence of lanthanide contraction on the formation of new lanthanoid thioarsenates. <i>Dalton Transactions</i> , 2015, 44, 7203-7212. | 1.6 | 23 |
| 67 | A series of new manganese thioarsenates(ν) based on different unsaturated [Mn(amine) _x] ²⁺ complexes. <i>Dalton Transactions</i> , 2015, 44, 16430-16438. | 1.6 | 13 |
| 68 | A new solvothermal route to crystalline proustite Ag ₃ As ₃ S ₃ with photocatalytic properties. <i>Inorganic Chemistry Communication</i> , 2014, 46, 17-20. | 1.8 | 11 |
| 69 | Three new vanadoborates functionalized with zinc complexes. <i>Inorganic Chemistry Communication</i> , 2014, 43, 101-104. | 1.8 | 15 |
| 70 | [Ni(dap) ₃] ₄ [As ₁₀ Cu ₂ S ₁₈]: a new thioarsenate containing the rare [As ₃ CuS ₆] cluster with mixed-valence As ²⁺ /As ³⁺ ions. <i>Dalton Transactions</i> , 2014, 43, 3055-3058. | 1.6 | 24 |
| 71 | A series of new 3-D boratopolyoxovanadates containing five types of [KxOy] _n building units. <i>CrystEngComm</i> , 2014, 16, 4236. | 1.3 | 19 |
| 72 | Solvothermal syntheses of lanthanide thiogermanates displaying three new structural moieties. <i>RSC Advances</i> , 2014, 4, 38682. | 1.7 | 17 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Solvothermal syntheses and characterizations of three new holmium selenidostannates(IV): a rare example of adamantane-like $[\text{Sn}_4\text{Se}_{10}]_4^{4-}$ selenidostannate(IV) with lanthanide complexes. Dalton Transactions, 2014, 43, 12306. | 1.6 | 13 |
| 74 | Two Tetra-Cd ^{II} -Substituted Vanadogermanate Frameworks. Journal of the American Chemical Society, 2014, 136, 5065-5071. | 6.6 | 89 |
| 75 | The syntheses, structures and properties of three new lanthanoid thioarsenates: the only example of thioarsenate acting as a ligand to a lanthanide complex. Dalton Transactions, 2013, 42, 11155. | 1.6 | 20 |
| 76 | New 3-D polyoxovanadoborate architectures based on $[\text{V}_{12}\text{B}_{18}\text{O}_{60}]_{16}^{4-}$ clusters. CrystEngComm, 2013, 15, 5057. | 1.3 | 30 |
| 77 | Two new 3-D boratopolyoxovanadate architectures based on the $[\text{V}_{12}\text{B}_{16}\text{O}_{50}(\text{OH})_8]_{12}^{4-}$ cluster with different metal linkers. New Journal of Chemistry, 2013, 37, 4077. | 1.4 | 19 |
| 78 | A novel 1-D telluridoindate based on rare tetramer $[\text{In}_4\text{Te}_{10}]_4^{4-}$ unit with photocatalytic properties. CrystEngComm, 2013, 15, 1194-1198. | 1.3 | 10 |
| 79 | Solvothermal synthesis and characterization of thioindate-thioantimonates with transition-metal complexes: The first examples of the incorporation of transition metal ions into In-Sb frameworks. Dalton Transactions, 2013, 42, 1735-1742. | 1.6 | 22 |
| 80 | The first examples of thiogermanate anion $[\text{GeS}_3(\text{SH})]_3^{3-}$ as a bridging ligand to a lanthanide complex. Dalton Transactions, 2013, 42, 1961-1964. | 1.6 | 14 |
| 81 | $[\text{Ni}(\text{dien})_2]_3[\text{Ge}_3\text{Sb}_8\text{S}_{21}] \cdot 0.5\text{H}_2\text{O}$: A new 2-D layered thiogermanate-thioantimonate with metal complexes as template ions. Inorganic Chemistry Communication, 2013, 27, 92-96. | 1.8 | 16 |
| 82 | A new polymorph telluridoindate $[\text{In}(\text{en})_3][\text{In}_5\text{Te}_9(\text{en})_2]$ with photocatalytic properties. Inorganic Chemistry Communication, 2013, 28, 55-59. | 1.8 | 13 |
| 83 | Hydrothermal syntheses, crystal structures and characterization of new vanadoborates: The novel decorated cage cluster $[\text{V}_6\text{B}_{22}\text{O}_{44}(\text{OH})_{10}]$. Journal of Solid State Chemistry, 2013, 201, 79-84. | 1.4 | 26 |
| 84 | A 3-D chiral organic-inorganic hybrid zinc vanadate assembled from helical units. Dalton Transactions, 2013, 42, 5603-5606. | 1.6 | 20 |
| 85 | A novel 3-D chiral polyoxovanadate architecture based on breaking high symmetry of spherical $[\text{V}_{15}\text{O}_{36}\text{Cl}]_8^{8-}$ cluster. CrystEngComm, 2013, 15, 4593. | 1.3 | 20 |
| 86 | Solvothermal Syntheses and Characterization of Three Lanthanide Thioantimonates(V) with Mixed Ethylene Polyamines. Zeitschrift Fur Naturforschung - Section B Journal of Chemical Sciences, 2013, 68, 133-137. | 0.3 | 6 |
| 87 | A new 1-D extended vanadoborate containing triply bridged metal complex units. Inorganic Chemistry Communication, 2012, 25, 51-54. | 1.8 | 23 |
| 88 | One unprecedented 1-D europium thioindate-thioantimonate based on heterometallic mixed nitro-thioclusters with photoluminescent properties. Chemical Communications, 2012, 48, 2537. | 2.2 | 35 |
| 89 | A series of new lanthanoid thioantimonates: a rare example of organic-decorated cationic lanthanoid thioantimonate chains based on asymmetric dinuclear lanthanide complexes. CrystEngComm, 2012, 14, 5544. | 1.3 | 21 |
| 90 | Solvothermal Synthesis and Characterization of a Series of Lanthanide Thioantimonates(IV): The First Examples of Inorganic-Organic Hybrid Cationic Lanthanide Thioantimonates(IV). Inorganic Chemistry, 2012, 51, 2283-2290. | 1.9 | 36 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 91 | Solvothermal synthesis, crystal structures and properties of three new thiogermanates: the only example of the thiogermanate anion $[\text{Ge}_2\text{S}_6]^{4-}$ as a bridging ligand to a lanthanide complex ion. <i>CrystEngComm</i> , 2012, 14, 3464. | 1.3 | 26 |
| 92 | One novel 3-D vanadoborate with unusual 3-D Na $\text{O}^{\text{O}}\text{Na}$ network. <i>RSC Advances</i> , 2012, 2, 10937. | 1.7 | 29 |
| 93 | A New 1D Polyoxovanadate $[\text{Cu}(\text{en})_2\text{V}_{10}\text{O}_{28}] [\text{Cu}(\text{en})_2(\text{H}_2\text{O})_2] \cdot 2\text{H}_3\text{BO}_3 \cdot 2\text{H}_2\text{O}$ Containing the Rarely Non-condensed Boric acid. <i>Zeitschrift Fur Naturforschung - Section B Journal of Chemical Sciences</i> , 2012, 67, 860-864. | 0.3 | 3 |
| 94 | Novel lanthanoid thioantimonates: the first coexistence of different types of thioantimonate anions in the same framework. <i>Dalton Transactions</i> , 2012, 41, 11760. | 1.6 | 24 |
| 95 | Solvothermal synthesis and characterization of two 2-D layered germanium thioantimonates with transition-metal complexes. <i>Dalton Transactions</i> , 2011, 40, 11419. | 1.6 | 33 |
| 96 | A novel 3-D thioindate-thioantimonate based on the linkages of large heterometallic $\{\text{In}_2\text{Sb}_2\text{S}_9\}$ clusters and 1-D $[\text{In}_2\text{Sb}_2\text{S}_8]_n$ chains. <i>CrystEngComm</i> , 2011, 13, 5924. | 1.3 | 22 |
| 97 | Solvothermal synthesis and characterization of two novel europium thioantimonates(iii) containing $[\text{SbIII}_3]$ unit as an unusual chelating ligand. <i>CrystEngComm</i> , 2011, 13, 4806. | 1.3 | 19 |
| 98 | $[\text{M}(\text{dap})_3]\text{InSb}_3\text{S}_7$ (M = Co, Ni): Two Novel Open-Framework Thioindate \sim Thioantimonates with 8-, 12-, and 16-Ring Intersecting Channels. <i>Inorganic Chemistry</i> , 2011, 50, 415-417. | 1.9 | 41 |
| 99 | Two Organic Hybrid Thiogermanates $[\text{Ni}(\text{dien})_2]_2(\text{H}_2\text{dien})\text{Ge}_2\text{S}_6$ and $[\text{Ni}(\text{teta})_2]_2\text{Ge}_4\text{S}_{10}$. <i>Zeitschrift Fur Naturforschung - Section B Journal of Chemical Sciences</i> , 2011, 66, 659-663. | 0.3 | 3 |
| 100 | Hydrothermal Synthesis and Structure of Two 1-D Organic HybridMetal Sulfates. <i>Zeitschrift Fur Naturforschung - Section B Journal of Chemical Sciences</i> , 2011, 66, 1127-1130. | 0.3 | 0 |
| 101 | Solvothermal Syntheses and Crystal Structures of Two Thioarsenate Metal Complexes $[\text{Co}(\text{dien})_2]_3[\text{As}_3\text{S}_6]_2$ and $[\text{Mn}(\text{teta})_2]\text{As}_2\text{S}_5$. <i>Zeitschrift Fur Naturforschung - Section B Journal of Chemical Sciences</i> , 2011, 66, 366-370. | 0.3 | 1 |
| 102 | Hydrothermal Syntheses and Crystal Structures of Two New Heteropolyoxovanadoborates Containing $\{(\text{VO})_{12}\text{O}_6[\text{B}_3\text{O}_6(\text{OH})_6(\text{H}_2\text{O})]\}$ Cluster. <i>Journal of Cluster Science</i> , 2011, 22, 65-72. | 1.7 | 25 |
| 103 | Solvothermal syntheses and crystal structures of two new thiogermanates $[\text{M}(\text{dap})_3]_4\text{Ge}_4\text{S}_{10}\text{Cl}_4$ (M = Co, Ni) with metal complexes as counterions. <i>Monatshefte Für Chemie</i> , 2011, 142, 763-768. | 0.9 | 16 |
| 104 | Two Novel Adamantane Like Thio/Selenidogermanates with Complex Cations. <i>Zeitschrift Fur Anorganische Und Allgemeine Chemie</i> , 2011, 637, 1388-1393. | 0.6 | 19 |
| 105 | Solvothermal synthesis of two new thioantimonates with transition-metal complexes $[\text{Co}(\text{dien})_2]_4[\text{CoSb}_6\text{S}_{14}]$ and $[\text{Co}(\text{dien})_2]_2[\text{Sb}_4\text{S}_9]$. <i>Inorganic Chemistry Communication</i> , 2011, 14, 1286-1289. | 1.8 | 18 |
| 106 | The New Vanadoborate-supported Hexanuclear Zinc Complex $[\text{Zn}(\text{teta})_6][(\text{VO})_{12}\text{O}_6\text{B}_{18}\text{O}_{36}(\text{OH})_6] \cdot 4(\text{H}_2\text{O}) \cdot 8\text{H}_2\text{O}$. <i>Zeitschrift Fur Naturforschung - Section B Journal of Chemical Sciences</i> , 2011, 66, 115-118. | 0.3 | 4 |
| 107 | Determination of Cd in Water Samples by Hollow-Fiber Supported Liquid-Membrane Extraction Coupled with Thermospray-Flame Furnace Atomic-Absorption Spectrometry. <i>Spectroscopy Letters</i> , 2011, 44, 278-284. | 0.5 | 5 |
| 108 | Two New Hemidirected Lead(II) Complexes: $[\text{Pb}(\text{pcih})(\text{bha})]$ and $[\text{Pb}(\text{pcih})(\text{NO}_3)_3]$. <i>Zeitschrift Fur Naturforschung - Section B Journal of Chemical Sciences</i> , 2010, 65, 1084-1088. | 0.3 | 9 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Solvothermal Syntheses and Crystal Structures of Two Thiostannates(IV) $[M(\text{tepa})]_2(\frac{1}{4}\text{-Sn}_2\text{S}_6)$ ($M = \text{Fe}^{2+}$) <i>Tj ETQq1 1 0.784314 rgB</i> | 0.3 | 17 |
| 110 | Two 3-D supramolecular architecture based on the linkages of Co(II) complexes and lattice water molecules. <i>Structural Chemistry</i> , 2010, 21, 159-164. | 1.0 | 8 |
| 111 | A Series of Open-Frame Aluminoborates Templated by Transition-Metal Complexes. <i>Chemistry - A European Journal</i> , 2010, 16, 4852-4863. | 1.7 | 103 |
| 112 | A Series of Vanadogermanates from 1D Chain to 3D Framework Built by Ge ^{IV} -O Clusters and Transition-Metal-Complex Bridges. <i>Chemistry - A European Journal</i> , 2010, 16, 13253-13261. | 1.7 | 54 |
| 113 | Two Supramolecular Architectures Containing Dinuclear Thiostannate $[\text{Sn}_2\text{S}_6]$ Units. <i>Zeitschrift Fur Naturforschung - Section B Journal of Chemical Sciences</i> , 2010, 65, 1229-1234. | 0.3 | 17 |
| 114 | Two Novel Thioindate-Thioantimonate Compounds $[\text{Ni}(\text{dien})_2]_2\text{In}_2\text{Sb}_4\text{S}_{11}$ and $[\text{Ni}(\text{dien})_2]_3(\text{In}_3\text{Sb}_2\text{S}_9)_2 \cdot 2\text{H}_2\text{O}$ with Transition Metal Complexes. <i>Inorganic Chemistry</i> , 2010, 49, 9671-9676. | 1.9 | 39 |
| 115 | A New 2-D Network Containing $\{\text{As}_4\text{V}_6\text{O}_{42}(\text{H}_2\text{O})\}$ Cluster Units. <i>European Journal of Inorganic Chemistry</i> , 2009, 2009, 5075-5078. | 1.0 | 31 |
| 116 | Two 3-D Supramolecular Networks Containing Dimeric $\{\text{Cu}_2\text{X}_2\}$ Cluster Units. <i>Journal of Cluster Science</i> , 2009, 20, 555-563. | 1.7 | 8 |
| 117 | Solvothermal Synthesis of Two Cationic Indium Selenides with I^+ Ion as Counterion. <i>Zeitschrift Fur Anorganische Und Allgemeine Chemie</i> , 2009, 635, 151-155. | 0.6 | 8 |
| 118 | Solvothermal synthesis of Group 13-15 chalcogenidometalates with chelating organic amines. <i>Coordination Chemistry Reviews</i> , 2009, 253, 1221-1247. | 9.5 | 247 |
| 119 | A New 2D Hybrid Indate-Germanate Based on the Linkages of $\text{Ge}_4\text{O}_{10}\text{F}_2$ Clusters and $\{\text{In}_2\text{O}_6\}$ Dimers. <i>Inorganic Chemistry</i> , 2009, 48, 10895-10897. | 1.9 | 11 |
| 120 | Three 1-D selenidogallates $[\text{GaSe}_2]_n$, displaying conformational variations. <i>Journal of Coordination Chemistry</i> , 2009, 62, 1112-1120. | 0.8 | 6 |
| 121 | A new layered aluminoborate $[\text{Zn}(\text{dien})_2][\{\text{Al}(\text{OH})\}\{\text{B}_5\text{O}_9\text{F}\}]$ templated by transition metal complexes. <i>CrystEngComm</i> , 2009, 11, 2597. | 1.3 | 55 |
| 122 | A New Type of 1-D Thioindates, $[\text{M}(\text{en})_3]_0.5[\text{InS}_2]$ ($M = \text{Co}, \text{Ni}$), Synthesized by Solvothermal Reaction. <i>Zeitschrift Fur Naturforschung - Section B Journal of Chemical Sciences</i> , 2009, 64, 504-508. | 0.3 | 9 |
| 123 | Two novel extended lead(II) coordination polymers generated from bridging Schiff-base ligands. <i>Inorganic Chemistry Communication</i> , 2008, 11, 367-371. | 1.8 | 20 |
| 124 | The first polymeric thiogallate with lanthanide-containing counter cation, $[\text{Dy}_2(\text{en})_6(\frac{1}{4}\text{-OH})_2]\text{Ga}_4\text{S}_8$. <i>Inorganic Chemistry Communication</i> , 2008, 11, 1327-1329. | 1.8 | 9 |
| 125 | Structural Study of Organic-Inorganic Hybrid Thiogallates and Selenidogallates in View of Effects of the Chelate Amines. <i>Crystal Growth and Design</i> , 2008, 8, 2235-2240. | 1.4 | 37 |
| 126 | Cationic Main-Group Metal Chalcogenide with a 1-D Polymeric Structure, $[\{\text{In}(\text{dien})\}_2(\text{InTe})_4]\cdot\text{Cl}$. <i>Inorganic Chemistry</i> , 2008, 47, 8586-8588. | 1.9 | 21 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 127 | A Photoluminescent zig-zag 1D Coordination Polymer with Mixed Ligands [Zn(absa)(pcih)](CH ₃ OH). Zeitschrift Fur Naturforschung - Section B Journal of Chemical Sciences, 2008, 63, 1343-1346. | 0.3 | 4 |
| 128 | Synthesis of a Selenidostannate(IV), [Mn(tepa)Sn ₃ Se ₇] _n , Demonstrating the Transformation from Achiral to Chiral and Dimeric to Polymeric Structure. Inorganic Chemistry, 2007, 46, 1541-1543. | 1.9 | 69 |
| 129 | One-Dimensional Indium Sulfides with Transition Metal Complexes of Polyamines. Inorganic Chemistry, 2007, 46, 6347-6352. | 1.9 | 67 |
| 130 | Synthesis, crystal structures of [M(tepa)] ₂ (^{1/4} Sn ₂ Se ₆)(M=Fe and Co), showing a selenidostannate bridge for transition metal complex cations. Inorganic Chemistry Communication, 2007, 10, 348-351. | 1.8 | 36 |
| 131 | A new hemidirected lead(II) complex converted into holodirected by secondary Pb ^{II} ⋯N interactions. Inorganic Chemistry Communication, 2007, 10, 475-478. | 1.8 | 17 |
| 132 | Solvothermal Synthesis of two 2D Thioantimonates(III) with Metal Complexes as Template Ions, [M(dap) ₃]Sb ₄ S ₇ (M = Ni ²⁺ and Co ²⁺). Zeitschrift Fur Anorganische Und Allgemeine Chemie, 2007, 633, 2701-2705. | 0.6 | 29 |
| 133 | Solvothermal Synthesis of a Series of Telluridoindate Compounds Displaying Three Types of One-Dimensional Polymeric Anion Chains with the Formula {[InTe ₂]} _n . Crystal Growth and Design, 2007, 7, 1889-1892. | 1.4 | 42 |
| 134 | Synthesis and crystal structure of a three-dimensional network coordination polymer: [Pb(C ₅ H ₄ NCOO) ₂ (H ₂ O)] _n . Journal of Coordination Chemistry, 2006, 59, 1477-1482. | 0.8 | 20 |
| 135 | Luminescent 2-D Double-layered Polymer, [(CuI) ₄ (CH ₃ SCH ₃) ₃] _n , Containing Helical Chains Constructed by Flower-Basket-Shaped Cu ₄ I ₄ Clusters. Inorganic Chemistry, 2006, 45, 8486-8488. | 1.9 | 58 |
| 136 | Solvothermal Synthesis, Crystal Structures, and Properties of New Selenidoantimonates [Ln(en) ₄ (SbSe ₄)] (Ln = La, Nd) and [Sm(en) ₄]SbSe ₄ ·0.5en: The First Example of an SbSe ₄ ³⁻ Anion Acting as a Ligand to a Lanthanide Complex. European Journal of Inorganic Chemistry, 2006, 2006, 2760-2765. | 1.0 | 30 |
| 137 | Synthesis and crystal structure of a two-dimensional bi-layered supramolecular network compound [Cu ₂ (C ₇ H ₈ N ₄ O) ₄](ClO ₄) ₂ (C ₇ H ₈ N ₄ O=pyridine-3-carbaldehyde semicarbazone). Journal of Coordination Chemistry, 2006, 59, 1711-1717. | 0.8 | 2 |
| 138 | Catalytic behavior of silica-supported chitosan-platinum-iron complex for asymmetric hydrogenation of ketones. Polymers for Advanced Technologies, 2004, 15, 287-290. | 1.6 | 15 |