
Paulo R Menezes

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/636553/publications.pdf Version: 2024-02-01



DALLIO P MENEZES

#	Article	IF	CITATIONS
1	Global prevalence of dementia: a Delphi consensus study. Lancet, The, 2005, 366, 2112-2117.	13.7	4,496
2	Chronic non-communicable diseases in Brazil: burden and current challenges. Lancet, The, 2011, 377, 1949-1961.	13.7	979
3	Mapping genomic loci implicates genes and synaptic biology in schizophrenia. Nature, 2022, 604, 502-508.	27.8	929
4	The contribution of cannabis use to variation in the incidence of psychotic disorder across Europe (EU-GEI): a multicentre case-control study. Lancet Psychiatry,the, 2019, 6, 427-436.	7.4	528
5	Dementia diagnosis in developing countries: a cross-cultural validation study. Lancet, The, 2003, 361, 909-917.	13.7	489
6	Métodos para estimar razão de prevalência em estudos de corte transversal. Revista De Saude Publica, 2008, 42, 992-998.	1.7	381
7	Drug and Alcohol Problems among Individuals with Severe Mental Illnesses in South London. British Journal of Psychiatry, 1996, 168, 612-619.	2.8	354
8	Religious and spiritual interventions in mental health care: a systematic review and meta-analysis of randomized controlled clinical trials. Psychological Medicine, 2015, 45, 2937-2949.	4.5	251
9	Treated Incidence of Psychotic Disorders in the Multinational EU-GEI Study. JAMA Psychiatry, 2018, 75, 36.	11.0	235
10	IQ and non-clinical psychotic symptoms in 12-year-olds: results from the ALSPAC birth cohort. British Journal of Psychiatry, 2008, 193, 185-191.	2.8	233
11	Identifying Gene-Environment Interactions in Schizophrenia: Contemporary Challenges for Integrated, Large-scale Investigations. Schizophrenia Bulletin, 2014, 40, 729-736.	4.3	229
12	Care arrangements for people with dementia in developing countries. International Journal of Geriatric Psychiatry, 2004, 19, 170-177.	2.7	226
13	Prevalence of anxiety and depression during pregnancy in a private setting sample. Archives of Women's Mental Health, 2007, 10, 25-32.	2.6	225
14	Methods for estimating prevalence ratios in cross-sectional studies. Revista De Saude Publica, 2008, 42, 992-8.	1.7	187
15	Age-related gray matter volume changes in the brain during non-elderly adulthood. Neurobiology of Aging, 2011, 32, 354-368.	3.1	158
16	Substance misuse and risk of aggression and offending among the severely mentally ill. British Journal of Psychiatry, 1998, 172, 345-350.	2.8	152
17	Very low levels of education and cognitive reserve. Neurology, 2013, 81, 650-657.	1.1	133
18	High prevalence of dementia among older adults from poor socioeconomic backgrounds in São Paulo, Brazil. International Psychogeriatrics, 2008, 20, 394-405.	1.0	125

#	Article	IF	CITATIONS
19	Limitations of the Mini-Mental State Examination for screening dementia in a community with low socioeconomic status. European Archives of Psychiatry and Clinical Neuroscience, 2009, 259, 8-15.	3.2	120
20	Investigating whether adverse prenatal and perinatal events are associated with non-clinical psychotic symptoms at age 12 years in the ALSPAC birth cohort. Psychological Medicine, 2009, 39, 1457-1467.	4.5	102
21	Maternal tobacco, cannabis and alcohol use during pregnancy and risk of adolescent psychotic symptoms in offspring. British Journal of Psychiatry, 2009, 195, 294-300.	2.8	95
22	Preterm birth as a risk factor for postpartum depression: A systematic review and meta-analysis. Journal of Affective Disorders, 2019, 259, 392-403.	4.1	90
23	Behavioral and Psychological Symptoms of Dementia in developing countries. International Psychogeriatrics, 2004, 16, 441-459.	1.0	87
24	Common mental disorders during pregnancy: prevalence and associated factors among low-income women in São Paulo, Brazil. Archives of Women's Mental Health, 2009, 12, 335-343.	2.6	85
25	Risk factors across the life course and dementia in a Brazilian population: results from the Sao Paulo Ageing & Health Study (SPAH). International Journal of Epidemiology, 2008, 37, 879-890.	1.9	79
26	Caregiver burden in an elderly population with depression in São Paulo, Brazil. Social Psychiatry and Psychiatric Epidemiology, 2002, 37, 416-422.	3.1	77
27	Effects of education and culture on the validity of the Geriatric Mental State and its AGECAT algorithm. British Journal of Psychiatry, 2004, 185, 429-436.	2.8	76
28	The detection of depression in medical setting: A study with PRIME-MD. Journal of Affective Disorders, 2006, 91, 11-17.	4.1	74
29	Esgotamento profissional e transtornos mentais comuns em agentes comunitÃ;rios de saúde. Revista De Saude Publica, 2008, 42, 921-929.	1.7	73
30	School bullying: A systematic review of contextual-level risk factors in observational studies. Aggression and Violent Behavior, 2015, 22, 65-76.	2.1	70
31	Transdiagnostic dimensions of psychopathology at first episode psychosis: findings from the multinational EU-GEI study. Psychological Medicine, 2019, 49, 1378-1391.	4.5	69
32	Postpartum depression: in relation to life events and patterns of coping. Archives of Women's Mental Health, 2004, 7, 123-131.	2.6	68
33	Investigation of the possible association between nosocomial candiduria and candidaemia. Clinical Microbiology and Infection, 2006, 12, 538-543.	6.0	64
34	Cross-cultural validation of the Chalder Fatigue Questionnaire in Brazilian primary care. Journal of Psychosomatic Research, 2007, 62, 301-304.	2.6	64
35	Inequities in access to depression treatment: results of the Brazilian National Health Survey – PNS. International Journal for Equity in Health, 2016, 15, 154.	3.5	64
36	Impact of Adherence to Antiretroviral Therapy in HIV-1–Infected Patients at a University Public Service in Brazil. AIDS Patient Care and STDs, 2001, 15, 587-593.	2.5	63

#	Article	IF	CITATIONS
37	Subclinical hyperthyroidism and dementia: the Sao Paulo Ageing & Health Study (SPAH). BMC Public Health, 2010, 10, 298.	2.9	63
38	Investigating if psychosis-like symptoms (PLIKS) are associated with family history of schizophrenia or paternal age in the ALSPAC birth cohort. Schizophrenia Research, 2008, 104, 279-286.	2.0	62
39	Discriminating sarcopenia in community-dwelling older women with high frequency of overweight/obesity: the São Paulo Ageing & Health Study (SPAH). Osteoporosis International, 2013, 24, 595-603.	3.1	62
40	Complementary religious and spiritual interventions in physical health and quality of life: A systematic review of randomized controlled clinical trials. PLoS ONE, 2017, 12, e0186539.	2.5	62
41	Arg72Pro <i>TP53</i> polymorphism and cancer susceptibility: A comprehensive metaâ€analysis of 302 caseâ€control studies. International Journal of Cancer, 2011, 129, 920-930.	5.1	61
42	Brain Structural Variability due to Aging and Gender in Cognitively Healthy Elders: Results from the São Paulo Ageing and Health Study. American Journal of Neuroradiology, 2009, 30, 1850-1856.	2.4	60
43	Paternal and maternal ages at conception and risk of bipolar affective disorder in their offspring. Psychological Medicine, 2010, 40, 477-485.	4.5	60
44	Violence at work and depressive symptoms in primary health care teams: a cross-sectional study in Brazil. Social Psychiatry and Psychiatric Epidemiology, 2015, 50, 1347-1355.	3.1	60
45	Identification of Dementia Cases in the Community: A Brazilian Experience. Journal of the American Geriatrics Society, 2005, 53, 1738-1742.	2.6	59
46	Social disadvantage, linguistic distance, ethnic minority status and first-episode psychosis: results from the EU-GEI case–control study. Psychological Medicine, 2021, 51, 1536-1548.	4.5	58
47	ERICA: prevalence of common mental disorders in Brazilian adolescents. Revista De Saude Publica, 2016, 50, 14s.	1.7	57
48	Incidence of first-contact psychosis in São Paulo, Brazil. British Journal of Psychiatry, 2007, 191, s102-s106.	2.8	56
49	Grey matter abnormalities in Brazilians with first-episode psychosis. British Journal of Psychiatry, 2007, 191, s117-s122.	2.8	55
50	Antenatal Depression Strongly Predicts Postnatal Depression in Primary Health Care. Revista Brasileira De Psiquiatria, 2012, 34, 446-450.	1.7	55
51	Depression and health behaviors in Brazilian adults – PNS 2013. Revista De Saude Publica, 2017, 51, 8s.	1.7	55
52	The worldwide prevalence of insufficient physical activity in adolescents; a systematic review. Nutricion Hospitalaria, 2013, 28, 575-84.	0.3	55
53	XPC polymorphisms play a role in tissue-specific carcinogenesis: a meta-analysis. European Journal of Human Genetics, 2008, 16, 724-734.	2.8	54
54	Prevalence and risk factors of radiographic vertebral fracture in Brazilian community-dwelling elderly. Osteoporosis International, 2011, 22, 711-719.	3.1	53

#	Article	IF	CITATIONS
55	Lack of progression of brain abnormalities in first-episode psychosis: a longitudinal magnetic resonance imaging study. Psychological Medicine, 2011, 41, 1677-1689.	4.5	53
56	Late Entry into HIV Care: Estimated Impact on AIDS Mortality Rates in Brazil, 2003–2006. PLoS ONE, 2011, 6, e14585.	2.5	52
57	Neuroanatomical pattern classification in a population-based sample of first-episode schizophrenia. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2013, 43, 116-125.	4.8	52
58	Cognitive deficits in first-episode psychosis: A population-based study in São Paulo, Brazil. Schizophrenia Research, 2007, 90, 338-343.	2.0	51
59	Validity of the self reporting questionnaire-20 in epidemiological studies with older adults. Social Psychiatry and Psychiatric Epidemiology, 2009, 44, 247-254.	3.1	50
60	Service use and costs of people with dual diagnosis in South London. Acta Psychiatrica Scandinavica, 2000, 101, 464-472.	4.5	47
61	Prevalência do autorrelato de depressão no Brasil: resultados da Pesquisa Nacional de Saúde, 2013. Revista Brasileira De Epidemiologia, 2015, 18, 170-180.	0.8	47
62	Cannabis use, cognition and brain structure in first-episode psychosis. Schizophrenia Research, 2013, 147, 209-215.	2.0	46
63	Cognitive performance is related to cortical grey matter volumes in early stages of schizophrenia: A population-based study of first-episode psychosis. Schizophrenia Research, 2009, 113, 200-209.	2.0	45
64	A population-based morphometric MRI study in patients with first-episode psychotic bipolar disorder: comparison with geographically matched healthy controls and major depressive disorder subjects. Bipolar Disorders, 2011, 13, 28-40.	1.9	45
65	Risk factors for colonisation of newborn infants during an outbreak of extended-spectrum β-lactamase-producing Klebsiella pneumoniae in an intermediate-risk neonatal unit. Journal of Hospital Infection, 2009, 71, 340-347.	2.9	44
66	The role of literacy, occupation and income in dementia prevention: the São Paulo Ageing & Health Study (SPAH). International Psychogeriatrics, 2010, 22, 1209-1215.	1.0	44
67	Neuroanatomical Classification in a Population-Based Sample of Psychotic Major Depression and Bipolar I Disorder with 1 Year of Diagnostic Stability. BioMed Research International, 2014, 2014, 1-9.	1.9	44
68	Comparative epidemiology of chronic fatigue syndrome in Brazilian and British primary care: prevalence and recognition. British Journal of Psychiatry, 2009, 194, 117-122.	2.8	43
69	Temporal Relationship Between Intimate Partner Violence and Postpartum Depression in a Sample of Low Income Women. Maternal and Child Health Journal, 2013, 17, 1297-1303.	1.5	43
70	Suicidal ideation during pregnancy: prevalence and associated factors among low-income women in São Paulo, Brazil. Archives of Women's Mental Health, 2012, 15, 135-138.	2.6	41
71	Cytokine profile in first-episode psychosis, unaffected siblings and community-based controls: the effects of familial liability and childhood maltreatment. Psychological Medicine, 2020, 50, 1139-1147.	4.5	41
72	The EUropean Network of National Schizophrenia Networks Studying Gene–Environment Interactions (EU-GEI): Incidence and First-Episode Case–Control Programme. Social Psychiatry and Psychiatric Epidemiology, 2020, 55, 645-657.	3.1	41

#	Article	IF	CITATIONS
73	Age-Related Metabolic Profiles in Cognitively Healthy Elders: Results from a Voxel-Based [¹⁸ F]Fluorodeoxyglucose–Positron-Emission Tomography Study with Partial Volume Effects Correction: Fig 1 American Journal of Neuroradiology, 2011, 32, 560-565.	2.4	40
74	Population impact of depression on functional disability in elderly: results from "São Paulo Ageing & Health Study―(SPAH). European Archives of Psychiatry and Clinical Neuroscience, 2013, 263, 153-158.	3.2	40
75	Vertebral Fracture Assessment by Dual Xâ€Ray Absorptiometry: A Valid Tool to Detect Vertebral Fractures in Communityâ€Dwelling Older Adults in a Populationâ€Based Survey. Arthritis Care and Research, 2013, 65, 809-815.	3.4	40
76	Depression and excess mortality in the elderly living in low―and middleâ€income countries: Systematic review and metaâ€analysis. International Journal of Geriatric Psychiatry, 2019, 34, 22-30.	2.7	40
77	Effect of a Digital Intervention on Depressive Symptoms in Patients With Comorbid Hypertension or Diabetes in Brazil and Peru. JAMA - Journal of the American Medical Association, 2021, 325, 1852.	7.4	40
78	White-matter hyperintensities in first-episode psychosis. British Journal of Psychiatry, 2008, 193, 25-30.	2.8	39
79	Unplanned pregnancy and risk of maternal depression: secondary data analysis from a prospective pregnancy cohort. Psychology, Health and Medicine, 2017, 22, 65-74.	2.4	39
80	Daily use of high-potency cannabis is associated with more positive symptoms in first-episode psychosis patients: the EU-GEI case–control study. Psychological Medicine, 2021, 51, 1329-1337.	4.5	38
81	Prevalence of thyroid disorders among older people: results from the São Paulo Ageing & Health Study. Cadernos De Saude Publica, 2011, 27, 155-161.	1.0	37
82	Atherosclerosis and Dementia. Stroke, 2011, 42, 3614-3615.	2.0	37
83	Meta-Analysis of Sensorimotor Gating Deficits in Patients With Schizophrenia Evaluated by Prepulse Inhibition Test. Schizophrenia Bulletin, 2020, 46, 1482-1497.	4.3	37
84	Use of a Mobile Phone App to Treat Depression Comorbid With Hypertension or Diabetes: A Pilot Study in Brazil and Peru. JMIR Mental Health, 2019, 6, e11698.	3.3	36
85	Acinetobacter spp. are associated with a higher mortality in intensive care patients with bacteremia: a survival analysis. BMC Infectious Diseases, 2016, 16, 386.	2.9	35
86	The Relationship Between Mode of Delivery and Sexual Health Outcomes after Childbirth. Journal of Sexual Medicine, 2015, 12, 1212-1220.	0.6	34
87	<i>GRIN2B</i> promoter methylation deficits in early-onset schizophrenia and its association with cognitive function. Epigenomics, 2019, 11, 401-410.	2.1	34
88	Jumping to conclusions, general intelligence, and psychosis liability: findings from the multi-centre EU-GEI case-control study. Psychological Medicine, 2021, 51, 623-633.	4.5	34
89	Predictors of clinical and social outcomes after hospitalization in schizophrenia. European Archives of Psychiatry and Clinical Neuroscience, 1997, 247, 137-145.	3.2	33
90	Consumo de frutas e hortaliças por idosos de baixa renda na cidade de São Paulo. Revista De Saude Publica, 2009, 43, 806-813.	1.7	33

#	Article	IF	CITATIONS
91	Association of Appendicular Lean Mass, and Subcutaneous and Visceral Adipose Tissue With Mortality in Older Brazilians: The São Paulo Ageing & Health Study. Journal of Bone and Mineral Research, 2019, 34, 1264-1274.	2.8	33
92	Longitudinal follow-up of cavum septum pellucidum and adhesio interthalamica alterations in first-episode psychosis: a population-based MRI study. Psychological Medicine, 2012, 42, 2523-2534.	4.5	31
93	Serum phosphate and hip bone mineral density as additional factors for high vascular calcification scores in a community-dwelling: The São Paulo Ageing & Health Study (SPAH). Bone, 2013, 52, 354-359.	2.9	31
94	Factors associated with risk of malnutrition in the elderly in south-eastern Brazil. Revista Brasileira De Epidemiologia, 2017, 20, 598-610.	0.8	31
95	CHEMONUCLEOLYSIS IN LUMBAR DISC HERNIATION: A META-ANALYSIS. Clinics, 2007, 62, 175-180.	1.5	30
96	Maternity "Blues― Prevalence and Risk Factors. Spanish Journal of Psychology, 2008, 11, 593-599.	2.1	29
97	Relationship between Brain Age-Related Reduction in Gray Matter and Educational Attainment. PLoS ONE, 2015, 10, e0140945.	2.5	29
98	Patterns of health-related behaviours among adolescents: a cross-sectional study based on the National Survey of School Health Brazil 2012. BMJ Open, 2016, 6, e011571.	1.9	29
99	The Relationship Between Depressive/Anxiety Symptoms During Pregnancy/Postpartum and Sexual Life Decline after Delivery. Journal of Sexual Medicine, 2013, 10, 1343-1349.	0.6	28
100	Individual and contextual factors associated with verbal bullying among Brazilian adolescents. BMC Pediatrics, 2015, 15, 49.	1.7	28
101	Cannabis use as a risk factor for psychoticâ€like experiences: A systematic review of nonâ€clinical populations evaluated with the Community Assessment of Psychic Experiences. Microbial Biotechnology, 2018, 12, 1013-1023.	1.7	28
102	Frontal and anterior cingulate activation during overt verbal fluency in patients with first episode psychosis. Revista Brasileira De Psiquiatria, 2005, 27, 228-232.	1.7	27
103	Well-being and help-seeking: an exploratory study among final-year medical students. Revista Da Associação Médica Brasileira, 2011, 57, 379-386.	0.7	27
104	Coping Style and Depressive Symptomatology during Pregnancy in a Private Setting Sample. Spanish Journal of Psychology, 2012, 15, 295-305.	2.1	27
105	What determines continuing grey matter changes in first-episode schizophrenia and affective psychosis?. Psychological Medicine, 2015, 45, 817-828.	4.5	27
106	Perception of negative emotions in close relatives by patients with schizophrenia. British Journal of Clinical Psychology, 2001, 40, 167-175.	3.5	26
107	The Prevalence of Headache Among Elderly in a Low-Income Area of São Paulo, Brazil. Cephalalgia, 2008, 28, 329-333.	3.9	26
108	Incidence and risk factors for osteoporotic vertebral fracture in low-income community-dwelling elderly: a population-based prospective cohort study in Brazil. The São Paulo Ageing & Health (SPAH) Study. Osteoporosis International, 2014, 25, 2805-2815.	3.1	26

#	Article	IF	CITATIONS
109	Urbanicity and risk of first-episode psychosis: incidence study in Brazil. British Journal of Psychiatry, 2019, 215, 726-729.	2.8	25
110	Prevalence of cannabis use in Brazil: Data from the I Brazilian National Alcohol Survey (BNAS). Addictive Behaviors, 2010, 35, 190-193.	3.0	24
111	Immunogenicity and safety of two doses of the CoronaVac SARS-CoV-2 vaccine in SARS-CoV-2 seropositive and seronegative patients with autoimmune rheumatic diseases in Brazil: a subgroup analysis of a phase 4 prospective study. Lancet Rheumatology, The, 2022, 4, e113-e124.	3.9	24
112	Lateral ventricle differences between first-episode schizophrenia and first-episode psychotic bipolar disorder: A population-based morphometric MRI study. World Journal of Biological Psychiatry, 2010, 11, 873-887.	2.6	23
113	Common mental disorders during pregnancy and adverse obstetric outcomes. Journal of Psychosomatic Obstetrics and Gynaecology, 2010, 31, 229-235.	2.1	23
114	Education and WHO Recommendations for Fruit and Vegetable Intake Are Associated with Better Cognitive Function in a Disadvantaged Brazilian Elderly Population: A Population-Based Cross-Sectional Study. PLoS ONE, 2014, 9, e94042.	2.5	23
115	Cognitive functioning throughout adulthood and illness stages in individuals with psychotic disorders and their unaffected siblings. Molecular Psychiatry, 2021, 26, 4529-4543.	7.9	23
116	Obsessive-Compulsive Symptoms, Obsessive-Compulsive Disorder, and Related Disorders in Parkinson's Disease. Journal of Neuropsychiatry and Clinical Neurosciences, 2003, 15, 371-374.	1.8	23
117	Mortality among patients with non-affective functional psychoses in a metropolitan area of South-Eastern Brazil. Revista De Saude Publica, 1996, 30, 304-309.	1.7	22
118	Chronic fatigue among bank workers in Brazil. Occupational Medicine, 2002, 52, 187-194.	1.4	22
119	Cardiovascular risk factors associated with migraine among the elderly with a low income: The São Paulo Ageing & Health Study (SPAH). Cephalalgia, 2011, 31, 331-337.	3.9	22
120	Prevalence of hepatitis B virus infection and carriage after nineteen years of vaccination program in the Western Brazilian Amazon. Revista Da Sociedade Brasileira De Medicina Tropical, 2012, 45, 13-17.	0.9	22
121	Visceral fat measured by DXA is associated with increased risk of non-spine fractures in nonobese elderly women: a population-based prospective cohort analysis from the SĂ£o Paulo Ageing & Health (SPAH) Study. Osteoporosis International, 2016, 27, 3525-3533.	3.1	22
122	Early-life stress effects on BDNF DNA methylation in first-episode psychosis and in rats reared in isolation. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2021, 108, 110188.	4.8	22
123	Migration history and risk of psychosis: results from the multinational EU-GEI study. Psychological Medicine, 2022, 52, 2972-2984.	4.5	22
124	Relationship between obsessive-compulsive disorders and diseases affecting primarily the basal ganglia. Revista Do Hospital Das Clinicas, 1999, 54, 213-221.	0.5	21
125	Family context and duration of untreated psychosis (DUP): Results from the Sao Paulo Study. Schizophrenia Research, 2010, 119, 124-130.	2.0	21
126	Relationship Between Regional Brain Volumes and Cognitive Performance in the Healthy Aging: An MRI Study Using Voxel-Based Morphometry. Journal of Alzheimer's Disease, 2012, 31, 45-58.	2.6	21

#	Article	IF	CITATIONS
127	Anemia and dementia among the elderly: the São Paulo Ageing & Health Study. International Psychogeriatrics, 2012, 24, 74-81.	1.0	21
128	The relationship between indicators of socioeconomic status and cesarean section in public hospitals. Revista De Saude Publica, 2017, 51, 14.	1.7	21
129	Prevalence of substance misuse among individuals with severe mental illness in S�o Paulo. Social Psychiatry and Psychiatric Epidemiology, 2004, 39, 212-217.	3.1	20
130	Subtle Gray Matter Changes in Temporo-Parietal Cortex Associated with Cardiovascular Risk Factors. Journal of Alzheimer's Disease, 2011, 27, 575-589.	2.6	20
131	Brain activity patterns during phonological verbal fluency performance with varying levels of difficulty: A functional magnetic resonance imaging study in Portuguese-speaking healthy individuals. Journal of Clinical and Experimental Neuropsychology, 2011, 33, 864-873.	1.3	20
132	Fibrilação atrial e demência: resultados do São Paulo ageing & health study. Arquivos Brasileiros De Cardiologia, 2012, 99, 1108-1114.	0.8	20
133	Intimate Partner Violence Is Associated with Suicidality Among Low-Income Postpartum Women. Journal of Women's Health, 2018, 27, 171-178.	3.3	20
134	Structural validity of the Maslach Burnout Inventory and influence of depressive symptoms in banking workplace: Unfastening the occupational conundrum. Psychiatry Research, 2018, 267, 168-174.	3.3	20
135	Addressing Depression Comorbid With Diabetes or Hypertension in Resource-Poor Settings: A Qualitative Study About User Perception of a Nurse-Supported Smartphone App in Peru. JMIR Mental Health, 2019, 6, e11701.	3.3	20
136	Cognitive functioning in subjects with recent-onset psychosis from a low-middle-income environment: Multiple-domain deficits and longitudinal evaluation. Psychiatry Research, 2010, 179, 157-164.	3.3	19
137	Work-Related Depression in Primary Care Teams in Brazil. American Journal of Public Health, 2016, 106, 1990-1997.	2.7	19
138	Nutritional Risk and Quality of Life in Community-Dwelling Elderly: A Cross-Sectional Study. Journal of Nutrition, Health and Aging, 2018, 22, 111-116.	3.3	19
139	Public Stigma towards Older Adults with Depression: Findings from the São Paulo-Manaus Elderly in Primary Care Study. PLoS ONE, 2016, 11, e0157719.	2.5	19
140	Prevalência de Achados eletrocardiográficos no paciente idoso: estudo envelhecimento e saúde de São Paulo. Arquivos Brasileiros De Cardiologia, 2009, 93, 651-656.	0.8	18
141	Volume reduction of the corpus callosum and its relationship with deficits in interhemispheric transfer of information in recent-onset psychosis. Psychiatry Research - Neuroimaging, 2010, 184, 1-9.	1.8	18
142	The influence of lifestyle and gender on sickness absence in Brazilian workers. BMC Public Health, 2014, 14, 317.	2.9	18
143	Depressive symptoms and psychosocial aspects of work in bank employees. Occupational Medicine, 2016, 66, 54-61.	1.4	18
144	Lower education among low-income Brazilian adolescent females is associated with planned pregnancies. International Journal of Women's Health, 2017, Volume 9, 43-48.	2.6	18

#	Article	IF	CITATIONS
145	Recognition of Depressive Symptoms by Physicians. Clinics, 2009, 64, 629-635.	1.5	17
146	Loss of interest, depressed mood and impact on the quality of life: Cross-sectional survey. BMC Public Health, 2011, 11, 826.	2.9	17
147	Cardiovascular risk factors in patients with first-episode psychosis in São Paulo, Brazil. General Hospital Psychiatry, 2012, 34, 268-275.	2.4	17
148	The association between depressive disorders and health care utilization: results from the São Paulo Ageing and Health Study (SPAH). General Hospital Psychiatry, 2014, 36, 199-202.	2.4	17
149	The Independent Effects of Psychosocial Stressors on Subclinical Psychosis: Findings From the Multinational EU-GEI Study. Schizophrenia Bulletin, 2021, 47, 1674-1684.	4.3	17
150	Feasibility and reliability of the elderly version of the Camberwell Assessment of Needs (CANE): results from the São Paulo Ageing & Health Study. Revista Brasileira De Psiquiatria, 2009, 31, 34-38.	1.7	16
151	Osteoporotic Fractures in the Brazilian Community-Dwelling Elderly: Prevalence and Risk Factors. Journal of Clinical Densitometry, 2011, 14, 359-366.	1.2	16
152	Voxelwise evaluation of white matter volumes in first-episode psychosis. Psychiatry Research - Neuroimaging, 2012, 202, 198-205.	1.8	16
153	Building Research Capacity Across and Within Low- and Middle-Income Countries: The Collaborative Hubs for International Research on Mental Health. Academic Psychiatry, 2016, 40, 686-691.	0.9	16
154	High frequency of silent brain infarcts associated with cognitive deficits in an economically disadvantaged population. Clinics, 2017, 72, 474-480.	1.5	16
155	Type of delivery is not associated with maternal depression. Archives of Women's Mental Health, 2019, 22, 631-635.	2.6	16
156	The social adjustment of patients with schizophrenia: implications to the mental health policy in Brazil. Revista De Saude Publica, 1993, 27, 340-349.	1.7	15
157	Investigating the role played by social support in the association between religiosity and mental health in low income older adults: results from the São Paulo Ageing & Health Study (SPAH). Revista Brasileira De Psiquiatria, 2011, 33, 157-164.	1.7	15
158	Intimate partner violence among undergraduate students of two universities of the state of São Paulo, Brazil. Revista Brasileira De Epidemiologia, 2013, 16, 801-816.	0.8	15
159	Low plasma concentrations of N-methyl-d-aspartate receptor subunits as a possible biomarker for psychosis. Schizophrenia Research, 2018, 202, 55-63.	2.0	15
160	Prevalence of Depression and Depression Care for Populations Registered in Primary Care in Two Remote Cities in the Brazilian Amazon. PLoS ONE, 2016, 11, e0150046.	2.5	15
161	Cardiovascular risk in cognitively preserved elderlies is associated with glucose hypometabolism in the posterior cingulate cortex and precuneus regardless of brain atrophy and apolipoprotein gene variations. Age, 2013, 35, 777-792.	3.0	14
162	Corpus callosum volumes in the 5†years following the first-episode of schizophrenia: Effects of antipsychotics, chronicity and maturation. NeuroImage: Clinical, 2018, 18, 932-942.	2.7	14

#	Article	IF	CITATIONS
163	Premorbid Adjustment and IQ in Patients With First-Episode Psychosis: A Multisite Case-Control Study of Their Relationship With Cannabis Use. Schizophrenia Bulletin, 2020, 46, 517-529.	4.3	14
164	The awareness of chronic fatigue syndrome: A comparative study in Brazil and the United Kingdom. Journal of Psychosomatic Research, 2008, 64, 351-355.	2.6	13
165	Monitoring self-reported adherence to antiretroviral therapy in public HIV care facilities in Brazil. Medicine (United States), 2018, 97, S38-S45.	1.0	13
166	Prolonged Periods of Social Isolation From Weaning Reduce the Anti-inflammatory Cytokine IL-10 in Blood and Brain. Frontiers in Neuroscience, 2018, 12, 1011.	2.8	13
167	Subjective Memory Deficits in People with and without Dementia: Findings from the 10/66 Dementia Research Group Pilot Studies in Low―and Middleâ€Income Countries. Journal of the American Geriatrics Society, 2009, 57, 2118-2124.	2.6	12
168	The relationship between perinatal psychiatric disorders and contraception use among postpartum women. Contraception, 2013, 88, 498-502.	1.5	12
169	Identification and treatment of depression of older adults in primary care: findings from the São Paulo Ageing and Health Study. Family Practice, 2016, 33, 233-237.	1.9	12
170	The continuity of effect of schizophrenia polygenic risk score and patterns of cannabis use on transdiagnostic symptom dimensions at first-episode psychosis: findings from the EU-GEI study. Translational Psychiatry, 2021, 11, 423.	4.8	12
171	Reliability of the Brazilian version of the Camberwell Assessment of Needs (CAN) in first-episode psychosis cases in São Paulo, Brazil. Revista Brasileira De Psiquiatria, 2007, 29, 160-163.	1.7	11
172	Associations between OPG and RANKL polymorphisms, vertebral fractures, and abdominal aortic calcification in community-dwelling older subjects: the Sao Paulo Ageing & Health Study (SPAH). Osteoporosis International, 2016, 27, 3319-3329.	3.1	11
173	A study of the incidence and recognition of surgical glove perforation during obstetric and gynecological procedures. Archives of Gynecology and Obstetrics, 2004, 270, 263-264.	1.7	10
174	Early Parental Death and Risk of Psychosis in Offspring: A Six-Country Case-Control Study. Journal of Clinical Medicine, 2019, 8, 1081.	2.4	10
175	The independent and combined effects of cannabis use and systemic inflammation during the early stages of psychosis: exploring the two-hit hypothesis. Psychological Medicine, 2022, 52, 3874-3884.	4.5	10
176	Synergistic effects of childhood adversity and polygenic risk in first-episode psychosis: the EU-GEI study. Psychological Medicine, 0, , 1-9.	4.5	10
177	Corpus callosum volumes in recent-onset schizophrenia are correlated to positive symptom severity after 1year of follow-up. Schizophrenia Research, 2012, 137, 258-259.	2.0	9
178	The relationship between sleep disturbance in pregnancy and persistent common mental disorder in the perinatal period (sleep disturbance and persistent CMD). Journal of Mental Health, 2015, 24, 375-378.	1.9	9
179	Ethnic differences in tobacco use during pregnancy: findings from a primary care sample in São Paulo, Brazil. Ethnicity and Health, 2015, 20, 209-217.	2.5	9
180	A Method for Simultaneous Evaluation of Muscular and Neural Prepulse Inhibition. Frontiers in Neuroscience, 2018, 12, 654.	2.8	9

#	Article	IF	CITATIONS
181	Childhood Maltreatment, Educational Attainment, and IQ: Findings From a Multicentric Case-control Study of First-episode Psychosis (EU-GEI). Schizophrenia Bulletin, 2022, 48, 575-589.	4.3	9
182	Facial Emotion Recognition in Psychosis and Associations With Polygenic Risk for Schizophrenia: Findings From the Multi-Center EU-GEI Case–Control Study. Schizophrenia Bulletin, 2022, 48, 1104-1114.	4.3	9
183	Causes of recurrent or persistent anemia in older people from the results of the São Paulo Ageing & Health Study. Geriatrics and Gerontology International, 2013, 13, 204-208.	1.5	8
184	Management of depression during pregnancy in primary care in Brazil: a cluster ransomised trial. Lancet, The, 2014, 384, S21.	13.7	8
185	Epigenetic-mediated <i>N</i> -methyl-D-aspartate receptor changes in the brain of isolated reared rats. Epigenomics, 2020, 12, 1983-1997.	2.1	8
186	PrognÃ ³ stico da esquizofrenia. Revista Brasileira De Psiquiatria, 2000, 22, 18-20.	1.7	8
187	Characteristics of hospital-treated schizophrenia in Sĩز½o Paulo, Brazil. Social Psychiatry and Psychiatric Epidemiology, 1993, 28, 267-274.	3.1	7
188	BDNF Gene Polymorphism, Cognition and Symptom Severity in a Brazilian Population-Based Sample of First-Episode Psychosis Subjects. Revista Brasileira De Psiquiatria, 2012, 34, 219-232.	1.7	7
189	Framingham Coronary Heart Disease Risk Score Can be Predicted from Structural Brain Images in Elderly Subjects. Frontiers in Aging Neuroscience, 2014, 6, 300.	3.4	7
190	Psychosocial Work Conditions and Burnout Among Brazilian Bank Employees: A Cross-Sectional Study. Annals of Occupational Hygiene, 2016, 60, 567-580.	1.9	7
191	Plasma amino acids profile in first-episode psychosis, unaffected siblings and community-based controls. Scientific Reports, 2020, 10, 21423.	3.3	7
192	Recruitment, training and supervision of nurses and nurse assistants for a task-shifting depression intervention in two RCTs in Brazil and Peru. Human Resources for Health, 2021, 19, 16.	3.1	7
193	Perceived major experiences of discrimination, ethnic group, and risk of psychosis in a six-country caseâ°'control study. Psychological Medicine, 2022, 52, 3668-3676.	4.5	7
194	Burnout among primary health care workers in Brazil: results of a multilevel analysis. International Archives of Occupational and Environmental Health, 2021, 94, 1863-1875.	2.3	7
195	Validation of the Portuguese version of the Community Assessment of Psychic Experiences and characterization of psychotic experiences in a Brazilian sample. Revista Brasileira De Psiquiatria, 2020, 42, 389-397.	1.7	7
196	Prevalence of tobacco use in individuals with severe mental illnesses, São Paulo, Brazil. Revista De Saude Publica, 2007, 41, 510-516.	1.7	6
197	Commentary: Epidemiological mental health research: contribution from low- and middle-income countries is essential. International Journal of Epidemiology, 2014, 43, 301-303.	1.9	6
198	Persistent hypovitaminosis D and loss of hip bone mineral density over time as additional risk factors for recurrent falls in a population-based prospective cohort of elderly persons living in the community. The São Paulo Ageing & Health (SPAH) Study. Osteoporosis International, 2015, 26, 1535-1542.	3.1	6

#	Article	IF	CITATIONS
199	Pre-training inter-rater reliability of clinical instruments in an international psychosis research project. Schizophrenia Research, 2020, 230, 104-107.	2.0	6
200	Risk Factors for Death Among 120,804 Hospitalized Patients with Confirmed COVID-19 in São Paulo, Brazil. American Journal of Tropical Medicine and Hygiene, 2021, , .	1.4	6
201	Early Schizophrenia and Bipolar Disorder Patients Display Reduced Neural Prepulse Inhibition. Brain Sciences, 2022, 12, 93.	2.3	6
202	Trend in the prevalence of depressive symptoms in Brazil: results from the Brazilian National Health Survey 2013 and 2019. Cadernos De Saude Publica, 2022, 38, e00123421.	1.0	6
203	The relationship of childhood trauma and DNA methylation of NMDA receptor genes in first-episode schizophrenia. Epigenomics, 2021, 13, 927-937.	2.1	5
204	Excess mortality in a cohort of Brazilian patients with a median follow-up of 11Âyears after the first psychiatric hospital admission. Social Psychiatry and Psychiatric Epidemiology, 2023, 58, 319-330.	3.1	5
205	Longitudinal brain volumetric changes during one year in non-elderly healthy adults: a voxel-based morphometry study. Brazilian Journal of Medical and Biological Research, 2012, 45, 516-523.	1.5	4
206	Duration of Untreated Psychosis in First-Episode Psychosis is not Associated With Common Genetic Variants for Major Psychiatric Conditions: Results From the Multi-Center EU-GEI Study. Schizophrenia Bulletin, 2021, 47, 1653-1662.	4.3	4
207	The relationship of symptom dimensions with premorbid adjustment and cognitive characteristics at first episode psychosis: Findings from the EU-GEI study. Schizophrenia Research, 2021, 236, 69-79.	2.0	4
208	Frequência de sintomas no último ano de vida de idosos de baixa renda em São Paulo: estudo transversal com cuidadores informais. Revista Brasileira De Epidemiologia, 2011, 14, 75-85.	0.8	3
209	Prefrontal-Parietal White Matter Volumes in Healthy Elderlies Are Decreased in Proportion to the Degree of Cardiovascular Risk and Related to Inhibitory Control Deficits. Frontiers in Psychology, 2017, 8, 57.	2.1	3
210	Risk Factors for Low Muscle Mass in a Population-based Prospective Cohort of Brazilian Community-dwelling Older Women: The São Paulo Ageing & Health (SPAH) Study. Journal of Clinical Densitometry, 2020, 23, 503-510.	1.2	3
211	The social adjustment of people with severe mental illness in São Paulo, Brazil. Revista Brasileira De Psiquiatria, 2005, 27, 309-314.	1.7	2
212	Simultaneous evaluation of prepulse inhibition with EMG and EEG using advanced artifact removal techniques. , 2016, 2016, 5262-5265.		2
213	From where we came: Absence of internal migration effect on psychosis in two case-control Brazilian samples. Schizophrenia Research, 2019, 212, 241-242.	2.0	2
214	Prevalence of Older Adult Disability and Primary Health Care Responsiveness in Low-Income Communities. Life, 2020, 10, 133.	2.4	2
215	KLOTHO polymorphisms and age-related outcomes in community-dwelling older subjects: TheÂSão Paulo Ageing & Health (SPAH) Study. Scientific Reports, 2020, 10, 8574.	3.3	2
216	Regional gray matter reductions in elderly subjects with cardiovascular risk factors: A possible link between microvascular lesions and Alzheimer's disease. Journal of the Neurological Sciences, 2009, 283, 311-312.	0.6	1

#	Article	IF	CITATIONS
217	P2-541 The impact of depression and depressive symptoms on health service use: the Sao Paulo Ageing & Health Study (SPAH). Journal of Epidemiology and Community Health, 2011, 65, A370-A370.	3.7	1
218	Twoâ€Year Anemia Incidence and Causes in Older Individuals with Nearâ€Abnormal Baseline Hemoglobin Levels: The São Paulo Ageing and Health Study. Journal of the American Geriatrics Society, 2013, 61, 1419-1421.	2.6	1
219	Depression and violent crime: is there a relationship?. Evidence-Based Mental Health, 2015, 18, 114-114.	4.5	1
220	Prevalence and risk factors for cannabis use in low-income pregnant women in São Paulo, Brazil. Archives of Women's Mental Health, 2016, 19, 193-196.	2.6	1
221	S14. DNA METHYLATION CHANGES IN GABAERGIC AND GLUTAMATERGIC MARKERS IN EARLY SCHIZOPHRENIA. Schizophrenia Bulletin, 2018, 44, S329-S329.	4.3	1
222	Patient needs four years after first psychiatric hospitalization in a Brazilian cohort. Brazilian Journal of Medical and Biological Research, 2021, 54, e11447.	1.5	1
223	Plasma prevalence of anti-N-methyl-d-aspartate receptor IgG antibodies in early stages of psychosis. Ciencia E Saude Coletiva, 2021, 26, 1085-1094.	0.5	1
224	Study had methodological limitations. BMJ: British Medical Journal, 1996, 312, 512-513.	2.3	1
225	Sexual activity among female teenagers: a comparison between two groups of middle class adolescents from a private clinic according to pregnancy status. Revista Brasileira De Saude Materno Infantil, 2008, 8, 251-256.	0.5	1
226	Computerized psychiatric interviews: can they help to improve medical care?. Sao Paulo Medical Journal, 1998, 116, 1821-1822.	0.9	0
227	Evaluation of abnormal hippocampal volumes using voxel-based morphometry (VBM): Impact of variable smoothing filters in a study using elderly as a model. International Journal of Psychophysiology, 2010, 77, 263-263.	1.0	0
228	P2-509 Population impact of depression on functional disability in the elderly: results from the "SAo Paulo Ageing & Health Study" (SPAH). Journal of Epidemiology and Community Health, 2011, 65, A361-A362.	3.7	0
229	P2-527 Prevalence of depression in the Brazilian family healthcare strategy: a cross-sectional study in SAo Paulo and Manaus. Journal of Epidemiology and Community Health, 2011, 65, A366-A367.	3.7	0
230	Poster #42 CANNABIS EFFECTS ON BRAIN STRUCTURE IN FIRST-EPISODE PSYCHOSIS. Schizophrenia Research, 2012, 136, S200.	2.0	0
231	Poster #43 STRUCTURAL BRAIN CHANGES IN FIRST-EPISODE SCHIZOPHRENIA: A 4-5 YEAR FOLLOW-UP STUDY USING MRI. Schizophrenia Research, 2012, 136, S201.	2.0	0
232	Perinatal common mental disorders and breastfeeding duration: A cohort study from Brazil. Journal of Neonatal-Perinatal Medicine, 2012, 5, 135-142.	0.8	0
233	The relationship between socioeconomic indicators during pregnancy and gynecological appointment at any time after childbirth. International Journal for Equity in Health, 2015, 14, 64.	3.5	0
234	Re: Response to â€~Depressive symptoms and psychosocial aspects of work in bank employees'. Occupational Medicine, 2016, 66, 421.2-422.	1.4	0

0

#	Article	IF	CITATIONS
235	T131. INCIDENCE OF FIRST-EPISODE PSYCHOSIS IN THE CATCHMENT AREA OF RIBEIRÃO PRETO, BRAZIL. Schizophrenia Bulletin, 2018, 44, S166-S166.	4.3	0
236	Estado nutricional de idosos residentes nos municÃpios da Superintendência Regional de Saúde de Uberaba. , 2019, 98, 290-297.	0.1	0
237	Creación del Colegio Doctoral Tordesillas de Salud Pública e Historia de la Ciencia: un modelo de internacionalización académica. Educacion Medica, 2020, 21, 223-229.	0.3	0
238	S127. MORTALITY IN PSYCHIATRIC PATIENTS WITH AT LEAST NINE-YEAR FOLLOW-UP PERIOD AFTER THEIR FIRST HOSPITAL ADMISSION: A BRAZILIAN COHORT STUDY. Schizophrenia Bulletin, 2020, 46, S83-S84.	4.3	0
239	S128. IMPACT OF COMORBIDITIES ON SURVIVAL ESTIMATE FOR PATIENTS WITH MENTAL DISORDERS. Schizophrenia Bulletin, 2020, 46, S84-S84.	4.3	0
240	Cannabis Consumption Increases Risk of Psychosis in a Subgroup of Patients With High Peripheral Blood Inflammation. Biological Psychiatry, 2020, 87, S259.	1.3	0
241	Signals of Adaptive Immune Activation in Non-Clinical Populations With Psychotic Experiences. Biological Psychiatry, 2021, 89, S191.	1.3	0
242	Social and clinical features associated with duration of untreated psychosis in the Brazilian STREAM study. Asian Journal of Psychiatry, 2021, 65, 102823.	2.0	0
243	A Mobile Health Intervention for Patients With Depressive Symptoms: Protocol for an Economic Evaluation Alongside Two Randomized Trials in Brazil and Peru. JMIR Research Protocols, 2021, 10, e26164.	1.0	0

Incidence and Outcome of Schizophrenia Across the Globe. , 2010, , 3-17.