

Young-Min Park

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6366298/publications.pdf>

Version: 2024-02-01

22
papers

402
citations

758635

12
h-index

887659

17
g-index

22
all docs

22
docs citations

22
times ranked

651
citing authors

#	ARTICLE	IF	CITATIONS
1	Physical Activity Differentially Affects the Cecal Microbiota of Ovariectomized Female Rats Selectively Bred for High and Low Aerobic Capacity. <i>PLoS ONE</i> , 2015, 10, e0136150.	1.1	64
2	Female rats selectively bred for high intrinsic aerobic fitness are protected from ovariectomy-associated metabolic dysfunction. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2015, 308, R530-R542.	0.9	44
3	Soy Improves Cardiometabolic Health and Cecal Microbiota in Female Low-Fit Rats. <i>Scientific Reports</i> , 2017, 7, 9261.	1.6	43
4	Time since menopause and skeletal muscle estrogen receptors, PGC-1 α , and AMPK. <i>Menopause</i> , 2017, 24, 815-823.	0.8	34
5	Moderate amounts of fructose- or glucose-sweetened beverages do not differentially alter metabolic health in male and female adolescents , ,. <i>American Journal of Clinical Nutrition</i> , 2014, 100, 796-805.	2.2	30
6	Effects of intrinsic aerobic capacity and ovariectomy on voluntary wheel running and nucleus accumbens dopamine receptor gene expression. <i>Physiology and Behavior</i> , 2016, 164, 383-389.	1.0	30
7	Retention of sedentary obese visceral white adipose tissue phenotype with intermittent physical activity despite reduced adiposity. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2015, 309, R594-R602.	0.9	28
8	Acute estradiol treatment reduces skeletal muscle protein breakdown markers in early- but not late-postmenopausal women. <i>Steroids</i> , 2019, 146, 43-49.	0.8	22
9	Effects of ovariectomy and intrinsic aerobic capacity on tissue-specific insulin sensitivity. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2016, 310, E190-E199.	1.8	21
10	Appendicular lean mass is lower in late compared with early perimenopausal women: potential role of FSH. <i>Journal of Applied Physiology</i> , 2020, 128, 1373-1380.	1.2	19
11	Voluntary Running Attenuates Metabolic Dysfunction in Ovariectomized Low-Fit Rats. <i>Medicine and Science in Sports and Exercise</i> , 2017, 49, 254-264.	0.2	17
12	Ovariectomized Highly Fit Rats Are Protected against Diet-Induced Insulin Resistance. <i>Medicine and Science in Sports and Exercise</i> , 2016, 48, 1259-1269.	0.2	12
13	Prior exercise does not alter the incretin response to a subsequent meal in obese women. <i>Peptides</i> , 2015, 71, 94-99.	1.2	11
14	Voluntary wheel running improves adipose tissue immunometabolism in ovariectomized low-fit rats. <i>Adipocyte</i> , 2018, 7, 20-34.	1.3	10
15	Bone Mineral Density in Different Menopause Stages is Associated with Follicle Stimulating Hormone Levels in Healthy Women. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 1200.	1.2	9
16	Lessons Learned in Protein Precipitation Using a Membrane Emulsification Technique to Produce Reversible and Uniform Microbeads. <i>Pharmaceutics</i> , 2021, 13, 1738.	2.0	5
17	Menopause and the Loss of Skeletal Muscle Mass in Women. <i>Iranian Journal of Public Health</i> , 2021, 50, 413-414.	0.3	3
18	Chronic Ankle Instability Research and Cross-Cultural Adaptation Process Necessity for the Self-Reported Questionnaire. <i>Iranian Journal of Public Health</i> , 2021, 50, 838-839.	0.3	0

#	ARTICLE	IF	CITATIONS
19	Low intrinsic aerobic fitness increases susceptibility to OVX-induced obesity and insulin resistance in the absence of adipose tissue inflammation (1028.3). FASEB Journal, 2014, 28, 1028.3.	0.2	0
20	Intermittent Physical Activity Produces a Leaner but "Sedentary Obese" White Adipose Tissue Phenotype. FASEB Journal, 2015, 29, 1055.16.	0.2	0
21	Appendicular Muscle Mass is Lower in Late-Compared to Early-Perimenopausal Women. FASEB Journal, 2018, 32, lb397.	0.2	0
22	Endothelial Estrogen Receptor α/β Protein Ratio is Associated with Circulating Estrogen Levels in Healthy Women. FASEB Journal, 2019, 33, lb513.	0.2	0