

Jana HansÃ-kovÃ;

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6370326/publications.pdf>

Version: 2024-02-01

12
papers

315
citations

1040056

9
h-index

1199594

12
g-index

13
all docs

13
docs citations

13
times ranked

587
citing authors

#	ARTICLE	IF	CITATIONS
1	Postnatal induction of muscle fatty acid oxidation in mice differing in propensity to obesity: a role of pyruvate dehydrogenase. <i>International Journal of Obesity</i> , 2020, 44, 235-244.	3.4	6
2	Chronic n-3 fatty acid intake enhances insulin response to oral glucose and elevates GLP-1 in high-fat diet-fed obese mice. <i>Food and Function</i> , 2020, 11, 9764-9775.	4.6	9
3	Dysregulation of epicardial adipose tissue in cachexia due to heart failure: the role of natriuretic peptides and cardiolipin. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2020, 11, 1614-1627.	7.3	24
4	Triacylglycerol-Rich Oils of Marine Origin are Optimal Nutrients for Induction of Polyunsaturated Docosahexaenoic Acid Ester of Hydroxy Linoleic Acid (13-HDHAHLA) with Anti-Inflammatory Properties in Mice. <i>Molecular Nutrition and Food Research</i> , 2020, 64, e1901238.	3.3	26
5	Increased plasma levels of palmitoleic acid may contribute to beneficial effects of Krill oil on glucose homeostasis in dietary obese mice. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2020, 1865, 158732.	2.4	12
6	Metformin acutely lowers blood glucose levels by inhibition of intestinal glucose transport. <i>Scientific Reports</i> , 2019, 9, 6156.	3.3	78
7	Differential modulation of white adipose tissue endocannabinoid levels by n-3 fatty acids in obese mice and type 2 diabetic patients. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2018, 1863, 712-725.	2.4	22
8	Levels of palmitic acid ester of hydroxystearic acid (PAHSA) are reduced in the breast milk of obese mothers. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2018, 1863, 126-131.	2.4	53
9	Induction of lipogenesis in white fat during cold exposure in mice: link to lean phenotype. <i>International Journal of Obesity</i> , 2017, 41, 372-380.	3.4	38
10	Early metabolic differences between obesity-resistant and obesity-prone mice: role of adipokines. <i>Proceedings of the Nutrition Society</i> , 2016, 75, .	1.0	0
11	Early differences in metabolic flexibility between obesity-resistant and obesity-prone mice. <i>Biochimie</i> , 2016, 124, 163-170.	2.6	13
12	Plasma Acylcarnitines and Amino Acid Levels As an Early Complex Biomarker of Propensity to High-Fat Diet-Induced Obesity in Mice. <i>PLoS ONE</i> , 2016, 11, e0155776.	2.5	13