Rikke M Zachar

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6370499/publications.pdf Version: 2024-02-01



RIKKE M ZACHAR

#	Article	IF	CITATIONS
1	The Epithelial Sodium Channel Î ³ -Subunit Is Processed Proteolytically in Human Kidney. Journal of the American Society of Nephrology: JASN, 2015, 26, 95-106.	3.0	55
2	Physiology and pathophysiology of the plasminogen system in the kidney. Pflugers Archiv European Journal of Physiology, 2017, 469, 1415-1423.	1.3	34
3	Urine exosomes from healthy and hypertensive pregnancies display elevated level of α-subunit and cleaved α- and γ-subunits of the epithelial sodium channel—ENaC. Pflugers Archiv European Journal of Physiology, 2017, 469, 1107-1119.	1.3	28
4	Albuminuria in kidney transplant recipients is associated with increased urinary serine proteases and activation of the epithelial sodium channel. American Journal of Physiology - Renal Physiology, 2018, 315, F151-F160.	1.3	26
5	Dietary Na+ intake in healthy humans changes the urine extracellular vesicle prostasin abundance while the vesicle excretion rate, NCC, and ENaC are not altered. American Journal of Physiology - Renal Physiology, 2019, 317, F1612-F1622.	1.3	12
6	The epithelial Na+ channel α- and γ-subunits are cleaved at predicted furin-cleavage sites, glycosylated and membrane associated in human kidney. Pflugers Archiv European Journal of Physiology, 2019, 471, 1383-1396.	1.3	10
7	Proteinuria is accompanied by intratubular complement activation and apical membrane deposition of C3dg and C5b-9 in kidney transplant recipients. American Journal of Physiology - Renal Physiology, 2022, 322, F150-F163.	1.3	9
8	Hydronephrosis is associated with elevated plasmin in urine in pediatric patients and rats and changes in NCC and γ-ENaC abundance in rat kidney. American Journal of Physiology - Renal Physiology, 2018, 315, F547-F557.	1.3	5