Chia-Ying Li

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/6376664/chia-ying-li-publications-by-year.pdf

Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

36	919	19	30
papers	citations	h-index	g-index
48	1,022	3.5	3.53
ext. papers	ext. citations	avg, IF	L-index

#	Paper	IF	Citations
36	Genomes of leafy and leafless Platanthera orchids illuminate the evolution of mycoheterotrophy <i>Nature Plants</i> , 2022 , 8, 373-388	11.5	4
35	Natural Cembrane Diterpenoids From the Soft Coral Sinularia querciformis. <i>Natural Product Communications</i> , 2021 , 16, 1934578X2110592	0.9	1
34	A Rapid and Feasible H-NMR Quantification Method of Ephedrine Alkaloids in Herbal Preparations. <i>Molecules</i> , 2021 , 26,	4.8	2
33	Direct synthesis of monolayer gold nanoparticles on epoxy based photoresist by photoreduction and application to surface-enhanced Raman sensing. <i>Materials and Design</i> , 2021 , 197, 109211	8.1	5
32	Tyrosinase Inhibitory Effect, Antioxidant and Anticancer Activities of Bioactive Compounds in Ripe Hog Plum (Spondias Pinnata) Fruit Extracts. <i>Oriental Journal of Chemistry</i> , 2019 , 35, 916-926	0.8	3
31	Separation and identification of a novel tadalafil analogue adulterant in a dietary supplement. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment</i> , 2016 , 33, 179-85	3.2	5
30	Cytotoxic phenanthroindolizidine alkaloids from the roots of Ficus septica. <i>Planta Medica</i> , 2009 , 75, 11	52 5. 6	21
29	A rapid and simple determination of protoberberine alkaloids in Rhizoma Coptidis by 1H NMR and its application for quality control of commercial prescriptions. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2009 , 49, 1272-6	3.5	45
28	Quality assessment of Radix Codonopsis by quantitative nuclear magnetic resonance. <i>Journal of Chromatography A</i> , 2009 , 1216, 2124-9	4.5	43
27	A novel homodimeric geranyl diphosphate synthase from the orchid Phalaenopsis bellina lacking a DD(X)2-4D motif. <i>Plant Journal</i> , 2008 , 55, 719-33	6.9	74
26	New diterpenoids and the bioactivity of Erythrophleum fordii. <i>Bioorganic and Medicinal Chemistry</i> , 2008 , 16, 9867-70	3.4	25
25	Isolation and identification of antiplatelet aggregatory principles from the leaves of Piper lolot. <i>Journal of Agricultural and Food Chemistry</i> , 2007 , 55, 9436-42	5.7	21
24	High-performance liquid chromatographic method for simultaneous quantification of eight major biologically active ingredients in XDa-Chai-Hu-TangXpreparation. <i>Biomedical Chromatography</i> , 2006 , 20, 305-8	1.7	14
23	Delayed treatment with nicotinamide inhibits brain energy depletion, improves cerebral microperfusion, reduces brain infarct volume, but does not alter neurobehavioral outcome following permanent focal cerebral ischemia in Sprague Dawley rats. <i>Current Neurovascular</i>	1.8	21
22	Research, 2006, 3, 203-13 Antidepressant principles of the roots of Polygala tenuifolia. <i>Journal of Natural Products</i> , 2006, 69, 130	5 z₽ 9	38
21	Constituents from the leaves of Phellodendron amurense and their antioxidant activity. <i>Chemical and Pharmaceutical Bulletin</i> , 2006 , 54, 1308-11	1.9	15
20	A rapid and simple determination of protoberberine alkaloids in cortex phellodendri by 1H NMR and its application for quality control of commercial traditional Chinese medicine prescriptions. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2006 , 40, 173-8	3.5	52

(1999-2005)

19	Phenanthroindolizidine alkaloids from the stems of Ficus septica. <i>Journal of Natural Products</i> , 2005 , 68, 1071-5	4.9	66	
18	The epimerization of sesamin and asarinin. <i>Journal of Natural Products</i> , 2005 , 68, 1622-4	4.9	36	
17	Constituents of leaves of Phellodendron japonicum MAXIM. and their antioxidant activity. <i>Chemical and Pharmaceutical Bulletin</i> , 2005 , 53, 1118-21	1.9	16	
16	Quantitative analysis of camptothecin derivatives in Nothapodytes foetida using 1H-NMR method. <i>Chemical and Pharmaceutical Bulletin</i> , 2005 , 53, 347-9	1.9	30	
15	Anthraquinones from Ophiorrhiza hayatana OHWI. Chemical and Pharmaceutical Bulletin, 2005, 53, 123	1 2-15 9	9	
14	Flavonoids and coumarins from Leaves of Phellodendron chinense. <i>Planta Medica</i> , 2004 , 70, 183-5	3.1	11	
13	Efficient 1H nuclear magnetic resonance method for improved quality control analyses of Ginkgo constituents. <i>Journal of Agricultural and Food Chemistry</i> , 2004 , 52, 3721-5	5.7	43	
12	Antityrosinase principles and constituents of the petals of Crocus sativus. <i>Journal of Natural Products</i> , 2004 , 67, 437-40	4.9	56	
11	A novel phenylpropenoyl sulfonic acid and a new chlorophyll from the leaves of Petasites formosanus KITAMURA. <i>Chemical and Pharmaceutical Bulletin</i> , 2004 , 52, 1151-2	1.9	13	
10	Constituents from the leaves of Phellodendron amurense var. wilsonii and their bioactivity. <i>Journal of Natural Products</i> , 2003 , 66, 1207-11	4.9	39	
9	Constituents of Leaves of Phellodendron chinense var. glabriusculum. Heterocycles, 2003, 60, 397	0.8	14	
8	Constituents of the Leaves of Petasites formosanus and Their Antioxidative Activity. <i>Heterocycles</i> , 2003 , 60, 1881	0.8	5	
7	Non-Alkaloidal Constituents from the Stem of Ficus Septica. <i>Journal of the Chinese Chemical Society</i> , 2002 , 49, 113-116	1.5	16	
6	Constituents of the pollen of Crocus sativus L. and their tyrosinase inhibitory activity. <i>Chemical and Pharmaceutical Bulletin</i> , 2002 , 50, 1305-9	1.9	26	
5	Constituents of the stigmas of Crocus sativus and their tyrosinase inhibitory activity. <i>Journal of Natural Products</i> , 2002 , 65, 1452-6	4.9	33	
4	Limonoids and alkaloids of the root bark of Dictamnus angustifolius. <i>Phytochemistry</i> , 1999 , 50, 509-512	2 4	49	
3	Four aristolochic acid esters of rearranged ent-elemane sesquiterpenes from aristolochia heterophylla. <i>Journal of Natural Products</i> , 1999 , 62, 348-51	4.9	13	
2	The bakkenolides from the root of Petasites formosanus and their cytotoxicity. <i>Chemical and Pharmaceutical Bulletin</i> , 1999 , 47, 375-82	1.9	46	

Severibuxine, a new quinolin-2,4-dione and other constituents from Severinia buxifolia. *Phytochemistry*, **1998**, 49, 1467-1470

4 9