
LÃ;szlÃ³ Kalabay

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6381272/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Correlation of maternal serum fetuin/alpha2-HS-glycoprotein concentration with maternal insulin resistance and anthropometric parameters of neonates in normal pregnancy and gestational diabetes. European Journal of Endocrinology, 2002, 147, 243-248.	3.7	71
2	Human fetuin/α2HS-glycoprotein level as a novel indicator of liver cell function and short-term mortality in patients with liver cirrhosis and liver cancer. European Journal of Gastroenterology and Hepatology, 2002, 14, 389-394.	1.6	67
3	Association of increased serum heat shock protein 70 and C-reactive protein concentrations and decreased serum α 2 -HS glycoprotein concentration with the syndrome of hemolysis, elevated liver enzymes, and low platelet count. Journal of Reproductive Immunology, 2007, 73, 172-179.	1.9	50
4	Affective temperament, history of suicide attempt and family history of suicide in general practice patients. Journal of Affective Disorders, 2013, 149, 350-354.	4.1	46
5	Helicobacter pylori Infection in Connective Tissue Disorders is Associated with High Levels of Antibodies to Mycobacterial hsp65 but not to Human hsp60. Helicobacter, 2002, 7, 250-256.	3.5	45
6	Serum fetuin-A in metabolic and inflammatory pathways in patients with myocardial infarction. European Journal of Clinical Investigation, 2011, 41, 703-709.	3.4	38
7	Potential correlates of burnout among general practitioners and residents in Hungary: the significant role of gender, age, dependant care and experience. BMC Family Practice, 2018, 19, 193.	2.9	37
8	Preeclampsia is associated with decreased serum α2-HS glycoprotein (fetuin-A) concentration. Hypertension Research, 2009, 32, 665-669.	2.7	36
9	Human serum fetuin Alα2HS-glycoprotein level is associated with long-term survival in patients with alcoholic liver cirrhosis, comparison with the Child-Pugh and MELD scores. BMC Gastroenterology, 2007, 7, 15.	2.0	35
10	Personality and cardiovascular risk: Association between hypertension and affective temperaments—a cross-sectional observational study in primary care settings. European Journal of General Practice, 2014, 20, 247-252.	2.0	33
11	Alpha2-HS glycoprotein: a protein in search of a function. Diabetes/Metabolism Research and Reviews, 2002, 18, 311-314.	4.0	28
12	Primary care obesity management in Hungary: evaluation of the knowledge, practice and attitudes of family physicians. BMC Family Practice, 2013, 14, 156.	2.9	27
13	Paradoxical alteration of acute-phase protein levels in patients with chronic hepatitis C treated with IFN-Â2b. International Immunology, 2004, 16, 51-54.	4.0	23
14	Socio-demographic characteristics, health behaviour, co-morbidity and accidents in snorers: a population survey. Sleep and Breathing, 2011, 15, 809-818.	1.7	23
15	Affective Temperaments Contribute to Cardiac Complications in Hypertension Independently of Depression. Psychotherapy and Psychosomatics, 2014, 83, 187-189.	8.8	18
16	Psychometric Properties and Construct Validity of the Brief Wisconsin Inventory of Smoking Dependence Motives in an Internet-Based Sample of Treatment-Seeking Hungarian Smokers. Nicotine and Tobacco Research, 2011, 13, 273-281.	2.6	16
17	Decreased serum alfa2-HS-glycoprotein concentration in patients with primary biliary cirrhosis. Journal of Hepatology, 1997, 26, 1426-1427.	3.7	15
18	The association of affective temperaments with smoking initiation and maintenance in adult primary care patients. Journal of Affective Disorders, 2015, 172, 397-402.	4.1	14

LÃiszlÃ³ Kalabay

#	Article	IF	CITATIONS
19	Identification of hypertensive patients with dominant affective temperaments might improve the psychopathological and cardiovascular risk stratification: a pilot, case–control study. Annals of General Psychiatry, 2015, 14, 33.	2.7	11
20	Dose sparing and the lack of a dose–response relationship with an influenza vaccine in adult and elderly patients – a randomized, doubleâ€blind clinical trial. British Journal of Clinical Pharmacology, 2017, 83, 1912-1920.	2.4	11
21	Electrophoretic and isoelectric focusing analysis of human recombinant alpha2-HS glycoprotein produced in insect cells: Analysis of the post-translational events. Electrophoresis, 1996, 17, 529-532.	2.4	10
22	Family history of suicide: A clinical marker for major depression in primary care practice?. Journal of Affective Disorders, 2009, 117, 202-204.	4.1	10
23	Serum Ghrelin Level and TNF-α/Ghrelin Ratio in Patients with Previous Myocardial Infarction. Archives of Medical Research, 2012, 43, 548-554.	3.3	10
24	IL-6 AND LIPOPROTEIN (a) [LP(a)] CONCENTRATIONS ARE RELATED ONLY IN PATIENTS WITH HIGH APO(a) ISOFORMS IN MONOCLONAL GAMMOPATHY. Cytokine, 2002, 18, 340-343.	3.2	8
25	Fetuin-A serum levels in patients with aortic aneurysms of Marfan syndrome and atherosclerosis. European Journal of Clinical Investigation, 2011, 41, 176-182.	3.4	8
26	Association of Human Fetuin-A rs4917 Polymorphism with Obesity in 2 Cohorts. Journal of Investigative Medicine, 2015, 63, 548-553.	1.6	8
27	Case report of exercise and statin-fibrate combination therapy-caused myopathy in a patient with metabolic syndrome: contradictions between the two main therapeutic pathways. BMC Research Notes, 2013, 6, 52.	1.4	7
28	Perceived cardiovascular risk and attitude towards lifestyle change. Family Practice, 2018, 35, 712-717.	1.9	6
29	Diagnosis and successful management of calciphylaxis in a pancreas–kidney transplant patient. Nephrology Dialysis Transplantation, 2005, 20, 1520-1521.	0.7	5
30	Human Fetuin-A Rs4918 Polymorphism and its Association with Obesity in Healthy Persons and in Patients with Myocardial Infarction in Two Hungarian Cohorts. Medical Science Monitor, 2016, 22, 2742-2750.	1.1	5
31	Fibronectin on the surface of human lymphocytes. Immunology Letters, 1985, 9, 301-305.	2.5	4
32	The Diagnosis and Therapy of <i>Helicobacter pylori</i> Infection in Hungary: Comparison of Strategies Applied by Family Physicians and Internists. Digestive Diseases, 2019, 37, 434-443.	1.9	4
33	To prevent being stressed-out: Allostatic overload and resilience of general practitioners in the era of COVID-19. A cross-sectional observational study. European Journal of General Practice, 2021, 27, 277-285.	2.0	4
34	Serum fetuin-A, tumor necrosis factor alpha and C-reactive protein concentrations in patients with hereditary angioedema with C1-inhibitor deficiency. Orphanet Journal of Rare Diseases, 2019, 14, 67.	2.7	3
35	Licensing the first reduced, 6â€ ⁻ µg dose whole virion, aluminum adjuvanted seasonal influenza vaccine – A randomized-controlled multicenter trial. Vaccine, 2019, 37, 258-264.	3.8	3
36	Care management of patients with high cardiovascular risk in Hungary an international and Hungarian longitudinal comparison of target level achievement. BMC Family Practice, 2020, 21, 83.	2.9	3

LÃiszlÃ³ Kalabay

#	Article	IF	CITATIONS
37	Genetic Influence on Capillary Oxygen Saturation: A Twin Study. Lung, 2014, 192, 429-434.	3.3	2
38	Association of PPAR Alpha Intron 7 G/C, PPAR Gamma 2 Pro12Ala, and C161T Polymorphisms with Serum Fetuin-A Concentrations. PPAR Research, 2017, 2017, 1-8.	2.4	2
39	Diagnosis and successful management of calciphylaxis in a pancreas–kidney transplant patient. Nephrology Dialysis Transplantation, 2005, 20, 2295-2295.	0.7	1
40	Semmelweis Caring University Model Program Based on the Development of a Center of Preventive Services: Health for All Employees at a University Occupational Setting. Frontiers in Public Health, 2021, 9, 727668.	2.7	1
41	An appraisal: how notifiable infectious diseases are reported by Hungarian family physicians. BMC Infectious Diseases, 2018, 18, 45.	2.9	Ο
42	Evaluation of primary care services in Hungary: a comprehensive description of provision, professional competences, cooperation, financing, and infrastructure, based on the findings of the Hungarian-arm of the QUALICOPC study. Primary Health Care Research and Development, 2021, 22, e36.	1.2	0
43	The Missing Link: A Case of Severe Adverse Reaction to Histamine in Food and Beverages. American Journal of Case Reports, 2022, 23, e934212.	0.8	Ο
44	Serum fetuin-A level is independent of Helicobacter pylori postinfection status in systemic lupus erythematosus. Acta Microbiologica Et Immunologica Hungarica, 2022, , .	0.8	0