

Dorte Glintborg

List of Publications by Citations

Source: <https://exaly.com/author-pdf/6385749/dorte-glintborg-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

126
papers

3,284
citations

30
h-index

53
g-index

140
ext. papers

3,884
ext. citations

4
avg, IF

5.35
L-index

#	Paper	IF	Citations
126	Mitochondrial respiration is decreased in skeletal muscle of patients with type 2 diabetes. <i>Diabetes</i> , 2007 , 56, 1592-9	0.9	397
125	Reduced expression of nuclear-encoded genes involved in mitochondrial oxidative metabolism in skeletal muscle of insulin-resistant women with polycystic ovary syndrome. <i>Diabetes</i> , 2007 , 56, 2349-55	0.9	135
124	Association of pioglitazone treatment with decreased bone mineral density in obese premenopausal patients with polycystic ovary syndrome: a randomized, placebo-controlled trial. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008 , 93, 1696-701	5.6	114
123	An update on the pathogenesis, inflammation, and metabolism in hirsutism and polycystic ovary syndrome. <i>Gynecological Endocrinology</i> , 2010 , 26, 281-96	2.4	103
122	Impaired insulin-stimulated phosphorylation of Akt and AS160 in skeletal muscle of women with polycystic ovary syndrome is reversed by pioglitazone treatment. <i>Diabetes</i> , 2008 , 57, 357-66	0.9	103
121	Prevalence of endocrine diseases and abnormal glucose tolerance tests in 340 Caucasian premenopausal women with hirsutism as the referral diagnosis. <i>Fertility and Sterility</i> , 2004 , 82, 1570-9	4.8	99
120	Evaluation of metabolic risk markers in polycystic ovary syndrome (PCOS). Adiponectin, ghrelin, leptin and body composition in hirsute PCOS patients and controls. <i>European Journal of Endocrinology</i> , 2006 , 155, 337-45	6.5	93
119	Development and Risk Factors of Type 2 Diabetes in a Nationwide Population of Women With Polycystic Ovary Syndrome. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017 , 102, 3848-3857	5.6	90
118	Cardiovascular disease in a nationwide population of Danish women with polycystic ovary syndrome. <i>Cardiovascular Diabetology</i> , 2018 , 17, 37	8.7	89
117	Morbidity and medicine prescriptions in a nationwide Danish population of patients diagnosed with polycystic ovary syndrome. <i>European Journal of Endocrinology</i> , 2015 , 172, 627-38	6.5	83
116	Soluble CD36 and risk markers of insulin resistance and atherosclerosis are elevated in polycystic ovary syndrome and significantly reduced during pioglitazone treatment. <i>Diabetes Care</i> , 2008 , 31, 328-34	14.6	78
115	MON-LB131 Perfluoroalkyl Substance Exposure Was Negatively Associated With Cortisone Levels in Pregnancy. <i>Journal of the Endocrine Society</i> , 2020 , 4,	0.4	78
114	A 12-month randomized crossover study on the effects of lanreotide Autogel and octreotide long-acting repeatable on GH and IGF-I in patients with acromegaly. <i>Clinical Endocrinology</i> , 2008 , 68, 473-80	3.4	63
113	Normo- and hyperandrogenic women with polycystic ovary syndrome exhibit an adverse metabolic profile through life. <i>Fertility and Sterility</i> , 2017 , 107, 788-795.e2	4.8	60
112	Plasma monocyte chemoattractant protein-1 (MCP-1) and macrophage inflammatory protein-1alpha are increased in patients with polycystic ovary syndrome (PCOS) and associated with adiposity, but unaffected by pioglitazone treatment. <i>Clinical Endocrinology</i> , 2009 , 71, 652-8	3.4	58
111	Pioglitazone enhances mitochondrial biogenesis and ribosomal protein biosynthesis in skeletal muscle in polycystic ovary syndrome. <i>PLoS ONE</i> , 2008 , 3, e2466	3.7	58
110	Body composition is improved during 12 months treatment with metformin alone or combined with oral contraceptives compared with treatment with oral contraceptives in polycystic ovary syndrome. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014 , 99, 2584-91	5.6	55

109	Bioactive insulin-like growth factor (IGF) I and IGF-binding protein-1 in anorexia nervosa. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2007 , 92, 2323-9	5.6	50
108	The use of medication against attention deficit hyperactivity disorder in Denmark: a drug use study from a national perspective. <i>European Journal of Clinical Pharmacology</i> , 2012 , 68, 1443-50	2.8	49
107	Effect of pioglitazone on glucose metabolism and luteinizing hormone secretion in women with polycystic ovary syndrome. <i>Fertility and Sterility</i> , 2006 , 86, 385-97	4.8	49
106	Ethnic differences in Rotterdam criteria and metabolic risk factors in a multiethnic group of women with PCOS studied in Denmark. <i>Clinical Endocrinology</i> , 2010 , 73, 732-8	3.4	47
105	Gestational Diabetes Mellitus: Does One Size Fit All? A Challenge to Uniform Worldwide Diagnostic Thresholds. <i>Diabetes Care</i> , 2018 , 41, 1339-1342	14.6	44
104	The use of medication against attention deficit/hyperactivity disorder in Denmark: a drug use study from a patient perspective. <i>European Journal of Clinical Pharmacology</i> , 2013 , 69, 589-98	2.8	40
103	The effect of menopause on carotid artery remodeling, insulin sensitivity, and plasma adiponectin in healthy women. <i>American Journal of Hypertension</i> , 2009 , 22, 364-70	2.3	39
102	Effect of testosterone on insulin sensitivity, oxidative metabolism and body composition in aging men with type 2 diabetes on metformin monotherapy. <i>Diabetes, Obesity and Metabolism</i> , 2016 , 18, 980-9	6.7	37
101	Association of polycystic ovary syndrome susceptibility single nucleotide polymorphism rs2479106 and PCOS in Caucasian patients with PCOS or hirsutism as referral diagnosis. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2012 , 163, 39-42	2.4	36
100	Significantly higher adrenocorticotropin-stimulated cortisol and 17-hydroxyprogesterone levels in 337 consecutive, premenopausal, caucasian, hirsute patients compared with healthy controls. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2005 , 90, 1347-53	5.6	34
99	Higher bone mineral density in Caucasian, hirsute patients of reproductive age. Positive correlation of testosterone levels with bone mineral density in hirsutism. <i>Clinical Endocrinology</i> , 2005 , 62, 683-91	3.4	33
98	Pioglitazone treatment increases spontaneous growth hormone (GH) secretion and stimulated GH levels in polycystic ovary syndrome. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2005 , 90, 5605-12	5.6	33
97	Hemoglobin A1c as a tool for the diagnosis of type 2 diabetes in 208 premenopausal women with polycystic ovary syndrome. <i>Fertility and Sterility</i> , 2011 , 96, 1275-80	4.8	31
96	Impaired insulin activation and dephosphorylation of glycogen synthase in skeletal muscle of women with polycystic ovary syndrome is reversed by pioglitazone treatment. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008 , 93, 3618-26	5.6	29
95	Total and high molecular weight (HMW) adiponectin levels and measures of glucose and lipid metabolism following pioglitazone treatment in a randomized placebo-controlled study in polycystic ovary syndrome. <i>Clinical Endocrinology</i> , 2008 , 68, 165-74	3.4	29
94	MANAGEMENT OF ENDOCRINE DISEASE: Morbidity in polycystic ovary syndrome. <i>European Journal of Endocrinology</i> , 2017 , 176, R53-R65	6.5	28
93	The prevalence of Type 2 diabetes is not increased in normal-weight women with PCOS. <i>Human Reproduction</i> , 2017 , 32, 2279-2286	5.7	27
92	Hyperandrogenism and phenotypes of polycystic ovary syndrome are not associated with differences in obstetric outcomes. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2015 , 94, 204-11	3.8	26

91	Prolactin is associated with metabolic risk and cortisol in 1007 women with polycystic ovary syndrome. <i>Human Reproduction</i> , 2014 , 29, 1773-9	5.7	26
90	Free carnitine and acylcarnitines in obese patients with polycystic ovary syndrome and effects of pioglitazone treatment. <i>Fertility and Sterility</i> , 2012 , 98, 1620-6.e1	4.8	26
89	Genetic alterations within the DENND1A gene in patients with polycystic ovary syndrome (PCOS). <i>PLoS ONE</i> , 2013 , 8, e77186	3.7	26
88	Intact primary mitochondrial function in myotubes established from women with PCOS. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011 , 96, E1298-302	5.6	25
87	Perfluoroalkyl substances and glycemic status in pregnant Danish women: The Odense Child Cohort. <i>Environment International</i> , 2018 , 116, 101-107	12.9	24
86	Increased thrombin generation in women with polycystic ovary syndrome: A pilot study on the effect of metformin and oral contraceptives. <i>Metabolism: Clinical and Experimental</i> , 2015 , 64, 1272-8	12.7	23
85	Comparison of regional fat mass measurement by whole body DXA scans and anthropometric measures to predict insulin resistance in women with polycystic ovary syndrome and controls. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2016 , 95, 1235-1243	3.8	23
84	Smoking is associated with increased adrenal responsiveness, decreased prolactin levels and a more adverse lipid profile in 650 white patients with polycystic ovary syndrome. <i>Gynecological Endocrinology</i> , 2012 , 28, 170-4	2.4	22
83	Insulin resistance is not conserved in myotubes established from women with PCOS. <i>PLoS ONE</i> , 2010 , 5, e14469	3.7	22
82	Polycystic ovary syndrome and hyperglycaemia in pregnancy. A narrative review and results from a prospective Danish cohort study. <i>Diabetes Research and Clinical Practice</i> , 2018 , 145, 167-177	7.4	21
81	The prevalence of endometrial hyperplasia and endometrial cancer in women with polycystic ovary syndrome or hyperandrogenism. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2012 , 91, 1173-6	3.8	21
80	A randomized placebo-controlled study on the effects of pioglitazone on cortisol metabolism in polycystic ovary syndrome. <i>Fertility and Sterility</i> , 2009 , 91, 842-50	4.8	21
79	Fracture Risk Is Decreased in Women With Polycystic Ovary Syndrome: A Register-Based and Population-Based Cohort Study. <i>Journal of Bone and Mineral Research</i> , 2016 , 31, 709-17	6.3	20
78	Birth weight and polycystic ovary syndrome in adult life: a register-based study on 523,757 Danish women born 1973-1991. <i>Fertility and Sterility</i> , 2013 , 99, 777-82	4.8	20
77	Differential DNA methylation patterns of polycystic ovarian syndrome in whole blood of Chinese women. <i>Oncotarget</i> , 2017 , 8, 20656-20666	3.3	19
76	Thiazolidinedione treatment in PCOS--an update. <i>Gynecological Endocrinology</i> , 2010 , 26, 791-803	2.4	19
75	Age associated differences in prevalence of individual rotterdam criteria and metabolic risk factors during reproductive age in 446 caucasian women with polycystic ovary syndrome. <i>Hormone and Metabolic Research</i> , 2012 , 44, 694-8	3.1	18
74	Testosterone Levels in Third Trimester in Polycystic Ovary Syndrome: Odense Child Cohort. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018 , 103, 3819-3827	5.6	18

73	The Pharmacogenetics of Metformin in Women with Polycystic Ovary Syndrome: A Randomized Trial. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2018 , 122, 239-244	3.1	17
72	Effects of exercise and group counselling on body composition and VO2max in overweight women with polycystic ovary syndrome. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2013 , 92, 272-7	3.8	17
71	Adiponectin, interleukin-6, monocyte chemoattractant protein-1, and regional fat mass during 12-month randomized treatment with metformin and/or oral contraceptives in polycystic ovary syndrome. <i>Journal of Endocrinological Investigation</i> , 2014 , 37, 757-764	5.2	16
70	The relationship between health-related quality of life, obesity and testosterone levels in older men. <i>Age and Ageing</i> , 2014 , 43, 280-4	3	15
69	Strength training and testosterone treatment have opposing effects on migration inhibitor factor levels in ageing men. <i>Mediators of Inflammation</i> , 2013 , 2013, 539156	4.3	15
68	Myocardial response to a triathlon in male athletes evaluated by Doppler tissue imaging and biochemical parameters. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2008 , 18, 698-705	4.6	15
67	Effect of oral contraceptives and/or metformin on GLP-1 secretion and reactive hypoglycaemia in polycystic ovary syndrome. <i>Endocrine Connections</i> , 2017 , 6, 267-277	3.5	13
66	Anogenital distance in children born of mothers with polycystic ovary syndrome: the Odense Child Cohort. <i>Human Reproduction</i> , 2019 , 34, 2061-2070	5.7	13
65	Prescription of antidepressants is increased in Danish patients with polycystic ovary syndrome and is associated with hyperandrogenism. A population-based cohort study. <i>Clinical Endocrinology</i> , 2014 , 80, 884-9	3.4	13
64	Testosterone treatment increases androgen receptor and aromatase gene expression in myotubes from patients with PCOS and controls, but does not induce insulin resistance. <i>Biochemical and Biophysical Research Communications</i> , 2014 , 451, 622-6	3.4	13
63	Medical comorbidity in polycystic ovary syndrome with special focus on cardiometabolic, autoimmune, hepatic and cancer diseases: an updated review. <i>Current Opinion in Obstetrics and Gynecology</i> , 2017 , 29, 390-396	2.4	13
62	Insulin resistance in pregnant women with and without polycystic ovary syndrome, and measures of body composition in offspring at birth and three years of age. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2017 , 96, 1307-1314	3.8	13
61	A rare cause of Cushing's syndrome: an ACTH-secreting pheochromocytoma. <i>BMJ Case Reports</i> , 2014 , 2014,	0.9	13
60	Awareness of polycystic ovary syndrome among obstetrician-gynecologists and endocrinologists in Northern Europe. <i>PLoS ONE</i> , 2019 , 14, e0226074	3.7	13
59	Metabolic Syndrome in Hyperprolactinemia. <i>Frontiers of Hormone Research</i> , 2018 , 49, 29-47	3.5	12
58	Diagnosis and follow-up of type 2 diabetes in women with PCOS: a role for OGTT?. <i>European Journal of Endocrinology</i> , 2018 , 179, D1-D14	6.5	12
57	Supportive relationships--psychological effects of group counselling in women with polycystic ovary syndrome (PCOS). <i>Communication and Medicine</i> , 2012 , 9, 125-31	0	12
56	Endocrine and metabolic characteristics in polycystic ovary syndrome. <i>Danish Medical Journal</i> , 2016 , 63,	3.8	12

55	Osteoprotegerin levels decrease during testosterone therapy in aging men and are associated with changed distribution of regional fat. <i>Hormone and Metabolic Research</i> , 2013 , 45, 308-13	3.1	10
54	Bone mineral density and vitamin D in PCOS and hirsutism. <i>Expert Review of Endocrinology and Metabolism</i> , 2013 , 8, 449-459	4.1	10
53	Prenatal Exposures to Perfluoroalkyl Acids and Associations with Markers of Adiposity and Plasma Lipids in Infancy: An Odense Child Cohort Study. <i>Environmental Health Perspectives</i> , 2020 , 128, 77001	8.4	9
52	Prevalence and possible mechanisms of reactive hypoglycemia in polycystic ovary syndrome. <i>Human Reproduction</i> , 2016 , 31, 1105-12	5.7	9
51	Testosterone replacement therapy of opioid-induced male hypogonadism improved body composition but not pain perception: a double-blind, randomized, and placebo-controlled trial. <i>European Journal of Endocrinology</i> , 2020 , 182, 539-548	6.5	9
50	Total testosterone levels are often more than three times elevated in patients with androgen-secreting tumours. <i>BMJ Case Reports</i> , 2015 , 2015,	0.9	8
49	The effect of dietary carbohydrates in women with polycystic ovary syndrome: a systematic review. <i>Minerva Endocrinologica</i> , 2016 , 41, 57-69	1.9	8
48	Prenatal exposure to perfluorodecanoic acid is associated with lower circulating concentration of adrenal steroid metabolites during mini puberty in human female infants. The Odense Child Cohort. <i>Environmental Research</i> , 2020 , 182, 109101	7.9	7
47	Effect of 12-month treatment with metformin and/or oral contraceptives on health-related quality of life in polycystic ovary syndrome. <i>Gynecological Endocrinology</i> , 2018 , 34, 859-863	2.4	7
46	General practitioners and hospital physicians' preference for morphine or oxycodone as first-time choice for a strong opioid: a National Register-based study. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2013 , 112, 110-5	3.1	7
45	Long-Term Effect of Interactive Online Dietician Weight Loss Advice in General Practice (LIVA) Protocol for a Randomized Controlled Trial. <i>International Journal of Family Medicine</i> , 2014 , 2014, 245347		7
44	Antipsychotic treatment for children and adolescents with schizophrenia spectrum disorders: protocol for a network meta-analysis of randomised trials. <i>BMJ Open</i> , 2014 , 4, e005708	3	7
43	Increased risk of thyroid disease in Danish women with polycystic ovary syndrome: a cohort study. <i>Endocrine Connections</i> , 2019 , 8, 1405-1415	3.5	7
42	Maternal phthalate exposure associated with decreased testosterone/LH ratio in male offspring during mini-puberty. Odense Child Cohort. <i>Environment International</i> , 2020 , 144, 106025	12.9	7
41	Metformin decreases miR-122, miR-223 and miR-29a in women with polycystic ovary syndrome. <i>Endocrine Connections</i> , 2020 , 9, 1075-1084	3.5	6
40	Differential effects of strength training and testosterone treatment on soluble CD36 in aging men: Possible relation to changes in body composition. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2015 , 75, 659-66	2	6
39	Medical treatment and comorbidity in polycystic ovary syndrome: An updated review. <i>Current Opinion in Endocrine and Metabolic Research</i> , 2020 , 12, 33-40	1.7	5
38	Fibrin lysability is associated with central obesity and inflammation in women with polycystic ovary syndrome. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2020 , 99, 1078-1084	3.8	5

37	Adrenal activity and metabolic risk during randomized escitalopram or placebo treatment in PCOS. <i>Endocrine Connections</i> , 2018 , 7, 479-489	3.5	5
36	Mitochondria, glycogen, and lipid droplets in skeletal muscle during testosterone treatment and strength training: a randomized, double-blinded, placebo-controlled trial. <i>Andrology</i> , 2018 , 6, 547-555	4.2	5
35	Blood pressure in 3-year-old girls associates inversely with umbilical cord serum 25-hydroxyvitamin D: an Odense Child Cohort study. <i>Endocrine Connections</i> , 2018 , 7, 1236-1244	3.5	5
34	Should we individualize lipid profiling in women with polycystic ovary syndrome?. <i>Human Reproduction</i> , 2016 , 31, 2791-2795	5.7	5
33	Motivational interviewing in obese women with polycystic ovary syndrome - a pilot study. <i>Gynecological Endocrinology</i> , 2019 , 35, 76-80	2.4	5
32	Socioeconomic status in Danish women with polycystic ovary syndrome: A register-based cohort study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2019 , 98, 440-450	3.8	5
31	Effects of testosterone therapy on constructs related to aggression in transgender men: A systematic review. <i>Hormones and Behavior</i> , 2021 , 128, 104912	3.7	5
30	MANAGEMENT OF ENDOCRINE DISEASE: Optimal feminizing hormone treatment in transgender people. <i>European Journal of Endocrinology</i> , 2021 , 185, R49-R63	6.5	5
29	Blood pressure and hypertension during pregnancy in women with polycystic ovary syndrome: Odense Child Cohort. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2020 , 99, 1354-1363	3.8	4
28	Third trimester cortisol status is associated with offspring sex and polycystic ovary syndrome status: Odense Child Cohort. <i>Fertility and Sterility</i> , 2019 , 112, 764-772	4.8	4
27	Plasma osteoprotegerin is associated with testosterone levels but unaffected by pioglitazone treatment in patients with polycystic ovary syndrome. <i>Journal of Endocrinological Investigation</i> , 2013 , 36, 460-5	5.2	4
26	Perfluoroalkyl Substance Exposure Early In Pregnancy Was Negatively Associated With Late Pregnancy Cortisone Levels. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020 , 105,	5.6	3
25	Response to Letter to the Editor: "Development and Risk Factors of Type 2 Diabetes in a Nationwide Population of Women With Polycystic Ovary Syndrome". <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018 , 103, 362-363	5.6	3
24	Danish physicians' preferences for prescribing escitalopram over citalopram and sertraline to treatment-naïve patients: a national, register-based study. <i>European Journal of Clinical Pharmacology</i> , 2013 , 69, 1167-71	2.8	3
23	Clinical manifest x-linked recessive adrenoleukodystrophy in a female. <i>Case Reports in Neurological Medicine</i> , 2013 , 2013, 491790	0.7	3
22	Risk of impaired glucose tolerance in normal weight hirsute women during four years observation. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2010 , 89, 1091-5	3.8	3
21	Maternal prolactin is associated with glucose status and PCOS in pregnancy: Odense Child Cohort. <i>European Journal of Endocrinology</i> , 2020 , 183, 307-316	6.5	3
20	Licorice-induced apparent mineralocorticoid excess compounded by excessive use of terbutaline and high water intake. <i>BMJ Case Reports</i> , 2018 , 2018,	0.9	3

19	Masculinising testosterone treatment and effects on preclinical cardiovascular disease, muscle strength and power, aggression, physical fitness and respiratory function in transgender men: protocol for a 10-year, prospective, observational cohort study in Denmark at the Body Identity Clinic (BIC). <i>BMJ Open</i> , 2020 , 10, e045714	3	2
18	Adapting fasting plasma glucose threshold for GDM diagnosis according to the population distribution - An approach to the Danish paradox. <i>Diabetes Research and Clinical Practice</i> , 2021 , 175, 108832	7.4	2
17	Higher blood pressure in normal weight women with PCOS compared to controls. <i>Endocrine Connections</i> , 2021 , 10, 154-163	3.5	2
16	Recurrent autoimmune hypophysitis treated with rituximab: a case report.. <i>Journal of Medical Case Reports</i> , 2021 , 15, 591	1.2	2
15	Free thyroxine in early pregnancy is an independent negative predictor of 3rd trimester HbA1c. Odense child cohort. <i>Clinical Endocrinology</i> , 2021 , 95, 508-519	3.4	1
14	Pregnancy or cord 25-hydroxyvitamin D is not associated with measures of body fat or adiposity in children from three months to three years of age. An Odense Child Cohort study. <i>Clinical Nutrition</i> , 2020 , 39, 1832-1839	5.9	1
13	Accelerated osteoarthritis in women with polycystic ovary syndrome: a prospective nationwide registry-based cohort study. <i>Arthritis Research and Therapy</i> , 2021 , 23, 225	5.7	1
12	Birthweight, Childhood Body Mass Index, Height and Growth, and Risk of Polycystic Ovary Syndrome. <i>Obesity Facts</i> , 2021 , 14, 283-290	5.1	1
11	Maternal Testosterone Concentrations in Third Trimester and Offspring Handgrip Strength at 5 Years: Odense Child Cohort.. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2022 ,	5.6	1
10	Gender incongruence in Denmark, a quantitative assessment. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2021 , 100, 1800-1805	3.8	0
9	Socioeconomic status in Danish transgender persons: a nationwide register-based cohort study. <i>Endocrine Connections</i> , 2021 , 10, 1155-1166	3.5	0
8	Maternal polycystic ovary syndrome and attention deficit hyperactivity disorder in offspring at 3 years of age: Odense Child Cohort. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2021 , 100, 2053-2065	3.8	0
7	Systematic genetic screening in a prospective group of Danish patients with pheochromocytoma. <i>Research and Reports in Urology</i> , 2017 , 9, 113-119	1.3	
6	Prenatal Androgen Exposure and Traits of Autism Spectrum Disorder in the Offspring: Odense Child Cohort.. <i>Journal of Autism and Developmental Disorders</i> , 2022 , 1	4.6	
5	Awareness of polycystic ovary syndrome among obstetrician-gynecologists and endocrinologists in Northern Europe 2019 , 14, e0226074		
4	Awareness of polycystic ovary syndrome among obstetrician-gynecologists and endocrinologists in Northern Europe 2019 , 14, e0226074		
3	Awareness of polycystic ovary syndrome among obstetrician-gynecologists and endocrinologists in Northern Europe 2019 , 14, e0226074		
2	Awareness of polycystic ovary syndrome among obstetrician-gynecologists and endocrinologists in Northern Europe 2019 , 14, e0226074		

- 1 Higher free thyroxine associated with PFAS exposure in first trimester. The Odense Child Cohort.. *Environmental Research*, **2022**, 212, 113492 7.9