Kai-Hsiang Chuang

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6394056/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	The basal forebrain volume reduction detected by MRI does not necessarily link with the cholinergic neuronal loss in the Alzheimer's disease mouse model. Neurobiology of Aging, 2022, 117, 24-32.	1.5	2
2	Deep learning network for integrated coil inhomogeneity correction and brain extraction of mixed MRI data. Scientific Reports, 2022, 12, .	1.6	2
3	Optimizing diffusion MRI acquisition efficiency of rodent brain using simultaneous multislice EPI. NMR in Biomedicine, 2021, 34, e4398.	1.6	3
4	Quantitative Analysis of Renal Perfusion by Arterial Spin Labeling. Methods in Molecular Biology, 2021, 2216, 655-666.	0.4	0
5	Renal Blood Flow Using Arterial Spin Labeling (ASL) MRI: Experimental Protocol and Principles. Methods in Molecular Biology, 2021, 2216, 443-453.	0.4	1
6	Task-related brain functional network reconfigurations relate to motor recovery in chronic subcortical stroke. Scientific Reports, 2021, 11, 8442.	1.6	19
7	Brain Functional Changes in Stroke Following Rehabilitation Using Brain-Computer Interface-Assisted Motor Imagery With and Without tDCS: A Pilot Study. Frontiers in Human Neuroscience, 2021, 15, 692304.	1.0	16
8	Dysfunction in nonsense-mediated decay, protein homeostasis, mitochondrial function, and brain connectivity in ALS-FUS mice with cognitive deficits. Acta Neuropathologica Communications, 2021, 9, 9.	2.4	17
9	Neurogenic-dependent changes in hippocampal circuitry underlie the procognitive effect of exercise in aging mice. IScience, 2021, 24, 103450.	1.9	9
10	Protocol for a pilot randomised controlled trial of mindfulness-based cognitive therapy in youth with inflammatory bowel disease and depression. BMJ Open, 2019, 9, e025568.	0.8	9
11	Ultrafast fMRI of the rodent brain using simultaneous multi-slice EPI. NeuroImage, 2019, 195, 48-58.	2.1	27
12	Evaluation of nuisance removal for functional MRI of rodent brain. NeuroImage, 2019, 188, 694-709.	2.1	30
13	Magnetic resonance imaging quantification and biodistribution of magnetic nanoparticles using <i>T</i> ₁ -enhanced contrast. Journal of Materials Chemistry B, 2018, 6, 1470-1478.	2.9	6
14	Differential Amplitude of Low-Frequency Fluctuations in brain networks after BCI Training with and without tDCS in Stroke. , 2018, 2018, 1050-1053.		5
15	Connectomic imaging reveals Huntingtonâ€related pathological and pharmaceutical effects in a mouse model. NMR in Biomedicine, 2018, 31, e4007.	1.6	18
16	Cerebro-Cerebellar Pathways for Verbal Working Memory. Frontiers in Human Neuroscience, 2018, 12, 530.	1.0	4
17	GABAergic effect on resting-state functional connectivity: Dynamics under pharmacological antagonism. Neurolmage, 2017, 149, 53-62.	2.1	19
18	Intravoxel incoherent motion and diffusion tensor imaging of early renal fibrosis induced in a murine model of streptozotocin induced diabetes. Magnetic Resonance Imaging, 2017, 38, 71-76.	1.0	28

#	Article	IF	CITATIONS
19	Functional networks and network perturbations in rodents. NeuroImage, 2017, 163, 419-436.	2.1	55
20	Brain plasticity following MI-BCI training combined with tDCS in a randomized trial in chronic subcortical stroke subjects: a preliminary study. Scientific Reports, 2017, 7, 9222.	1.6	51
21	Transgenic Mice Overexpressing the Divalent Metal Transporter 1 Exhibit Iron Accumulation and Enhanced Parkin Expression in the Brain. NeuroMolecular Medicine, 2017, 19, 375-386.	1.8	21
22	MEMRI detects neuronal activity and connectivity in hypothalamic neural circuit responding to leptin. NeuroImage, 2017, 147, 904-915.	2.1	3
23	Caloric restriction preserves memory and reduces anxiety of aging mice with early enhancement of neurovascular functions. Aging, 2016, 8, 2814-2826.	1.4	62
24	Synthesis of Ferromagnetic Fe _{0.6} Mn _{0.4} O Nanoflowers as a New Class of Magnetic Theranostic Platform for In Vivo T ₁ â€T ₂ Dualâ€Mode Magnetic Resonance Imaging and Magnetic Hyperthermia Therapy. Advanced Healthcare Materials, 2016, 5, 2092-2104	3.9	75
25	Octreotide Functionalized Nano-Contrast Agent for Targeted Magnetic Resonance Imaging. Biomacromolecules, 2016, 17, 3902-3910.	2.6	19
26	Modality specificity in the cerebro-cerebellar neurocircuitry during working memory. Behavioural Brain Research, 2016, 305, 164-173.	1.2	17
27	Laquinimod rescues striatal, cortical and white matter pathology and results in modest behavioural improvements in the YAC128 model of Huntington disease. Scientific Reports, 2016, 6, 31652.	1.6	59
28	Multispectral optoacoustic and MRI coregistration for molecular imaging of orthotopic model of human glioblastoma. Journal of Biophotonics, 2016, 9, 701-708.	1.1	35
29	Structural and molecular myelination deficits occur prior to neuronal loss in the YAC128 and BACHD models of Huntington disease. Human Molecular Genetics, 2016, 25, ddw122.	1.4	62
30	Resting state functional connectivity data supports detection of cognition in the rodent brain. Data in Brief, 2016, 7, 1156-1164.	0.5	7
31	A method for the automatic segmentation of brown adipose tissue. Magnetic Resonance Materials in Physics, Biology, and Medicine, 2016, 29, 287-299.	1.1	17
32	Temporal dynamics of visual working memory. NeuroImage, 2016, 124, 1021-1030.	2.1	23
33	Functional connectivity MRI tracks memory networks after maze learning in rodents. NeuroImage, 2016, 127, 196-202.	2.1	30
34	Spatial memory training induces morphological changes detected by manganese-enhanced MRI in the hippocampal CA3 mossy fiber terminal zone. NeuroImage, 2016, 128, 227-237.	2.1	7
35	Optimization of arterial spin labeling MRI for quantitative tumor perfusion in a mouse xenograft model. NMR in Biomedicine, 2015, 28, 988-997.	1.6	5
36	Molecular Photoacoustic Imaging of Orthotopic Glioblastoma. , 2015, , .		1

3

#	Article	IF	CITATIONS
37	Dependence of BOLD signal fluctuation on arterial blood CO2 and O2: Implication for resting-state functional connectivity. NeuroImage, 2015, 117, 29-39.	2.1	33
38	Combining EPI and motion correction for fMRI human brain images with big motion. , 2015, 2015, 5449-52.		1
39	Pharmacokinetics of Gd(DO3A‣ys) and MR imaging studies in an orthotopic U87MG glioma tumor model. Contrast Media and Molecular Imaging, 2015, 10, 237-244.	0.4	7
40	Evaluation of EPI distortion correction methods for quantitative MRI of the brain at high magnetic field. Magnetic Resonance Imaging, 2015, 33, 1098-1105.	1.0	42
41	Non-canonical NF-κB signalling and ETS1/2 cooperatively drive C250T mutant TERT promoterÂactivation. Nature Cell Biology, 2015, 17, 1327-1338.	4.6	178
42	Intravoxel incoherent imaging of renal fibrosis induced in a murine model of unilateral ureteral obstruction. Magnetic Resonance Imaging, 2015, 33, 1324-1328.	1.0	25
43	Evaluation of thermo-triggered drug release in intramuscular-transplanted tumors using thermosensitive polymer-modified liposomes and MRI. Nanomedicine: Nanotechnology, Biology, and Medicine, 2015, 11, 229-238.	1.7	41
44	Dynamic Contrast-Enhanced MRI of Mouse Liver: A Feasibility Study Using a Dual-Input Two-Compartment Tracer Kinetic Model. Journal of Biomedical Science and Engineering, 2015, 08, 90-97.	0.2	0
45	Early detection of antiangiogenic treatment responses in a mouse xenograft tumor model using quantitative perfusion <scp>MRI</scp> . Cancer Medicine, 2014, 3, 47-60.	1.3	17
46	Neural correlate of resting-state functional connectivity under α2 adrenergic receptor agonist, medetomidine. NeuroImage, 2014, 84, 27-34.	2.1	42
47	A facile synthetic approach to a biodegradable polydisulfide MRI contrast agent. Journal of Materials Chemistry B, 2014, 2, 5295-5301.	2.9	12
48	Pharmacological insight into neurotransmission origins of resting-state functional connectivity: α2-adrenergic agonist vs antagonist. Neurolmage, 2014, 103, 364-373.	2.1	16
49	Detection of functional connectivity in the resting mouse brain. NeuroImage, 2014, 86, 417-424.	2.1	96
50	An intravascular MRI contrast agent based on Gd(DO3A-Lys) for tumor angiography. Biomaterials, 2014, 35, 327-336.	5.7	19
51	Functional Networks in Parallel with Cortical Development Associate with Executive Functions in Children. Cerebral Cortex, 2014, 24, 1937-1947.	1.6	37
52	Quantitative mouse renal perfusion using arterial spin labeling. NMR in Biomedicine, 2013, 26, 1225-1232.	1.6	30
53	A Variational Surface Deformation and Subdivision-Based Modeling Framework for Noisy and Small n-Furcated Tube-Like Structures. IEEE Transactions on Biomedical Engineering, 2013, 60, 1589-1598.	2.5	3
54	Specific Inhibition of p25/Cdk5 Activity by the Cdk5 Inhibitory Peptide Reduces Neurodegeneration <i>In Vivo</i> . Journal of Neuroscience, 2013, 33, 334-343.	1.7	87

#	Article	IF	CITATIONS
55	Imaging Brain Deoxyglucose Uptake and Metabolism by Glucocest MRI. Journal of Cerebral Blood Flow and Metabolism, 2013, 33, 1270-1278.	2.4	150
56	Superparamagnetic Nanostructures for Offâ€Resonance Magnetic Resonance Spectroscopic Imaging. Advanced Functional Materials, 2013, 23, 496-505.	7.8	18
57	Magnetic Resonance Imaging (MRI) Contrast Agents for Tumor Diagnosis. Journal of Healthcare Engineering, 2013, 4, 23-46.	1.1	51
58	Atlas-based automatic mouse brain image segmentation revisited: model complexity vs. image registration. Magnetic Resonance Imaging, 2012, 30, 789-798.	1.0	60
59	Stejskal–tanner equation derived in full. Concepts in Magnetic Resonance Part A: Bridging Education and Research, 2012, 40A, 205-214.	0.2	38
60	Synthesis of Manganese Ferrite/Graphene Oxide Nanocomposites for Biomedical Applications. Small, 2012, 8, 3620-3630.	5.2	113
61	Gadolinium chelate with DO3A conjugated 2-(diphenylphosphoryl)-ethyldiphenylphosphonium cation as potential tumor-selective MRI contrast agent. Biomaterials, 2012, 33, 9225-9231.	5.7	18
62	Pharmacological modulation of functional connectivity: α2-adrenergic receptor agonist alters synchrony but not neural activation. NeuroImage, 2012, 60, 436-446.	2.1	45
63	Quantitative measurement of changes in calcium channel activity in vivo utilizing dynamic manganese-enhanced MRI (dMEMRI). NeuroImage, 2012, 60, 392-399.	2.1	6
64	Gd(iii) chelates for MRI contrast agents: from high relaxivity to "smartâ€; from blood pool to blood–brain barrier permeable. MedChemComm, 2012, 3, 552.	3.5	93
65	Optimization of flowâ€sensitive alternating inversion recovery (FAIR) for perfusion functional MRI of rodent brain. NMR in Biomedicine, 2012, 25, 1209-1216.	1.6	19
66	Functional MRI of Neural Plasticity and Drug Effect in the Brain. Advances in Intelligent and Soft Computing, 2012, , 17-25.	0.2	0
67	Quantitative and Noninvasive Assessment of Prenatal X-Ray-Induced CNS Abnormalities Using Magnetic Resonance Imaging. Radiation Research, 2011, 175, 1-9.	0.7	14
68	Mapping plasticity in the forepaw digit barrel subfield of rat brains using functional MRI. NeuroImage, 2011, 54, 1122-1129.	2.1	8
69	Relationship between blood and myocardium manganese levels during manganeseâ€enhanced MRI (MEMRI) with <i>T</i> ₁ mapping in rats. NMR in Biomedicine, 2011, 24, 46-53.	1.6	12
70	Robust Automatic Rodent Brain Extraction Using 3-D Pulse-Coupled Neural Networks (PCNN). IEEE Transactions on Image Processing, 2011, 20, 2554-2564.	6.0	117
71	Novel synthesis of superparamagnetic magnetite nanoclusters for biomedical applications. Journal of Materials Chemistry, 2011, 21, 14717.	6.7	69
72	Multimodal tumor imaging by iron oxides and quantum dots formulated in poly (lactic) Tj ETQq0 0 0 rgBT /Ove	rlock 10 Tf 5.7	50 67 Td (acio

2969-2978.

1

#	Article	IF	CITATIONS
73	Development of Intravascular Contrast Agents for MRI Using Gadolinium Chelates. ChemMedChem, 2011, 6, 781-787.	1.6	9
74	Vitamin E (d-alpha-tocopheryl-co-poly(ethylene glycol) 1000 succinate) micelles-superparamagnetic iron oxide nanoparticles for enhanced thermotherapy and MRI. Biomaterials, 2011, 32, 5663-5672.	5.7	90
75	Imaging metabolic syndrome. EMBO Molecular Medicine, 2010, 2, 196-210.	3.3	10
76	Superparamagnetic iron oxide – Loaded poly (lactic acid)-d-α-tocopherol polyethylene glycol 1000 succinate copolymer nanoparticles as MRI contrast agent. Biomaterials, 2010, 31, 5588-5597.	5.7	103
77	Facile synthesis of water-stable magnetite nanoparticles for clinical MRI and magnetic hyperthermia applications. Nanomedicine, 2010, 5, 1571-1584.	1.7	61
78	In vivo detection of individual glomeruli in the rodent olfactory bulb using manganese enhanced MRI. NeuroImage, 2010, 49, 1350-1356.	2.1	31
79	Ipsilateral cortical fMRI responses after peripheral nerve damage in rats reflect increased interneuron activity. Proceedings of the National Academy of Sciences of the United States of America, 2009, 106, 14114-14119.	3.3	67
80	Temporal changes in the <i>T</i> ₁ and <i>T</i> ₂ relaxation rates (Δ <i>R</i> ₁ and Ĩ" <i>R</i> ₂) in the rat brain are consistent with the tissueâ€clearance rates of elemental manganese. Magnetic Resonance in Medicine, 2009, 61, 1528-1532.	1.9	55
81	Accounting for nonspecific enhancement in neuronal tract tracing using manganese enhanced magnetic resonance imaging. Magnetic Resonance Imaging, 2009, 27, 594-600.	1.0	38
82	Layer specific tracing of corticocortical and thalamocortical connectivity in the rodent using manganese enhanced MRI. NeuroImage, 2009, 44, 923-931.	2.1	45
83	Manganese enhanced MRI reveals functional circuitry in response to odorant stimuliâ~†. NeuroImage, 2009, 44, 363-372.	2.1	61
84	37.3: Smart Transflective Display Integrated with PDLC and OPVâ€Embeddedâ€OLED. Digest of Technical Papers SID International Symposium, 2009, 40, 530-531.	0.1	1
85	Vascular space occupancyâ€dependent functional MRI by tissue suppression. Journal of Magnetic Resonance Imaging, 2008, 28, 219-226.	1.9	15
86	Monitoring dynamic alterations in calcium homeostasis by <i>T</i> ₁ â€weighted and <i>T</i> ₁ â€mapping cardiac manganeseâ€enhanced MRI in a murine myocardial infarction model. NMR in Biomedicine, 2008, 21, 1102-1111.	1.6	37
87	Brain Redox Imaging Using Blood—Brain Barrier-Permeable Nitroxide MRI Contrast Agent. Journal of Cerebral Blood Flow and Metabolism, 2008, 28, 1165-1174.	2.4	86
88	Absence of gender effect on children with attention-deficit/hyperactivity disorder as assessed by optimized voxel-based morphometry. Psychiatry Research - Neuroimaging, 2008, 164, 245-253.	0.9	37
89	Mapping resting-state functional connectivity using perfusion MRI. NeuroImage, 2008, 40, 1595-1605.	2.1	109

90 A 300MHz, 48mW analog front-end design for IEEE 802.3an 10GBase-T Ethernet. , 2008, , .

#	Article	IF	CITATIONS
91	RF power duty cycle restrictions of continuous arterial spin labeling coil for 7.0 tesla perfusion magnetic resonance imaging. , 2007, , .		1
92	Functional MRI detection of bilateral cortical reorganization in the rodent brain following peripheral nerve deafferentation. NeuroImage, 2007, 37, 262-273.	2.1	75
93	Improved cardiac manganese-enhanced MRI (MEMRI) with T <inf>1</inf> mapping in rodent. , 2007, , .		1
94	Dissociated roles of the middle frontal gyri in the processing of Chinese characters. NeuroReport, 2006, 17, 1397-1401.	0.6	42
95	Cell labeling for magnetic resonance imaging with theT1 agent manganese chloride. NMR in Biomedicine, 2006, 19, 50-59.	1.6	77
96	Improved neuronal tract tracing using manganese enhanced magnetic resonance imaging with fastT1 mapping. Magnetic Resonance in Medicine, 2006, 55, 604-611.	1.9	77
97	Application of model-free analysis in the MR assessment of pulmonary perfusion dynamics. Magnetic Resonance in Medicine, 2005, 54, 299-308.	1.9	7
98	Neuronal Specificity of Acupuncture Response: A fMRI Study with Electroacupuncture. NeuroImage, 2002, 16, 1028-1037.	2.1	258
99	Tissue segmentation-assisted analysis of fMRI for human motor response: an approach combining artificial neural network and fuzzy C means. Journal of Digital Imaging, 2001, 14, 38-47.	1.6	6
100	IMPACT: Image-based physiological artifacts estimation and correction technique for functional MRI. Magnetic Resonance in Medicine, 2001, 46, 344-353.	1.9	66
101	IMAGE ANALYSIS OF FUNCTIONAL MAGNETIC RESONANCE IMAGING. Biomedical Engineering - Applications, Basis and Communications, 2001, 13, 248-255.	0.3	4
102	Model-free functional MRI analysis using Kohonen clustering neural network and fuzzy C-means. IEEE Transactions on Medical Imaging, 1999, 18, 1117-1128.	5.4	141