

# Michael Gerard McDonell

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6401962/publications.pdf>

Version: 2024-02-01

86  
papers

1,532  
citations

304743

22  
h-index

395702

33  
g-index

88  
all docs

88  
docs citations

88  
times ranked

1610  
citing authors

#	ARTICLE	IF	CITATIONS
1	Randomized Controlled Trial of Contingency Management for Stimulant Use in Community Mental Health Patients With Serious Mental Illness. <i>American Journal of Psychiatry</i> , 2013, 170, 94-101.	7.2	108
2	Increased distance was associated with lower daily attendance to an opioid treatment program in Spokane County Washington. <i>Journal of Substance Abuse Treatment</i> , 2018, 93, 26-30.	2.8	63
3	Racial-Ethnic Disparities in First-Episode Psychosis Treatment Outcomes From the RAISE-ETP Study. <i>Psychiatric Services</i> , 2018, 69, 1138-1145.	2.0	61
4	A review of contingency management for the treatment of substance-use disorders: adaptation for underserved populations, use of experimental technologies, and personalized optimization strategies. <i>Substance Abuse and Rehabilitation</i> , 2018, Volume 9, 43-57.	4.8	60
5	The Effect of Serious Mental Illness on the Risk of Rehospitalization Among Patients With Diabetes. <i>Psychosomatics</i> , 2014, 55, 134-143.	2.5	58
6	A Randomized Controlled Trial of Ethyl Glucuronide-Based Contingency Management for Outpatients With Co-Occurring Alcohol Use Disorders and Serious Mental Illness. <i>American Journal of Psychiatry</i> , 2017, 174, 370-377.	7.2	54
7	Impact of tobacco, alcohol and cannabis use on treatment outcomes among patients experiencing first episode psychosis: Data from the national RAISE-ETP study. <i>Microbial Biotechnology</i> , 2019, 13, 142-146.	1.7	48
8	A Pilot Evaluation of Dialectical Behavioural Therapy in Adolescent Long-Term Inpatient Care. <i>Child and Adolescent Mental Health</i> , 2010, 15, 193-196.	3.5	47
9	Using ethyl glucuronide in urine to detect light and heavy drinking in alcohol dependent outpatients. <i>Drug and Alcohol Dependence</i> , 2015, 157, 184-187.	3.2	45
10	Early life trauma, post-traumatic stress disorder, and allostatic load in a sample of American Indian adults. <i>American Journal of Human Biology</i> , 2017, 29, e22943.	1.6	44
11	Disparities in Access to Opioid Treatment Programs and Office-Based Buprenorphine Treatment Across the Rural-Urban and Area Deprivation Continua: A US Nationwide Small Area Analysis. <i>Value in Health</i> , 2021, 24, 188-195.	0.3	44
12	User Experience Evaluation of a Smoking Cessation App in People With Serious Mental Illness. <i>Nicotine and Tobacco Research</i> , 2016, 18, 1032-1038.	2.6	43
13	An economic evaluation of a contingency-management intervention for stimulant use among community mental health patients with serious mental illness. <i>Drug and Alcohol Dependence</i> , 2015, 153, 293-299.	3.2	34
14	The dissemination and implementation of contingency management for substance use disorders: A systematic review.. <i>Psychology of Addictive Behaviors</i> , 2020, 34, 99-110.	2.1	33
15	Determining Ethyl Glucuronide Cutoffs When Detecting Self-Reported Alcohol Use in Addiction Treatment Patients. <i>Alcoholism: Clinical and Experimental Research</i> , 2015, 39, 905-910.	2.4	31
16	A "missing not at random" (MNAR) and "missing at random" (MAR) growth model comparison with a buprenorphine/naloxone clinical trial. <i>Addiction</i> , 2015, 110, 51-58.	3.3	29
17	Multiple-Family Group Treatment of Outpatients With Schizophrenia: Impact on Service Utilization. <i>Family Process</i> , 2006, 45, 359-373.	2.6	28
18	VOUCHER-BASED REINFORCEMENT FOR ALCOHOL ABSTINENCE USING THE ETHYL-GLUCURONIDE ALCOHOL BIOMARKER. <i>Journal of Applied Behavior Analysis</i> , 2012, 45, 161-165.	2.7	28

#	ARTICLE	IF	CITATIONS
19	Global Appraisal of Individual Needs Short Screener (GSS): Psychometric Properties and Performance as a Screening Measure in Adolescents. <i>American Journal of Drug and Alcohol Abuse</i> , 2009, 35, 157-160.	2.1	27
20	High levels of agreement between clinic-based ethyl glucuronide (EtG) immunoassays and laboratory-based mass spectrometry. <i>American Journal of Drug and Alcohol Abuse</i> , 2015, 41, 246-250.	2.1	26
21	Focus groups to increase the cultural acceptability of a contingency management intervention for American Indian and Alaska Native Communities. <i>Journal of Substance Abuse Treatment</i> , 2018, 90, 57-63.	2.8	26
22	Longitudinal missing data strategies for substance use clinical trials using generalized estimating equations: an example with a buprenorphine trial. <i>Human Psychopharmacology</i> , 2013, 28, 506-515.	1.5	23
23	Contingency management targeting abstinence is effective in reducing depressive and anxiety symptoms among crack cocaine-dependent individuals.. <i>Experimental and Clinical Psychopharmacology</i> , 2017, 25, 466-472.	1.8	22
24	Three-year retention in methadone opioid agonist treatment: A survival analysis of clients by dose, area deprivation, and availability of alcohol and cannabis outlets. <i>Drug and Alcohol Dependence</i> , 2018, 193, 63-68.	3.2	22
25	A culturally-tailored behavioral intervention trial for alcohol use disorders in three American Indian communities: Rationale, design, and methods. <i>Contemporary Clinical Trials</i> , 2016, 47, 93-100.	1.8	21
26	Utility of Point-of-care Urine Drug Tests in the Treatment of Primary Care Patients With Drug Use Disorders. <i>Journal of Addiction Medicine</i> , 2016, 10, 196-201.	2.6	20
27	Use of a Mental Health Center Collaborative Care Team to Improve Diabetes Care and Outcomes for Patients With Psychosis. <i>Psychiatric Services</i> , 2018, 69, 349-352.	2.0	19
28	Barriers to Metabolic Care for Adults with Serious Mental Illness: Provider Perspectives. <i>International Journal of Psychiatry in Medicine</i> , 2011, 41, 379-387.	1.8	18
29	Pilot Study of an Integrated Smartphone and Breathalyzer Contingency Management Intervention for Alcohol Use. <i>Journal of Addiction Medicine</i> , 2020, 14, 193-198.	2.6	18
30	Post-traumatic stress disorder among persons with HIV who engage in heavy alcohol consumption in southwestern Uganda. <i>BMC Psychiatry</i> , 2021, 21, 457.	2.6	18
31	Availability of licensed cannabis businesses in relation to area deprivation in Washington state: A spatiotemporal analysis of cannabis business presence between 2014 and 2017. <i>Drug and Alcohol Review</i> , 2019, 38, 790-797.	2.1	17
32	Spatial access to opioid treatment program and alcohol and cannabis outlets: analysis of missed doses of methadone during the first, second, and third 90 days of treatment. <i>American Journal of Drug and Alcohol Abuse</i> , 2020, 46, 78-87.	2.1	17
33	Preliminary Evaluation of Washington State's Early Intervention Program for First-Episode Psychosis. <i>Psychiatric Services</i> , 2020, 71, 228-235.	2.0	17
34	Effect of Incentives for Alcohol Abstinence in Partnership With 3 American Indian and Alaska Native Communities. <i>JAMA Psychiatry</i> , 2021, 78, 599.	11.0	16
35	Predictors of stimulant abuse treatment outcomes in severely mentally ill outpatients. <i>Drug and Alcohol Dependence</i> , 2013, 131, 162-165.	3.2	15
36	Sociodemographic Characteristics, Patterns of Crack Use, Concomitant Substance Use Disorders, and Psychiatric Symptomatology in Treatment-Seeking Crack-Dependent Individuals in Brazil. <i>Journal of Psychoactive Drugs</i> , 2018, 50, 367-372.	1.7	15

#	ARTICLE	IF	CITATIONS
37	Association of Co-Occurring Serious Mental Illness with Emergency Hospitalization in People with Chronic Kidney Disease. <i>American Journal of Nephrology</i> , 2014, 39, 260-267.	3.1	14
38	Family Experiences Prior to the Initiation of Care for First-Episode Psychosis: A Meta-Synthesis of Qualitative Studies. <i>Journal of Child and Family Studies</i> , 2020, 29, 2530-2541.	1.3	13
39	Preliminary findings: Contingency management targeting psycho-stimulant use results in secondary decreases in smoking for severely mentally ill adults. <i>American Journal on Addictions</i> , 2014, 23, 407-410.	1.4	12
40	The impact of early family contact on quality of life among non-Hispanic Blacks and Whites in the RAISE-ETP trial. <i>Schizophrenia Research</i> , 2020, 216, 523-525.	2.0	12
41	The rewarding recovery study: a randomized controlled trial of incentives for alcohol and drug abstinence with a rural American Indian community. <i>Addiction</i> , 2021, 116, 1569-1579.	3.3	12
42	Access to medication-assisted treatment in the United States: Comparison of travel time to opioid treatment programs and office-based buprenorphine treatment. <i>Drug and Alcohol Dependence</i> , 2021, 224, 108727.	3.2	12
43	Associations Between Physiological Signals Captured Using Wearable Sensors and Self-reported Outcomes Among Adults in Alcohol Use Disorder Recovery: Development and Usability Study. <i>JMIR Formative Research</i> , 2021, 5, e27891.	1.4	12
44	Pilot investigation: randomized-controlled analog trial for alcohol and tobacco smoking co-addiction using contingency management. <i>Behavioural Pharmacology</i> , 2018, 29, 462-468.	1.7	11
45	Crack cocaine users views regarding treatment with contingency management in Brazil. <i>Substance Abuse Treatment, Prevention, and Policy</i> , 2018, 13, 7.	2.2	11
46	Impact of first episode psychosis treatment on heavy cannabis use: Secondary analysis on RAISE-ETP study. <i>Schizophrenia Research</i> , 2019, 211, 86-87.	2.0	10
47	HIV and syphilis infections and associated factors among patients in treatment at a Specialist Alcohol, Tobacco, and Drugs Center in São Paulo—Brazil. <i>Trends in Psychiatry and Psychotherapy</i> , 2020, 42, 1-6.	0.8	10
48	Contingency management intervention targeting co-addiction of alcohol and drugs among American Indian adults: Design, methodology, and baseline data. <i>Clinical Trials</i> , 2018, 15, 587-599.	1.6	9
49	Understanding differences in family engagement and provider outreach in New Journeys: A coordinated specialty care program for first episode psychosis. <i>Psychiatry Research</i> , 2020, 291, 113286.	3.3	9
50	Pilot investigation of a phosphatidylethanol-based contingency management intervention targeting alcohol use.. <i>Psychology of Addictive Behaviors</i> , 2017, 31, 608-613.	2.1	9
51	A pilot study of the accuracy of onsite immunoassay urinalysis of illicit drug use in seriously mentally ill outpatients. <i>American Journal of Drug and Alcohol Abuse</i> , 2011, 37, 137-140.	2.1	8
52	Validation of the Co-occurring Disorder Quadrant Model. <i>Journal of Psychoactive Drugs</i> , 2012, 44, 266-273.	1.7	8
53	Decreases in smoking during treatment for methamphetamine-use disorders: preliminary evidence. <i>Behavioural Pharmacology</i> , 2018, 29, 370-374.	1.7	8
54	Homelessness Predicts Attrition but Not Alcohol Abstinence in Outpatients Experiencing Co-Occurring Alcohol Dependence and Serious Mental Illness. <i>Substance Abuse</i> , 2018, 39, 271-274.	2.3	8

#	ARTICLE	IF	CITATIONS
55	Effectiveness of incorporating contingency management into a public treatment program for people who use crack cocaine in Brazil. A single-blind randomized controlled trial. <i>International Journal of Drug Policy</i> , 2022, 99, 103464.	3.3	8
56	High Agreement Between Benchtop and Point-of-Care Dipcard Tests for Ethyl Glucuronide. <i>Therapeutic Drug Monitoring</i> , 2017, 39, 461-462.	2.0	7
57	Proximity to Screening Site, Rurality, and Neighborhood Disadvantage: Treatment Status among Individuals with Sexually Transmitted Infections in Yakima County, Washington. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 2679.	2.6	7
58	The Drinkersâ€™™ Intervention to Prevent Tuberculosis (DIPT) trial among heavy drinkers living with HIV in Uganda: study protocol of a 2Ã—2 factorial trial. <i>Trials</i> , 2021, 22, 355.	1.6	7
59	Clinical Prevalence and Correlates of Substance Use in Adolescent Psychiatric Emergency Patients. <i>Pediatric Emergency Care</i> , 2011, 27, 384-389.	0.9	6
60	Pretreatment ethyl glucuronide levels predict response to a contingency management intervention for alcohol use disorders among adults with serious mental illness. <i>American Journal on Addictions</i> , 2017, 26, 673-675.	1.4	6
61	Perspectives on a contingency management intervention for alcohol use among consumers with serious mental illness.. <i>Psychiatric Rehabilitation Journal</i> , 2019, 42, 26-31.	1.1	6
62	Variations in national availability of waived buprenorphine prescribers by racial and ethnic composition of zip codes. <i>Substance Abuse Treatment, Prevention, and Policy</i> , 2022, 17, .	2.2	6
63	Interaction between pre-treatment drug use and heterogeneity of psychiatric diagnosis predicts outcomes in outpatients with co-occurring disorders. <i>Psychiatry Research</i> , 2018, 260, 233-235.	3.3	5
64	Using a randomized controlled trial to test whether modifications to contingency management improve outcomes for heavy drinkers with serious mental illness. <i>Contemporary Clinical Trials</i> , 2018, 69, 92-98.	1.8	5
65	Cannabis use in first episode psychosis: what we have tried and why it hasnâ€™™t worked. <i>BMC Medicine</i> , 2019, 17, 194.	5.5	5
66	Mixed-methods trial of a phosphatidylethanol-based contingency management intervention to initiate and maintain alcohol abstinence in formerly homeless adults with alcohol use disorders. <i>Contemporary Clinical Trials Communications</i> , 2021, 22, 100757.	1.1	5
67	Predictors of tobacco and alcohol co-use from ages 15 to 32: The Amsterdam Growth and Health Longitudinal Study.. <i>Experimental and Clinical Psychopharmacology</i> , 2018, 26, 549-559.	1.8	5
68	Contingency management for alcohol use disorder reduces cannabis use among American Indian and Alaska Native adults. <i>Journal of Substance Abuse Treatment</i> , 2022, 137, 108693.	2.8	5
69	Prevalence and Correlates of Cannabis Use in Outpatients with Serious Mental Illness Receiving Treatment for Alcohol Use Disorders. <i>Cannabis and Cannabinoid Research</i> , 2017, 2, 133-138.	2.9	4
70	Serious Mental Illness in Heavy Drinkers Is Associated with Poor Treatment Outcomes in Outpatients with Co-occurring Disorders. <i>International Journal of Mental Health and Addiction</i> , 2018, 16, 672-679.	7.4	4
71	Prevalence and Indicators of Household Smoking Bans Among American Indians. <i>Journal of Community Health</i> , 2018, 43, 746-755.	3.8	4
72	Contingency management is effective in promoting abstinence and retention in treatment among crack cocaine users with a previous history of poor treatment response: a crossover trial. <i>Psicologia: Reflexao E Critica</i> , 2019, 32, 14.	0.9	4

#	ARTICLE	IF	CITATIONS
73	Onset of alcohol use disorder among alcohol initiates by race/ethnicity. <i>Alcohol</i> , 2021, 97, 13-21.	1.7	4
74	Assessing the Interest and Cultural Congruence of Contingency Management as an Intervention for Alcohol Misuse Among Younger American Indian Adults. <i>American Indian and Alaska Native Mental Health Research</i> , 2019, 26, 38-57.	0.8	4
75	Cannabis use and metabolic syndrome among clients with first episode psychosis. <i>Microbial Biotechnology</i> , 2021, 15, 1051-1055.	1.7	3
76	Geographic Disparities in Access to Specialty Care Programs for Early Psychosis in Washington State. <i>Administration and Policy in Mental Health and Mental Health Services Research</i> , 2022, 49, 5-12.	2.1	3
77	Development of Automated Reinforcement Management System (ARMS): Protocol for a Phase I Feasibility and Usability Study. <i>JMIR Formative Research</i> , 2021, 5, e25796.	1.4	3
78	Assessing the Predictive Validity of the Stages of Change Readiness and Treatment Eagerness Scale (SOCRATES) in Alaska Native and American Indian People. <i>Journal of Addiction Medicine</i> , 2020, 14, e241-e246.	2.6	3
79	Serious Mental Illness in Heavy Drinkers Is Associated with Poor Treatment Outcomes in Outpatients with Co-occurring Disorders. <i>International Journal of Mental Health and Addiction</i> , 2018, 16, 672-679.	7.4	3
80	Concordance Between Point-of-Care Urine Ethyl Glucuronide Alcohol Tests and Self-Reported Alcohol Use in Persons with HIV in Uganda. <i>AIDS and Behavior</i> , 2022, 26, 2539-2547.	2.7	3
81	Response to Urine Drug Testing in a Family Residency Practice. <i>Journal of Addiction Medicine</i> , 2017, 11, 243.	2.6	1
82	Developing and implementing a culturally informed Family Motivational Engagement Strategy (FAMES) to increase family engagement in first episode psychosis programs: mixed methods pilot study protocol. <i>BMJ Open</i> , 2020, 10, e036907.	1.9	1
83	Lessons Shared: Implementation of a Randomized Clinical Trial for Alcohol Use Disorders with Five American Indian and Alaska Native Communities. <i>Collaborations (Coral Gables, Fla )</i> , 2021, 4, .	0.1	1
84	Automated Reinforcement Management System (ARMS): focused phase I provider feedback. <i>Addiction Science &amp; Clinical Practice</i> , 2022, 17, 20.	2.6	1
85	Rising strong: a housing-based intervention to prevent childhood out-of-home placements from families impacted by housing instability, opioid use, and other substance use disorders. <i>Social Work in Mental Health</i> , 2021, 19, 542-557.	1.0	0
86	Contingency Management as a Behavioral Approach in Addiction Treatment. , 2021, , 417-432.		0