
## Michael Gerard McDonell

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/6401962/publications.pdf

Version: 2024-02-01



#	Article	IF	CITATIONS
1	Geographic Disparities in Access to Specialty Care Programs for Early Psychosis in Washington State. Administration and Policy in Mental Health and Mental Health Services Research, 2022, 49, 5-12.	2.1	3
2	Effectiveness of incorporating contingency management into a public treatment program for people who use crack cocaine in Brazil. A single-blind randomized controlled trial. International Journal of Drug Policy, 2022, 99, 103464.	3.3	8
3	Concordance Between Point-of-Care Urine Ethyl Glucuronide Alcohol Tests and Self-Reported Alcohol Use in Persons with HIV in Uganda. AIDS and Behavior, 2022, 26, 2539-2547.	2.7	3
4	Contingency management for alcohol use disorder reduces cannabis use among American Indian and Alaska Native adults. Journal of Substance Abuse Treatment, 2022, 137, 108693.	2.8	5
5	Automated Reinforcement Management System (ARMS): focused phase I provider feedback. Addiction Science & Clinical Practice, 2022, 17, 20.	2.6	1
6	Variations in national availability of waivered buprenorphine prescribers by racial and ethnic composition of zip codes. Substance Abuse Treatment, Prevention, and Policy, 2022, 17, .	2.2	6
7	Cannabis use and metabolic syndrome among clients with first episode psychosis. Microbial Biotechnology, 2021, 15, 1051-1055.	1.7	3
8	Disparities in Access to Opioid Treatment Programs and Office-Based Buprenorphine Treatment Across the Rural-Urban and Area Deprivation Continua: A US Nationwide Small Area Analysis. Value in Health, 2021, 24, 188-195.	0.3	44
9	The rewarding recovery study: a randomized controlled trial of incentives for alcohol and drug abstinence with a rural American Indian community. Addiction, 2021, 116, 1569-1579.	3.3	12
10	Lessons Shared: Implementation of a Randomized Clinical Trial for Alcohol Use Disorders with Five American Indian and Alaska Native Communities. Collaborations (Coral Gables, Fla ), 2021, 4, .	0.1	1
11	The Drinkers' Intervention to Prevent Tuberculosis (DIPT) trial among heavy drinkers living with HIV in Uganda: study protocol of a 2×2 factorial trial. Trials, 2021, 22, 355.	1.6	7
12	Rising strong: a housing-based intervention to prevent childhood out-of-home placements from families impacted by housing instability, opioid use, and other substance use disorders. Social Work in Mental Health, 2021, 19, 542-557.	1.0	0
13	Mixed-methods trial of a phosphatidylethanol-based contingency management intervention to initiate and maintain alcohol abstinence in formerly homeless adults with alcohol use disorders. Contemporary Clinical Trials Communications, 2021, 22, 100757.	1.1	5
14	Effect of Incentives for Alcohol Abstinence in Partnership With 3 American Indian and Alaska Native Communities. JAMA Psychiatry, 2021, 78, 599.	11.0	16
15	Access to medication-assisted treatment in the United States: Comparison of travel time to opioid treatment programs and office-based buprenorphine treatment. Drug and Alcohol Dependence, 2021, 224, 108727.	3.2	12
16	Associations Between Physiological Signals Captured Using Wearable Sensors and Self-reported Outcomes Among Adults in Alcohol Use Disorder Recovery: Development and Usability Study. JMIR Formative Research, 2021, 5, e27891.	1.4	12
17	Development of Automated Reinforcement Management System (ARMS): Protocol for a Phase I Feasibility and Usability Study. JMIR Formative Research, 2021, 5, e25796.	1.4	3
18	Onset of alcohol use disorder among alcohol initiates by race/ethnicity. Alcohol, 2021, 97, 13-21.	1.7	4

#	Article	IF	CITATIONS
19	Post-traumatic stress disorder among persons with HIV who engage in heavy alcohol consumption in southwestern Uganda. BMC Psychiatry, 2021, 21, 457.	2.6	18
20	Contingency Management as a Behavioral Approach in Addiction Treatment. , 2021, , 417-432.		0
21	Spatial access to opioid treatment program and alcohol and cannabis outlets: analysis of missed doses of methadone during the first, second, and third 90 days of treatment. American Journal of Drug and Alcohol Abuse, 2020, 46, 78-87.	2.1	17
22	Family Experiences Prior to the Initiation of Care for First-Episode Psychosis: A Meta-Synthesis of Qualitative Studies. Journal of Child and Family Studies, 2020, 29, 2530-2541.	1.3	13
23	Pilot Study of an Integrated Smartphone and Breathalyzer Contingency Management Intervention for Alcohol Use. Journal of Addiction Medicine, 2020, 14, 193-198.	2.6	18
24	Preliminary Evaluation of Washington State's Early Intervention Program for First-Episode Psychosis. Psychiatric Services, 2020, 71, 228-235.	2.0	17
25	Understanding differences in family engagement and provider outreach in New Journeys: A coordinated specialty care program for first episode psychosis. Psychiatry Research, 2020, 291, 113286.	3.3	9
26	Developing and implementing a culturally informed FAmily Motivational Engagement Strategy (FAMES) to increase family engagement in first episode psychosis programs: mixed methods pilot study protocol. BMJ Open, 2020, 10, e036907.	1.9	1
27	The impact of early family contact on quality of life among non-Hispanic Blacks and Whites in the RAISE-ETP trial. Schizophrenia Research, 2020, 216, 523-525.	2.0	12
28	Proximity to Screening Site, Rurality, and Neighborhood Disadvantage: Treatment Status among Individuals with Sexually Transmitted Infections in Yakima County, Washington. International Journal of Environmental Research and Public Health, 2020, 17, 2679.	2.6	7
29	The dissemination and implementation of contingency management for substance use disorders: A systematic review Psychology of Addictive Behaviors, 2020, 34, 99-110.	2.1	33
30	Assessing the Predictive Validity of the Stages of Change Readiness and Treatment Eagerness Scale (SOCRATES) in Alaska Native and American Indian People. Journal of Addiction Medicine, 2020, 14, e241-e246.	2.6	3
31	HIV and syphilis infections and associated factors among patients in treatment at a Specialist Alcohol, Tobacco, and Drugs Center in São Paulo's "Cracolândia― Trends in Psychiatry and Psychotherapy, 2020, 42, 1-6.	0.8	10
32	Impact of first episode psychosis treatment on heavy cannabis use: Secondary analysis on RAISE-ETP study. Schizophrenia Research, 2019, 211, 86-87.	2.0	10
33	Cannabis use in first episode psychosis: what we have tried and why it hasn't worked. BMC Medicine, 2019, 17, 194.	5.5	5
34	Availability of licensed cannabis businesses in relation to area deprivation in Washington state: A spatiotemporal analysis of cannabis business presence between 2014 and 2017. Drug and Alcohol Review, 2019, 38, 790-797.	2.1	17
35	Contingency management is effective in promoting abstinence and retention in treatment among crack cocaine users with a previous history of poor treatment response: a crossover trial. Psicologia: Reflexao E Critica, 2019, 32, 14.	0.9	4
36	Impact of tobacco, alcohol and cannabis use on treatment outcomes among patients experiencing first episode psychosis: Data from the national RAISEâ€ETP study. Microbial Biotechnology, 2019, 13, 142-146.	1.7	48

#	Article	IF	CITATIONS
37	Perspectives on a contingency management intervention for alcohol use among consumers with serious mental illness Psychiatric Rehabilitation Journal, 2019, 42, 26-31.	1.1	6
38	Assessing the Interest and Cultural Congruence of Contingency Management as an Intervention for Alcohol Misuse Among Younger American Indian Adults. American Indian and Alaska Native Mental Health Research, 2019, 26, 38-57.	0.8	4
39	Serious Mental Illness in Heavy Drinkers Is Associated with Poor Treatment Outcomes in Outpatients with Co-occurring Disorders. International Journal of Mental Health and Addiction, 2018, 16, 672-679.	7.4	4
40	Pilot investigation: randomized-controlled analog trial for alcohol and tobacco smoking co-addiction using contingency management. Behavioural Pharmacology, 2018, 29, 462-468.	1.7	11
41	Use of a Mental Health Center Collaborative Care Team to Improve Diabetes Care and Outcomes for Patients With Psychosis. Psychiatric Services, 2018, 69, 349-352.	2.0	19
42	Prevalence and Indicators of Household Smoking Bans Among American Indians. Journal of Community Health, 2018, 43, 746-755.	3.8	4
43	Sociodemographic Characteristics, Patterns of Crack Use, Concomitant Substance Use Disorders, and Psychiatric Symptomatology in Treatment-Seeking Crack-Dependent Individuals in Brazil. Journal of Psychoactive Drugs, 2018, 50, 367-372.	1.7	15
44	Decreases in smoking during treatment for methamphetamine-use disorders: preliminary evidence. Behavioural Pharmacology, 2018, 29, 370-374.	1.7	8
45	Interaction between pre-treatment drug use and heterogeneity of psychiatric diagnosis predicts outcomes in outpatients with co-occurring disorders. Psychiatry Research, 2018, 260, 233-235.	3.3	5
46	Homelessness Predicts Attrition but Not Alcohol Abstinence in Outpatients Experiencing Co-Occurring Alcohol Dependence and Serious Mental Illness. Substance Abuse, 2018, 39, 271-274.	2.3	8
47	Three-year retention in methadone opioid agonist treatment: A survival analysis of clients by dose, area deprivation, and availability of alcohol and cannabis outlets. Drug and Alcohol Dependence, 2018, 193, 63-68.	3.2	22
48	Contingency management intervention targeting co-addiction of alcohol and drugs among American Indian adults: Design, methodology, and baseline data. Clinical Trials, 2018, 15, 587-599.	1.6	9
49	A review of contingency management for the treatment of substance-use disorders: adaptation for underserved populations, use of experimental technologies, and personalized optimization strategies. Substance Abuse and Rehabilitation, 2018, Volume 9, 43-57.	4.8	60
50	Racial-Ethnic Disparities in First-Episode Psychosis Treatment Outcomes From the RAISE-ETP Study. Psychiatric Services, 2018, 69, 1138-1145.	2.0	61
51	Using a randomized controlled trial to test whether modifications to contingency management improve outcomes for heavy drinkers with serious mental illness. Contemporary Clinical Trials, 2018, 69, 92-98.	1.8	5
52	Increased distance was associated with lower daily attendance to an opioid treatment program in Spokane County Washington. Journal of Substance Abuse Treatment, 2018, 93, 26-30.	2.8	63
53	Crack cocaine users views regarding treatment with contingency management in Brazil. Substance Abuse Treatment, Prevention, and Policy, 2018, 13, 7.	2.2	11
54	Focus groups to increase the cultural acceptability of a contingency management intervention for American Indian and Alaska Native Communities. Journal of Substance Abuse Treatment, 2018, 90, 57-63.	2.8	26

#	Article	IF	CITATIONS
55	Predictors of tobacco and alcohol co-use from ages 15 to 32: The Amsterdam Growth and Health Longitudinal Study Experimental and Clinical Psychopharmacology, 2018, 26, 549-559.	1.8	5
56	Serious Mental Illness in Heavy Drinkers Is Associated with Poor Treatment Outcomes in Outpatients with Co-occurring Disorders. International Journal of Mental Health and Addiction, 2018, 16, 672-679.	7.4	3
57	A Randomized Controlled Trial of Ethyl Glucuronide-Based Contingency Management for Outpatients With Co-Occurring Alcohol Use Disorders and Serious Mental Illness. American Journal of Psychiatry, 2017, 174, 370-377.	7.2	54
58	High Agreement Between Benchtop and Point-of-Care Dipcard Tests for Ethyl Glucuronide. Therapeutic Drug Monitoring, 2017, 39, 461-462.	2.0	7
59	Early life trauma, postâ€traumatic stress disorder, and allostatic load in a sample of American Indian adults. American Journal of Human Biology, 2017, 29, e22943.	1.6	44
60	Pretreatment ethyl glucuronide levels predict response to a contingency management intervention for alcohol use disorders among adults with serious mental illness. American Journal on Addictions, 2017, 26, 673-675.	1.4	6
61	Prevalence and Correlates of Cannabis Use in Outpatients with Serious Mental Illness Receiving Treatment for Alcohol Use Disorders. Cannabis and Cannabinoid Research, 2017, 2, 133-138.	2.9	4
62	Response to Urine Drug Testing in a Family Residency Practice. Journal of Addiction Medicine, 2017, 11, 243.	2.6	1
63	Contingency management targeting abstinence is effective in reducing depressive and anxiety symptoms among crack cocaine-dependent individuals Experimental and Clinical Psychopharmacology, 2017, 25, 466-472.	1.8	22
64	Pilot investigation of a phosphatidylethanol-based contingency management intervention targeting alcohol use Psychology of Addictive Behaviors, 2017, 31, 608-613.	2.1	9
65	Utility of Point-of-care Urine Drug Tests in the Treatment of Primary Care Patients With Drug Use Disorders. Journal of Addiction Medicine, 2016, 10, 196-201.	2.6	20
66	A culturally-tailored behavioral intervention trial for alcohol use disorders in three American Indian communities: Rationale, design, and methods. Contemporary Clinical Trials, 2016, 47, 93-100.	1.8	21
67	User Experience Evaluation of a Smoking Cessation App in People With Serious Mental Illness. Nicotine and Tobacco Research, 2016, 18, 1032-1038.	2.6	43
68	Determining Ethyl Glucuronide Cutoffs When Detecting Selfâ€Reported Alcohol Use in Addiction Treatment Patients. Alcoholism: Clinical and Experimental Research, 2015, 39, 905-910.	2.4	31
69	Using ethyl glucuronide in urine to detect light and heavy drinking in alcohol dependent outpatients. Drug and Alcohol Dependence, 2015, 157, 184-187.	3.2	45
70	An economic evaluation of a contingency-management intervention for stimulant use among community mental health patients with serious mental illness. Drug and Alcohol Dependence, 2015, 153, 293-299.	3.2	34
71	High levels of agreement between clinic-based ethyl glucuronide (EtG) immunoassays and laboratory-based mass spectrometry. American Journal of Drug and Alcohol Abuse, 2015, 41, 246-250.	2.1	26
72	A â€~missing not at random' ( <scp>MNAR</scp> ) and â€~missing at random' ( <scp>MAR</scp> ) growth	mggel	29

comparison with a buprenorphine/naloxone clinical trial. Addiction, 2015, 110, 51-58.

#	Article	IF	CITATIONS
73	Preliminary findings: Contingency management targeting psycho-stimulant use results in secondary decreases in smoking for severely mentally ill adults. American Journal on Addictions, 2014, 23, 407-410.	1.4	12
74	The Effect of Serious Mental Illness on the Risk of Rehospitalization Among Patients With Diabetes. Psychosomatics, 2014, 55, 134-143.	2.5	58
75	Association of Co-Occurring Serious Mental Illness with Emergency Hospitalization in People with Chronic Kidney Disease. American Journal of Nephrology, 2014, 39, 260-267.	3.1	14
76	Longitudinal missing data strategies for substance use clinical trials using generalized estimating equations: an example with a buprenorphine trial. Human Psychopharmacology, 2013, 28, 506-515.	1.5	23
77	Predictors of stimulant abuse treatment outcomes in severely mentally ill outpatients. Drug and Alcohol Dependence, 2013, 131, 162-165.	3.2	15
78	Randomized Controlled Trial of Contingency Management for Stimulant Use in Community Mental Health Patients With Serious Mental Illness. American Journal of Psychiatry, 2013, 170, 94-101.	7.2	108
79	Validation of the Co-occurring Disorder Quadrant Model. Journal of Psychoactive Drugs, 2012, 44, 266-273.	1.7	8
80	VOUCHERâ€BASED REINFORCEMENT FOR ALCOHOL ABSTINENCE USING THE ETHYLâ€GLUCURONIDE ALCOHOL BIOMARKER. Journal of Applied Behavior Analysis, 2012, 45, 161-165.	2.7	28
81	Clinical Prevalence and Correlates of Substance Use in Adolescent Psychiatric Emergency Patients. Pediatric Emergency Care, 2011, 27, 384-389.	0.9	6
82	Barriers to Metabolic Care for Adults with Serious Mental Illness: Provider Perspectives. International Journal of Psychiatry in Medicine, 2011, 41, 379-387.	1.8	18
83	A pilot study of the accuracy of onsite immunoassay urinalysis of illicit drug use in seriously mentally ill outpatients. American Journal of Drug and Alcohol Abuse, 2011, 37, 137-140.	2.1	8
84	A Pilot Evaluation of Dialectical Behavioural Therapy in Adolescent Longâ€Term Inpatient Care. Child and Adolescent Mental Health, 2010, 15, 193-196.	3.5	47
85	Global Appraisal of Individual Needs Short Screener (GSS): Psychometric Properties and Performance as a Screening Measure in Adolescents. American Journal of Drug and Alcohol Abuse, 2009, 35, 157-160.	2.1	27
86	Multiple-Family Group Treatment of Outpatients With Schizophrenia: Impact on Service Utilization. Family Process, 2006, 45, 359-373.	2.6	28