

Jung Hun Kang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6403/publications.pdf>

Version: 2024-02-01

75
papers

1,028
citations

430843

18
h-index

501174

28
g-index

81
all docs

81
docs citations

81
times ranked

1786
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Safe use of opioids. Journal of the Korean Medical Association, 2022, 65, 57-65. | 0.3 | 1 |
| 2 | Palliative Sedation in End-of-Life Patients in Eastern Asia: A Narrative Review. Cancer Research and Treatment, 2022, 54, 644-650. | 3.0 | 3 |
| 3 | Modified FOLFIRINOX as a Second-Line Treatment for Patients with Gemcitabine-Failed Advanced Biliary Tract Cancer: A Prospective Multicenter Phase II Study. Cancers, 2022, 14, 1950. | 3.7 | 3 |
| 4 | Survey of Opioid Risk Tool Among Cancer Patients Receiving Opioid Analgesics. Journal of Korean Medical Science, 2022, 37, . | 2.5 | 0 |
| 5 | Modified FOLFIRINOX versus S-1 as second-line chemotherapy in patients with gemcitabine-failed metastatic pancreatic cancer: A randomized phase III trial (MPACA-3).. Journal of Clinical Oncology, 2021, 39, 4119-4119. | 1.6 | 1 |
| 6 | Korean Version of the Patient Dignity Inventory: Translation and Validation in Patients With Advanced Cancer. Journal of Pain and Symptom Management, 2021, 62, 416-424.e2. | 1.2 | 0 |
| 7 | Modified FOLFIRINOX versus S-1 as second-line chemotherapy in gemcitabine-failed metastatic pancreatic cancer patients: A randomised controlled trial (MPACA-3). European Journal of Cancer, 2021, 157, 21-30. | 2.8 | 18 |
| 8 | Cachexia index as a potential biomarker for cancer cachexia and a prognostic indicator in diffuse large B-cell lymphoma. Journal of Cachexia, Sarcopenia and Muscle, 2021, 12, 2211-2219. | 7.3 | 22 |
| 9 | Community Care for Cancer Patients in Rural Areas: An Integrated Regional Cancer Center and Public Health Center Partnership Model. The Korean Journal of Hospice and Palliative Care, 2021, 24, 226-234. | 0.7 | 2 |
| 10 | A Randomized Controlled Trial of Epidermal Growth Factor Ointment for Treating Epidermal Growth Factor Receptor Inhibitor-Induced Skin Toxicities. Oncologist, 2020, 25, e186-e193. | 3.7 | 10 |
| 11 | Association of illness understanding with advance care planning and end-of-life care preferences for advanced cancer patients and their family members. Supportive Care in Cancer, 2020, 28, 2959-2967. | 2.2 | 25 |
| 12 | Debatable Issues in Palliative Sedation; Authors' Reply to the Response by Robert Twycross. Journal of Pain and Symptom Management, 2020, 59, e2-e3. | 1.2 | 0 |
| 13 | rhEGF Treatment Improves EGFR Inhibitor-Induced Skin Barrier and Immune Defects. Cancers, 2020, 12, 3120. | 3.7 | 8 |
| 14 | Regorafenib in patients with advanced Child-Pugh B hepatocellular carcinoma: A multicentre retrospective study. Liver International, 2020, 40, 2544-2552. | 3.9 | 32 |
| 15 | Interference with daily functioning by breakthrough pain in patients with cancer. Supportive Care in Cancer, 2020, 28, 5177-5183. | 2.2 | 8 |
| 16 | The Impacts of Prognostic Awareness on Mood and Quality of Life Among Patients With Advanced Cancer. American Journal of Hospice and Palliative Medicine, 2020, 37, 904-912. | 1.4 | 18 |
| 17 | Clinical Practice Guideline for Care in the Last Days of Life. The Korean Journal of Hospice and Palliative Care, 2020, 23, 103-113. | 0.7 | 6 |
| 18 | Liposomal irinotecan plus fluorouracil/leucovorin versus FOLFIRINOX as the second-line chemotherapy for patients with metastatic pancreatic cancer: Multicenter study of the Korean Cancer Study Group (KCSG).. Journal of Clinical Oncology, 2020, 38, 4624-4624. | 1.6 | 0 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | The prognostic significance of the advanced lung cancer inflammation index(ALI) in patients with advanced biliary tract cancer: A retrospective study.. Journal of Clinical Oncology, 2020, 38, e16613-e16613. | 1.6 | 0 |
| 20 | The Attitudes of Physicians and the General Public toward Prognostic Disclosure of Different Serious Illnesses: a Korean Nationwide Study. Journal of Korean Medical Science, 2020, 35, e401. | 2.5 | 0 |
| 21 | Clinical Patterns of Continuous and Intermittent Palliative Sedation in Patients With Terminal Cancer: A Descriptive, Observational Study. Journal of Pain and Symptom Management, 2019, 58, 65-71. | 1.2 | 17 |
| 22 | Attitudes of the General Public, Cancer Patients, Family Caregivers, and Physicians Toward Advance Care Planning: A Nationwide Survey Before the Enforcement of the Life-Sustaining Treatment Decision-Making Act. Journal of Pain and Symptom Management, 2019, 57, 774-782. | 1.2 | 25 |
| 23 | Sedation for terminally ill cancer patients. Medicine (United States), 2019, 98, e14278. | 1.0 | 12 |
| 24 | Capecitabine plus Oxaliplatin as a Second-Line Therapy for Advanced Biliary Tract Cancers: A Multicenter, Open-Label, Phase II Trial. Journal of Cancer, 2019, 10, 6185-6190. | 2.5 | 7 |
| 25 | Safety and efficacy of trastuzumab administered as a 30-min infusion in patients with HER2-positive advanced gastric cancer. Cancer Chemotherapy and Pharmacology, 2019, 83, 501-508. | 2.3 | 6 |
| 26 | Preference and Performance Fidelity of Modified Korean Physician Order for Life-Sustaining Treatment (MK-POLST) Items in Hospice Patients with Cancer. The Korean Journal of Hospice and Palliative Care, 2019, 22, 198-206. | 0.7 | 1 |
| 27 | Priorities of a "good death" according to cancer patients, their family caregivers, physicians, and the general population: a nationwide survey. Supportive Care in Cancer, 2018, 26, 3479-3488. | 2.2 | 46 |
| 28 | Attitudes toward early palliative care in cancer patients and caregivers: a Korean nationwide survey. Cancer Medicine, 2018, 7, 1784-1793. | 2.8 | 22 |
| 29 | Development and validation of the quality care questionnaire "palliative care (QCQ-PC): patient-reported assessment of quality of palliative care. BMC Palliative Care, 2018, 17, 40. | 1.8 | 13 |
| 30 | The impact of caregiver's role preference on decisional conflicts and psychiatric distresses in decision making to help caregiver's disclosure of terminal disease status. Quality of Life Research, 2018, 27, 1571-1581. | 3.1 | 6 |
| 31 | Using primary tumor volumetry to predict treatment outcome for patients with oropharyngeal cancer who were treated with definitive chemoradiotherapy. Asia-Pacific Journal of Clinical Oncology, 2018, 14, e21-e28. | 1.1 | 1 |
| 32 | Discordance between Physician and the General Public Perceptions of Prognostic Disclosure to Children with Serious Illness: a Korean Nationwide Study. Journal of Korean Medical Science, 2018, 33, e327. | 2.5 | 7 |
| 33 | Compassion is a key quality for palliative care teams. Cancer Medicine, 2018, 7, 5325-5326. | 2.8 | 2 |
| 34 | Comparison of attitudes towards five end-of-life care interventions (active pain control, withdrawal) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5 caregivers, physicians and the general Korean population. BMJ Open, 2018, 8, e020519. | 1.9 | 20 |
| 35 | Attenuated FOLFIRINOX in the salvage treatment of gemcitabine-refractory advanced pancreatic cancer: a phase II study. Cancer Communications, 2018, 38, 1-8. | 9.2 | 23 |
| 36 | Craving Behavior from Opioid Addiction Controlled with Olanzapine in an Advanced Cancer Patient: A Case Report. Journal of Palliative Medicine, 2018, 21, 1367-1370. | 1.1 | 8 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Retrospective analysis of palliative chemotherapy for the patients with bladder adenocarcinoma: Korean Cancer Study Group Genitourinary and Gynecology Cancer Committee. Korean Journal of Internal Medicine, 2018, 33, 383-390. | 1.7 | 9 |
| 38 | Pazopanib for the Treatment of Non-clear Cell Renal Cell Carcinoma: A Single-Arm, Open-Label, Multicenter, Phase II Study. Cancer Research and Treatment, 2018, 50, 488-494. | 3.0 | 28 |
| 39 | Pembrolizumab for Refractory Metastatic Myxofibrosarcoma: A Case Report. Cancer Research and Treatment, 2018, 50, 1458-1461. | 3.0 | 13 |
| 40 | Successful Treatment with Transdermal Buprenorphine Patch in Opioid-Dependent Cancer Patients: Case Series. The Korean Journal of Hospice and Palliative Care, 2018, 21, 152-157. | 0.7 | 1 |
| 41 | Qualitative Study on the Perceptions of Terminally Ill Cancer Patients and Their Family Members Regarding End-of-Life Experiences Focusing on Palliative Sedation. Journal of Pain and Symptom Management, 2017, 53, 1010-1016. | 1.2 | 17 |
| 42 | Efficacy and Safety of FOLFIRI Regimen in Elderly Versus Nonelderly Patients with Metastatic Colorectal or Gastric Cancer. Oncologist, 2017, 22, 293-303. | 3.7 | 5 |
| 43 | Antiemetic Corticosteroid Rotation from Dexamethasone to Methylprednisolone to Prevent Dexamethasone-Induced Hiccup in Cancer Patients Treated with Chemotherapy: A Randomized, Single-Blind, Crossover Phase III Trial. Oncologist, 2017, 22, 1354-1361. | 3.7 | 24 |
| 44 | Molecular characterization of urothelial carcinoma of bladder and upper urinary tract.. Journal of Clinical Oncology, 2017, 35, 361-361. | 1.6 | 1 |
| 45 | A comparison of pectoralis versus lumbar skeletal muscle indices for defining sarcopenia in diffuse large B-cell lymphoma - two are better than one. Oncotarget, 2017, 8, 47007-47019. | 1.8 | 26 |
| 46 | Factors Influencing Clinicians' Choice of Adjuvant S-1 versus Capecitabine plus Oxaliplatin after Curative Gastrectomy in Patients with Gastric Cancer. Journal of Cancer, 2016, 7, 1711-1715. | 2.5 | 6 |
| 47 | Chemotherapy vs supportive care alone for relapsed gastric, gastroesophageal junction, and oesophageal adenocarcinoma: a meta-analysis of patient-level data. British Journal of Cancer, 2016, 114, 381-387. | 6.4 | 56 |
| 48 | Phase II trial of epidermal growth factor ointment for patients with Erlotinib-related skin effects. Supportive Care in Cancer, 2016, 24, 301-309. | 2.2 | 11 |
| 49 | Gender Differences in Hiccup Patients: Analysis of Published Case Reports and Case-Control Studies. Journal of Pain and Symptom Management, 2016, 51, 278-283. | 1.2 | 25 |
| 50 | Chemotherapy versus best supportive care in patients with advanced biliary tract carcinoma: A propensity score-matched analysis.. Journal of Clinical Oncology, 2016, 34, 316-316. | 1.6 | 1 |
| 51 | Is There a Role for Adjuvant Therapy in R0 Resected Gallbladder Cancer?: A Propensity Score-Matched Analysis. Cancer Research and Treatment, 2016, 48, 1274-1285. | 3.0 | 14 |
| 52 | Role of adjuvant therapy after R0 resection for patients with distal cholangiocarcinoma.. Journal of Clinical Oncology, 2016, 34, 355-355. | 1.6 | 4 |
| 53 | Attenuated FOLFIRINOX (5-FU/LV, irinotecan and oxaliplatin) as second-line chemotherapy for the Koreans (Asian population) with gemcitabine-resistant or "refractory advanced pancreatic cancer.. Journal of Clinical Oncology, 2016, 34, e15731-e15731. | 1.6 | 0 |
| 54 | Changes in health-related quality of life and quality of care among terminally ill cancer patients and survival prediction: Multicenter prospective cohort study. Palliative and Supportive Care, 2015, 13, 1103-1111. | 1.0 | 10 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Intraocular lymphoma in Korea: the Consortium for Improving Survival of Lymphoma (CISL) study. <i>Blood Research</i> , 2015, 50, 242. | 1.3 | 15 |
| 56 | Long-Term Intermittent Palliative Sedation for Refractory Symptoms at the End of Life in Two Cancer Patients. <i>Journal of Palliative Medicine</i> , 2015, 18, 807-810. | 1.1 | 16 |
| 57 | Minimal clinically important differences in the Edmonton Symptom Assessment Scale in cancer patients: A prospective study.. <i>Journal of Clinical Oncology</i> , 2015, 33, 9601-9601. | 1.6 | 0 |
| 58 | Comparison of Failure Patterns Between Conventional and Intensity-modulated Radiotherapy for Stage III and IV Head and Neck Squamous Cell Carcinoma. <i>Anticancer Research</i> , 2015, 35, 6833-40. | 1.1 | 5 |
| 59 | The Understanding of Terminal Cancer and Its Relationship with Attitudes toward End-of-Life Care Issues. <i>Medical Decision Making</i> , 2014, 34, 720-730. | 2.4 | 11 |
| 60 | Simvastatin plus capecitabine+ cisplatin versus placebo plus capecitabine+ cisplatin in patients with previously untreated advanced gastric cancer: A double-blind randomised phase 3 study. <i>European Journal of Cancer</i> , 2014, 50, 2822-2830. | 2.8 | 79 |
| 61 | Physical Symptoms and Psychiatric, Social, Spiritual and Economical Care Needs of Patients under Home-based Cancer Service. <i>The Korean Journal of Hospice and Palliative Care</i> , 2014, 17, 216-222. | 0.7 | 6 |
| 62 | Clinical Usefulness of Hydromorphone-OROS in Improving Sleep Disturbances in Korean Cancer Patients: A Multicenter, Prospective, Open-Label Study. <i>Cancer Research and Treatment</i> , 2014, 46, 331-338. | 3.0 | 6 |
| 63 | Natural history of untreated patients who had metastatic biliary tract cancer (BTC) with good performance status (PS).. <i>Journal of Clinical Oncology</i> , 2014, 32, e15106-e15106. | 1.6 | 0 |
| 64 | Clinical significance of the neutrophil+lymphocyte ratio in venous thromboembolism patients with lung cancer.. <i>Journal of Clinical Oncology</i> , 2014, 32, e19118-e19118. | 1.6 | 0 |
| 65 | The prognostic impact of the neutrophil-to-lymphocyte ratio in patients with small-cell lung cancer.. <i>Journal of Clinical Oncology</i> , 2014, 32, e18550-e18550. | 1.6 | 0 |
| 66 | The attitudes of Korean cancer patients, family caregivers, oncologists, and members of the general public toward advance directives. <i>Supportive Care in Cancer</i> , 2013, 21, 1437-1444. | 2.2 | 48 |
| 67 | Comprehensive approaches to managing delirium in patients with advanced cancer. <i>Cancer Treatment Reviews</i> , 2013, 39, 105-112. | 7.7 | 39 |
| 68 | Changes in Symptom Intensity Among Cancer Patients Receiving Outpatient Palliative Care. <i>Journal of Pain and Symptom Management</i> , 2013, 46, 652-660. | 1.2 | 68 |
| 69 | Treatment of Dexamethasone-Induced Hiccup in Chemotherapy Patients by Methylprednisolone Rotation. <i>Oncologist</i> , 2013, 18, 1229-1234. | 3.7 | 28 |
| 70 | The attitudes of Korean cancer patients, family caregivers, oncologists, and members of the general public toward advance directives. , 2013, 21, 1437. | | 1 |
| 71 | Corticosteroid Rotation to Alleviate Dexamethasone-Induced Hiccup: A Case Series at a Single Institution. <i>Journal of Pain and Symptom Management</i> , 2012, 43, 625-630. | 1.2 | 31 |
| 72 | Targeted agent use in patients with cancer at the end of life.. <i>Journal of Clinical Oncology</i> , 2012, 30, 215-215. | 1.6 | 1 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Pilot study of irinotecan/oxaliplatin (IROX) combination chemotherapy for patients with gemcitabine- and 5-fluorouracil- refractory pancreatic cancer. <i>Investigational New Drugs</i> , 2010, 28, 343-349. | 2.6 | 15 |
| 74 | Phase II trial of S-1 in combination with gemcitabine for chemo-naïve patients with locally advanced or metastatic pancreatic cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2009, 64, 707-713. | 2.3 | 28 |
| 75 | Clinical Features and Treatment Outcomes of Primary Testicular Lymphoma.. <i>Blood</i> , 2006, 108, 4670-4670. | 1.4 | 0 |