

Mustapha Oulad-Abdelghani

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6406122/publications.pdf>

Version: 2024-02-01

40
papers

4,291
citations

186265

28
h-index

302126

39
g-index

40
all docs

40
docs citations

40
times ranked

7046
citing authors

#	ARTICLE	IF	CITATIONS
1	H3K9 and H3K14 acetylation co-occur at many gene regulatory elements, while H3K14ac marks a subset of inactive inducible promoters in mouse embryonic stem cells. <i>BMC Genomics</i> , 2012, 13, 424.	2.8	409
2	Heterochromatin Formation in Mammalian Cells. <i>Molecular Cell</i> , 2001, 7, 729-739.	9.7	353
3	Loss of C9 <scp>ORF</scp> 72 impairs autophagy and synergizes with polyQ Ataxinâ€² to induce motor neuron dysfunction and cell death. <i>EMBO Journal</i> , 2016, 35, 1276-1297.	7.8	343
4	Nodal Antagonists in the Anterior Visceral Endoderm Prevent the Formation of Multiple Primitive Streaks. <i>Developmental Cell</i> , 2002, 3, 745-756.	7.0	330
5	Interaction with members of the heterochromatin protein 1 (HP1) family and histone deacetylation are differentially involved in transcriptional silencing by members of the TIF1 family. <i>EMBO Journal</i> , 1999, 18, 6385-6395.	7.8	325
6	Retinoic Acid Metabolism and Signaling Pathways in the Adult and Developing Mouse Testis. <i>Endocrinology</i> , 2006, 147, 96-110.	2.8	225
7	Translation of Expanded CGG Repeats into FMRpolyG Is Pathogenic and May Contribute to Fragile X Tremor Ataxia Syndrome. <i>Neuron</i> , 2017, 93, 331-347.	8.1	194
8	STRA8-deficient spermatocytes initiate, but fail to complete, meiosis and undergo premature chromosome condensation. <i>Journal of Cell Science</i> , 2008, 121, 3233-3242.	2.0	189
9	AP-2.2, a novel gene related to AP-2, is expressed in the forebrain, limbs and face during mouse embryogenesis. <i>Mechanisms of Development</i> , 1996, 54, 83-94.	1.7	175
10	Efficient Cloning of cDNAs of Retinoic Acid-Responsive Genes in P19 Embryonal Carcinoma Cells and Characterization of a Novel Mouse Gene, Stra1 (Mouse LERK-2/Eplg2). <i>Developmental Biology</i> , 1995, 170, 420-433.	2.0	168
11	The Major Yolk Protein Vitellogenin Interferes with the Anti-Plasmodium Response in the Malaria Mosquito <i>Anopheles gambiae</i> . <i>PLoS Biology</i> , 2010, 8, e1000434.	5.6	144
12	Murine spermatogonial stem cells: targeted transgene expression and purification in an active state. <i>EMBO Reports</i> , 2002, 3, 753-759.	4.5	121
13	AP-2.2: A Novel AP-2-Related Transcription Factor Induced by Retinoic Acid during Differentiation of P19 Embryonal Carcinoma Cells. <i>Experimental Cell Research</i> , 1996, 225, 338-347.	2.6	106
14	Sequence and expression pattern of the Stra7 (Gbx-2) homeobox-containing gene induced by retinoic acid in P19 embryonal carcinoma cells. <i>Developmental Dynamics</i> , 1995, 204, 372-382.	1.8	100
15	A new mouse member of the Wnt gene family, mWnt-8, is expressed during early embryogenesis and is ectopically induced by retinoic acid. <i>Mechanisms of Development</i> , 1996, 58, 141-152.	1.7	92
16	Meis2, a novel mouse Pbx-related homeobox gene induced by retinoic acid during differentiation of P19 embryonal carcinoma cells. <i>Developmental Dynamics</i> , 1997, 210, 173-183.	1.8	88
17	Pathogenic and Non-pathogenic Polyglutamine Tracts Have Similar Structural Properties: Towards a Length-dependent Toxicity Gradient. <i>Journal of Molecular Biology</i> , 2007, 371, 235-244.	4.2	86
18	Molecular Characterization of the Microsomal Tamoxifen Binding Site. <i>Journal of Biological Chemistry</i> , 2004, 279, 34048-34061.	3.4	84

#	ARTICLE	IF	CITATIONS
19	Poly(L-lysine) nanostructured particles for gene delivery and hormone stimulation. <i>Biomaterials</i> , 2010, 31, 1699-1706.	11.4	83
20	FMRpolyG-positive inclusions in CNS and non-CNS organs of a fragile X premutation carrier with fragile X-associated tremor/ataxia syndrome. <i>Acta Neuropathologica Communications</i> , 2014, 2, 162.	5.2	78
21	TIF1 \hat{I} , a Novel HP1-interacting Member of the Transcriptional Intermediary Factor 1 (TIF1) Family Expressed by Elongating Spermatids. <i>Journal of Biological Chemistry</i> , 2004, 279, 48329-48341.	3.4	67
22	Suppression of cervical carcinoma cell growth by intracytoplasmic codelivery of anti-oncoprotein E6 antibody and small interfering RNA. <i>Molecular Cancer Therapeutics</i> , 2007, 6, 1728-1735.	4.1	60
23	Helios Deficiency Has Minimal Impact on T Cell Development and Function. <i>Journal of Immunology</i> , 2009, 183, 2303-2311.	0.8	58
24	Uniform Widespread Nuclear Phosphorylation of Histone H2AX Is an Indicator of Lethal DNA Replication Stress. <i>Cancers</i> , 2019, 11, 355.	3.7	58
25	SiRNA-loaded multi-shell nanoparticles incorporated into a multilayered film as a reservoir for gene silencing. <i>Biomaterials</i> , 2010, 31, 6013-6018.	11.4	53
26	Down-regulation of the Phosphatidylinositol 3-Kinase/Akt Pathway Is Involved in Retinoic Acid-induced Phosphorylation, Degradation, and Transcriptional Activity of Retinoic Acid Receptor \hat{I}^2 . <i>Journal of Biological Chemistry</i> , 2002, 277, 24859-24862.	3.4	50
27	Targeting endogenous nuclear antigens by electrotransfer of monoclonal antibodies in living cells. <i>MAbs</i> , 2013, 5, 518-522.	5.2	46
28	Differential expression of retinoic acid-inducible (Stra) genes during mouse placentation. <i>Mechanisms of Development</i> , 2000, 92, 295-299.	1.7	42
29	Imaging of native transcription factors and histone phosphorylation at high resolution in live cells. <i>Journal of Cell Biology</i> , 2018, 217, 1537-1552.	5.2	35
30	Histone H3 Tails Containing Dimethylated Lysine and Adjacent Phosphorylated Serine Modifications Adopt a Specific Conformation during Mitosis and Meiosis. <i>Molecular and Cellular Biology</i> , 2008, 28, 1739-1754.	2.3	23
31	Targeting the replisome with transduced monoclonal antibodies triggers lethal DNA replication stress in cancer cells. <i>Experimental Cell Research</i> , 2016, 342, 145-158.	2.6	20
32	Restricted expression of a novel retinoic acid responsive gene during limb bud dorsoventral patterning and endochondral ossification. , 1996, 19, 66-73.		18
33	p38 \hat{I}^2 -Mediated Phosphorylation and Sumoylation of ATF7 Are Mutually Exclusive. <i>Journal of Molecular Biology</i> , 2008, 384, 980-991.	4.2	14
34	A Novel Nanobody Precisely Visualizes Phosphorylated Histone H2AX in Living Cancer Cells under Drug-Induced Replication Stress. <i>Cancers</i> , 2021, 13, 3317.	3.7	14
35	Single-chain Fv fragment antibodies selected from an intrabody library as effective mono- or bivalent reagents for in vitro protein detection. <i>Journal of Immunological Methods</i> , 2011, 369, 42-50.	1.4	10
36	Selective neutralization of APP-C99 with monoclonal antibodies reduces the production of Alzheimer's A \hat{I}^2 peptides. <i>Neurobiology of Aging</i> , 2012, 33, 2704-2714.	3.1	8

#	ARTICLE	IF	CITATIONS
37	Molecular Architecture of the Basal Transcription Factor B-TFIID. <i>Journal of Biological Chemistry</i> , 2004, 279, 21802-21807.	3.4	7
38	In vitro hydroquinone-induced instauration of histone bivalent mark on human retroelements (LINE-1) in HL60 cells. <i>Toxicology in Vitro</i> , 2017, 40, 1-10.	2.4	6
39	ATF7 is stabilized during mitosis in a CDK1-dependent manner and contributes to cyclin D1 expression. <i>Cell Cycle</i> , 2015, 14, 2655-2666.	2.6	5
40	Generation of Monoclonal Antibody Fragments Binding the Native β -Secretase Complex for Use in Structural Studies. <i>Biochemistry</i> , 2012, 51, 8779-8790.	2.5	4