

Fang Liu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/6413100/publications.pdf>

Version: 2024-02-01

30
papers

953
citations

567281

15
h-index

477307

29
g-index

31
all docs

31
docs citations

31
times ranked

1289
citing authors

#	ARTICLE	IF	CITATIONS
1	Melatonin-stimulated MSC-derived exosomes improve diabetic wound healing through regulating macrophage M1 and M2 polarization by targeting the PTEN/AKT pathway. <i>Stem Cell Research and Therapy</i> , 2020, 11, 259.	5.5	221
2	Exosomes derived from atorvastatin-pretreated MSC accelerate diabetic wound repair by enhancing angiogenesis via AKT/eNOS pathway. <i>Stem Cell Research and Therapy</i> , 2020, 11, 350.	5.5	173
3	SLC47A1 gene rs2289669 G>A variants enhance the glucose-lowering effect of metformin via delaying its excretion in Chinese type 2 diabetes patients. <i>Diabetes Research and Clinical Practice</i> , 2015, 109, 57-63.	2.8	48
4	Liraglutide-loaded PLGA/gelatin electrospun nanofibrous mats promote angiogenesis to accelerate diabetic wound healing via the modulation of miR-29b-3p. <i>Biomaterials Science</i> , 2020, 8, 4225-4238.	5.4	44
5	Development of Multimarker Diagnostic Models from Metabolomics Analysis for Gestational Diabetes Mellitus (GDM). <i>Molecular and Cellular Proteomics</i> , 2018, 17, 431-441.	3.8	43
6	Vitamin D deficiency increases the risk of peripheral neuropathy in Chinese patients with type 2 diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2017, 33, e2820.	4.0	41
7	Development of a novel RNAi therapy: Engineered miR-31 exosomes promoted the healing of diabetic wounds. <i>Bioactive Materials</i> , 2021, 6, 2841-2853.	15.6	40
8	SLC22A2 gene 808 G/T variant is related to plasma lactate concentration in Chinese type 2 diabetics treated with metformin. <i>Acta Pharmacologica Sinica</i> , 2010, 31, 184-190.	6.1	35
9	Heterogeneity of insulin resistance and beta cell dysfunction in gestational diabetes mellitus: a prospective cohort study of perinatal outcomes. <i>Journal of Translational Medicine</i> , 2018, 16, 289.	4.4	34
10	Association between serum cystatin C and diabetic peripheral neuropathy: a cross-sectional study of a Chinese type 2 diabetic population. <i>European Journal of Endocrinology</i> , 2014, 171, 641-648.	3.7	31
11	The γ 1123G>C Variant of PTPN22 Gene Promoter is Associated with Latent Autoimmune Diabetes in Adult Chinese Hans. <i>Cell Biochemistry and Biophysics</i> , 2012, 62, 273-279.	1.8	30
12	A high thyroid stimulating hormone level is associated with diabetic peripheral neuropathy in type 2 diabetes patients. <i>Diabetes Research and Clinical Practice</i> , 2016, 115, 122-129.	2.8	27
13	Polymorphism of Organic Cation Transporter 2 Improves Glucose-Lowering Effect of Metformin via Influencing Its Pharmacokinetics in Chinese Type 2 Diabetic Patients. <i>Molecular Diagnosis and Therapy</i> , 2015, 19, 25-33.	3.8	26
14	High cystatin C levels predict severe retinopathy in type 2 diabetes patients. <i>European Journal of Epidemiology</i> , 2013, 28, 775-778.	5.7	25
15	TRAF6 mediates high glucose-induced endothelial dysfunction. <i>Experimental Cell Research</i> , 2018, 370, 490-497.	2.6	20
16	Elevated First-Trimester Total Bile Acid is Associated with the Risk of Subsequent Gestational Diabetes. <i>Scientific Reports</i> , 2016, 6, 34070.	3.3	16
17	Low levels of ficolin-3 are associated with diabetic peripheral neuropathy. <i>Acta Diabetologica</i> , 2016, 53, 295-302.	2.5	16
18	Different effect of testosterone and oestrogen on urinary excretion of metformin via regulating OCTs and MATEs expression in the kidney of mice. <i>Journal of Cellular and Molecular Medicine</i> , 2016, 20, 2309-2317.	3.6	14

#	ARTICLE	IF	CITATIONS
19	Vibrating Perception Threshold and Body Mass Index Are Associated with Abnormal Foot Plantar Pressure in Type 2 Diabetes Outpatients. <i>Diabetes Technology and Therapeutics</i> , 2012, 14, 1053-1059.	4.4	10
20	Differentially methylated circulating DNA: A novel biomarker to monitor beta cell death. <i>Journal of Diabetes and Its Complications</i> , 2018, 32, 349-353.	2.3	10
21	The Gonadal Hormone Regulates the Plasma Lactate Levels in Type 2 Diabetes Treated With and Without Metformin. <i>Diabetes Technology and Therapeutics</i> , 2012, 14, 469-474.	4.4	9
22	Relationship between High Serum Cystatin C Levels and the Risk of Gestational Diabetes Mellitus. <i>PLoS ONE</i> , 2016, 11, e0147277.	2.5	9
23	Early elevated alkaline phosphatase increases the risk of large-for-gestational-age birth weight in pregnant women with normal glucose tolerance. <i>Diabetes Research and Clinical Practice</i> , 2018, 141, 209-216.	2.8	7
24	Serum IL-17A concentration and a IL17RA single nucleotide polymorphism contribute to the risk of autoimmune type 1 diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2022, 38, e3547.	4.0	6
25	Vibration perception threshold for sight-threatening retinopathy screening in type 2 diabetic outpatients. <i>Diabetes/Metabolism Research and Reviews</i> , 2013, 29, 525-531.	4.0	5
26	The value of ankle-brachial index screening for cardiovascular disease in type 2 diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2019, 35, e3076.	4.0	5
27	The role of complement factor H in gestational diabetes mellitus and pregnancy. <i>BMC Pregnancy and Childbirth</i> , 2021, 21, 562.	2.4	4
28	Index of Plantar Pressure Alters with Prolonged Diabetes Duration. <i>Diabetes Therapy</i> , 2019, 10, 2139-2152.	2.5	3
29	High level of complement factor Ba within first prenatal test of gestation increases the risk of subsequent gestational diabetes: a propensity score-matched study. <i>Gynecological Endocrinology</i> , 2022, 38, 158-163.	1.7	1
30	Adult multisystem Langerhans cell histiocytosis involving parathyroid glands and pancreas. <i>Chinese Medical Journal</i> , 2014, 127, 1597.	2.3	0