

Jong Bong Lee

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/6417051/jong-bong-lee-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

46
papers

587
citations

14
h-index

22
g-index

51
ext. papers

771
ext. citations

5.4
avg, IF

3.68
L-index

#	Paper	IF	Citations
46	Development and optimisation of simulated salivary fluid for biorelevant oral cavity dissolution. <i>European Journal of Pharmaceutics and Biopharmaceutics</i> , 2021 , 160, 125-133	5.7	3
45	Natural sesame oil is superior to pre-digested lipid formulations and purified triglycerides in promoting the intestinal lymphatic transport and systemic bioavailability of cannabidiol. <i>European Journal of Pharmaceutics and Biopharmaceutics</i> , 2021 , 162, 43-49	5.7	6
44	Designing topographically textured microparticles for induction and modulation of osteogenesis in mesenchymal stem cell engineering. <i>Biomaterials</i> , 2021 , 266, 120450	15.6	12
43	Targeted delivery of lopinavir to HIV reservoirs in the mesenteric lymphatic system by lipophilic ester prodrug approach. <i>Journal of Controlled Release</i> , 2021 , 329, 1077-1089	11.7	9
42	Inclusion of Medium-Chain Triglyceride in Lipid-Based Formulation of Cannabidiol Facilitates Micellar Solubilization In Vitro, but In Vivo Performance Remains Superior with Pure Sesame Oil Vehicle. <i>Pharmaceutics</i> , 2021 , 13,	6.4	3
41	Administration in fed state but not controlled release in the colon increases oral bioavailability of DF030263, a promising drug candidate for chronic lymphocytic leukemia. <i>European Journal of Pharmaceutics and Biopharmaceutics</i> , 2021 , 165, 106-112	5.7	
40	Development and validation of a cost-effective and sensitive bioanalytical HPLC-UV method for determination of lopinavir in rat and human plasma. <i>Biomedical Chromatography</i> , 2020 , 34, e4934	1.7	3
39	Development of a Population Pharmacokinetics-Based in vitro-in vivo Correlation Model for Drugs with Site-Dependent Absorption: the Acyclovir Case Study. <i>AAPS Journal</i> , 2020 , 22, 27	3.7	7
38	Interspecies prediction of pharmacokinetics and tissue distribution of doxorubicin by physiologically-based pharmacokinetic modeling. <i>Biopharmaceutics and Drug Disposition</i> , 2020 , 41, 192-205	1.7	1
37	Intradermal and transdermal drug delivery using microneedles - Fabrication, performance evaluation and application to lymphatic delivery. <i>Advanced Drug Delivery Reviews</i> , 2020 , 153, 195-215	18.5	46
36	Is rat a good model for assessment of particulate-based taste-masked formulations?. <i>European Journal of Pharmaceutics and Biopharmaceutics</i> , 2020 , 146, 1-9	5.7	2
35	Pharmacokinetic Modeling of the Impact of P-glycoprotein on Ondansetron Disposition in the Central Nervous System. <i>Pharmaceutical Research</i> , 2020 , 37, 205	4.5	
34	Predicting Intestinal and Hepatic First-Pass Metabolism of Orally Administered Testosterone Undecanoate. <i>Applied Sciences (Switzerland)</i> , 2020 , 10, 7283	2.6	1
33	Codrug Approach for the Potential Treatment of EML4-ALK Positive Lung Cancer. <i>ACS Medicinal Chemistry Letters</i> , 2020 , 11, 316-321	4.3	0
32	Physiologically Relevant In Vitro-In Vivo Correlation (IVIVC) Approach for Sildenafil with Site-Dependent Dissolution. <i>Pharmaceutics</i> , 2019 , 11,	6.4	11
31	Cardiac glycoside cerberin exerts anticancer activity through PI3K/AKT/mTOR signal transduction inhibition. <i>Cancer Letters</i> , 2019 , 453, 57-73	9.9	25
30	Prevention of paclitaxel-induced neuropathy by formulation approach. <i>Journal of Controlled Release</i> , 2019 , 303, 109-116	11.7	17

29	Simultaneous quantification of ondansetron and tariquidar in rat and human plasma using a high performance liquid chromatography-ultraviolet method. <i>Biomedical Chromatography</i> , 2019 , 33, e4653	1.7	3
28	Application of Size and Maturation Functions to Population Pharmacokinetic Modeling of Pediatric Patients. <i>Pharmaceutics</i> , 2019 , 11,	6.4	7
27	A novel nucleoside rescue metabolic pathway may be responsible for therapeutic effect of orally administered cordycepin. <i>Scientific Reports</i> , 2019 , 9, 15760	4.9	9
26	Solid lipid nanoparticles self-assembled from spray dried microparticles. <i>International Journal of Pharmaceutics</i> , 2019 , 572, 118784	6.5	3
25	Reduced variability in tacrolimus pharmacokinetics following intramuscular injection compared to oral administration in cynomolgus monkeys: Investigating optimal dosing regimens. <i>Journal of Pharmacological Sciences</i> , 2019 , 139, 65-71	3.7	1
24	Quantitative Prediction of Oral Bioavailability of a Lipophilic Antineoplastic Drug Bexarotene Administered in Lipidic Formulation Using a Combined In Vitro Lipolysis/Microsomal Metabolism Approach. <i>Journal of Pharmaceutical Sciences</i> , 2019 , 108, 1047-1052	3.9	5
23	Self-Assembling Benzothiazole-Based Gelators: A Mechanistic Understanding of in Vitro Bioactivation and Gelation. <i>Molecular Pharmaceutics</i> , 2018 , 15, 1578-1586	5.6	2
22	Lipophilic activated ester prodrug approach for drug delivery to the intestinal lymphatic system. <i>Journal of Controlled Release</i> , 2018 , 286, 10-19	11.7	26
21	Quantitative analysis of lab-to-lab variability in Caco-2 permeability assays. <i>European Journal of Pharmaceutics and Biopharmaceutics</i> , 2017 , 114, 38-42	5.7	34
20	Hyperlipidaemia alone and in combination with acidosis can increase the incidence and severity of statin-induced myotoxicity. <i>European Journal of Pharmaceutical Sciences</i> , 2017 , 100, 163-175	5.1	6
19	Development of Cordycepin Formulations for Preclinical and Clinical Studies. <i>AAPS PharmSciTech</i> , 2017 , 18, 3219-3226	3.9	12
18	Oral administration of cannabis with lipids leads to high levels of cannabinoids in the intestinal lymphatic system and prominent immunomodulation. <i>Scientific Reports</i> , 2017 , 7, 14542	4.9	53
17	Simple and sensitive HPLC-UV method for determination of bexarotene in rat plasma. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2017 , 1040, 73-80	3.2	11
16	The role of acid-base imbalance in statin-induced myotoxicity. <i>Translational Research</i> , 2016 , 174, 140-160, e14	11.4	22
15	In vitro anticancer properties and biological evaluation of novel natural alkaloid jerantinine B. <i>Cancer Letters</i> , 2016 , 370, 185-97	9.9	35
14	Dietary fats and pharmaceutical lipid excipients increase systemic exposure to orally administered cannabis and cannabis-based medicines. <i>American Journal of Translational Research (discontinued)</i> , 2016 , 8, 3448-59	3	41
13	Sensitive liquid chromatography-tandem mass spectrometry method for the simultaneous determination of benzyl butyl phthalate and its metabolites, monobenzyl phthalate and monobutyl phthalate, in rat plasma, urine, and various tissues collected from a toxicokinetic study. <i>Analytical and Bioanalytical Chemistry</i> , 2015 , 407, 7391-400	4.4	6
12	A sensitive LC-ESI-MS/MS method for the quantification of avobenzone in rat plasma and skin layers: Application to a topical administration study. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2015 , 1003, 41-6	3.2	5

11	Exposure assessment of epidermal growth factor to various tissues in mice after intravenous and subcutaneous administration. <i>Journal of Pharmacy and Pharmacology</i> , 2015 , 67, 1519-27	4.8	2
10	Simultaneous determination of phenoxyethanol and its major metabolite, phenoxyacetic acid, in rat biological matrices by LC-MS/MS with polarity switching: Application to ADME studies. <i>Talanta</i> , 2015 , 144, 29-38	6.2	9
9	A novel in vitro permeability assay using three-dimensional cell culture system. <i>Journal of Biotechnology</i> , 2015 , 205, 93-100	3.7	13
8	Population pharmacokinetics of a novel histone deacetylase inhibitor, cyclo{(2S)-2-amino-8-[(aminocarbonyl)hydrazono]decanoyl-1-L-tryptophyl-L-isoleucyl-(2R)-2-piperidinecarbonyl} (SD-2007), and its metabolic	3.2	1
7	Assessment of bisphenol A exposure in Korean pregnant women by physiologically based pharmacokinetic modeling. <i>Journal of Toxicology and Environmental Health - Part A: Current Issues</i> , 2010 , 73, 1586-98	3.2	18
6	Prediction of drug bioavailability in humans using immobilized artificial membrane phosphatidylcholine column chromatography and in vitro hepatic metabolic clearance. <i>Biomedical Chromatography</i> , 2009 , 23, 764-9	1.7	10
5	Determination of zearalenone by liquid chromatography/tandem mass spectrometry and application to a pharmacokinetic study. <i>Biomedical Chromatography</i> , 2009 , 23, 1014-21	1.7	25
4	Enhanced absorption and tissue distribution of paclitaxel following oral administration of DHP 107, a novel mucoadhesive lipid dosage form. <i>Cancer Chemotherapy and Pharmacology</i> , 2009 , 64, 87-94	3.5	15
3	Development of a Sensitive LC Assay with Fluorescence Detection for the Determination of Zearalenone in Rat Serum. <i>Chromatographia</i> , 2009 , 69, 295-299	2.1	9
2	Physiologically based pharmacokinetics of zearalenone. <i>Journal of Toxicology and Environmental Health - Part A: Current Issues</i> , 2009 , 72, 1395-405	3.2	27
1	Disposition, oral bioavailability, and tissue distribution of zearalenone in rats at various dose levels. <i>Journal of Toxicology and Environmental Health - Part A: Current Issues</i> , 2009 , 72, 1406-11	3.2	31