

Mustafa AetÄ°n

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/642253/publications.pdf>

Version: 2024-02-01

157
papers

1,577
citations

361045

20
h-index

476904

29
g-index

159
all docs

159
docs citations

159
times ranked

2481
citing authors

#	ARTICLE	IF	CITATIONS
1	Prediction of Coronary Artery Disease Severity Using CHADS2 and CHA2DS2-VASc Scores and a Newly Defined CHA2DS2-VASc-HS Score. American Journal of Cardiology, 2014, 113, 950-956.	0.7	74
2	Increased Platelet Distribution Width Is Associated With ST-Segment Elevation Myocardial Infarction and Thrombolysis Failure. Angiology, 2014, 65, 737-743.	0.8	50
3	Predictive value of elevated D-dimer in patients undergoing primary angioplasty for ST elevation myocardial infarction. Blood Coagulation and Fibrinolysis, 2013, 24, 704-710.	0.5	44
4	Relation of Epicardial Fat Thickness with Carotid Intima-Media Thickness in Patients with Type 2 Diabetes Mellitus. International Journal of Endocrinology, 2013, 2013, 1-6.	0.6	43
5	The degree of premature hair graying as an independent risk marker for coronary artery disease: a predictor of biological age rather than chronological age. Anatolian Journal of Cardiology, 2012, 12, 457-63.	0.4	35
6	Level of Selvester QRS score is predictive of ST-segment resolution and 30-day outcomes in patients with acute myocardial infarction undergoing primary coronary intervention. American Heart Journal, 2006, 151, 1239.e1-1239.e7.	1.2	34
7	Non-dipping pattern in untreated hypertensive patients is related to increased pulse wave velocity independent of raised nocturnal blood pressure. Blood Pressure, 2013, 22, 34-38.	0.7	32
8	Effect of epicardial adipose tissue on diastolic functions and left atrial dimension in untreated hypertensive patients with normal systolic function. Journal of Cardiology, 2013, 61, 359-364.	0.8	31
9	Effect of a successful percutaneous coronary intervention for chronic total occlusion on parameters of ventricular repolarization. Coronary Artery Disease, 2014, 25, 705-712.	0.3	31
10	CHA2DS2-VASc-HS score in non-ST elevation acute coronary syndrome patients: assessment of coronary artery disease severity and complexity and comparison to other scoring systems in the prediction of in-hospital major adverse cardiovascular events. Anatolian Journal of Cardiology, 2016, 16, 742-748.	0.5	31
11	TcR β 2-depleted haploidentical transplantation results in adult acute leukemia patients. Hematology, 2017, 22, 136-144.	0.7	31
12	Overexpression of cyclooxygenase-2 in multiple myeloma: Association with reduced survival. American Journal of Hematology, 2005, 80, 169-173.	2.0	28
13	Increased pulse wave velocity in patients with panic disorder: Independent vascular influence of panic disorder on arterial stiffness. Journal of Psychosomatic Research, 2012, 73, 145-148.	1.2	28
14	Efficacy and safety of oral deferasirox treatment in the posttransplant period for patients who have undergone allogeneic hematopoietic stem cell transplantation (alloHSCT). Annals of Hematology, 2012, 91, 743-749.	0.8	27
15	Comparison of low dose versus standard dose heparin for radial approach in elective coronary angiography?. International Journal of Cardiology, 2015, 187, 389-392.	0.8	27
16	The independent relationship of epicardial adipose tissue with carotid intima-media thickness and endothelial functions. Blood Pressure Monitoring, 2013, 18, 85-93.	0.4	26
17	New Markers for Ventricular Repolarization in Coronary Slow Flow: Tp α e Interval, Tp α e/QT Ratio, and Tp α e/QTc Ratio. Annals of Noninvasive Electrocardiology, 2015, 20, 338-344.	0.5	26
18	Increased Serum Bilirubin Level is Related to Good Collateral Development in Patients with Chronic Total Coronary Occlusion. Internal Medicine, 2012, 51, 249-255.	0.3	24

#	ARTICLE	IF	CITATIONS
19	The importance of fragmented QRS complexes in prediction of myocardial infarction and reperfusion parameters in patients undergoing primary percutaneous coronary intervention. <i>Turk Kardiyoloji Dernegi Arsivi</i> , 2012, 40, 213-222.	0.6	24
20	Effect of Smoking on Tp-e Interval, Tp-e/QT and Tp-e/QTc Ratios as Indices of Ventricular Arrhythmogenesis. <i>Heart Lung and Circulation</i> , 2014, 23, 827-832.	0.2	23
21	<i>Saprochaete capitata</i> as an emerging fungus among patients with haematological malignencies. <i>Mycoses</i> , 2015, 58, 491-497.	1.8	23
22	Negative Effects of Acute Sleep Deprivation on Left Ventricular Functions and Cardiac Repolarization in Healthy Young Adults. <i>PACE - Pacing and Clinical Electrophysiology</i> , 2015, 38, 713-722.	0.5	22
23	Predictive value of elevated soluble CD40 ligand in patients undergoing primary angioplasty for ST-segment elevation myocardial infarction. <i>Coronary Artery Disease</i> , 2014, 25, 558-564.	0.3	20
24	Low-Level Vitamin D Is Associated with Atrial Fibrillation in Patients with Chronic Heart Failure. <i>Advances in Clinical and Experimental Medicine</i> , 2016, 25, 51-57.	0.6	20
25	Predictive value of fibrinogen-to-albumin ratio in acute coronary syndrome. <i>Herz</i> , 2020, 45, 145-151.	0.4	19
26	COVID 19 disease independently predicted endothelial dysfunction measured by flow-mediated dilatation. <i>International Journal of Cardiovascular Imaging</i> , 2022, 38, 25-32.	0.7	19
27	Premature Hair Whitening is an Independent Predictor of Carotid Intima-media Thickness in Young and Middle-aged Men. <i>Internal Medicine</i> , 2013, 52, 29-36.	0.3	18
28	Prognostic value of elevated mean platelet volume in patients undergoing primary angioplasty for ST-elevation myocardial infarction. <i>Acta Cardiologica</i> , 2013, 68, 307-314.	0.3	18
29	Fragmented QRS on Admission Electrocardiography Predicts Long-Term Mortality in Patients with Non-ST-Segment Elevation Myocardial Infarction. <i>Annals of Noninvasive Electrocardiology</i> , 2016, 21, 352-357.	0.5	18
30	Metabolic syndrome is a predictor for an ECG sign of no-reflow after primary PCI in patients with acute ST-elevation myocardial infarction. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2008, 18, 441-447.	1.1	17
31	An increase in epicardial adipose tissue is strongly associated with carotid-intima media thickness and atherosclerotic plaque, but LDL only with the plaque. <i>Anatolian Journal of Cardiology</i> , 2017, 17, 56-63.	0.5	17
32	Relationship between fragmented QRS complexes and left ventricular systolic and diastolic functions. <i>Herz</i> , 2013, 38, 665-670.	0.4	16
33	Relation of Angiographic Thrombus Burden with Severity of Coronary Artery Disease in Patients with ST Segment Elevation Myocardial Infarction. <i>Medical Science Monitor</i> , 2015, 21, 3540-3546.	0.5	16
34	Androgenetic alopecia is associated with increased arterial stiffness in asymptomatic young adults. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2015, 29, 26-30.	1.3	16
35	The Independent Relationship of Systemic Inflammation With Fragmented QRS Complexes in Patients With Acute Coronary Syndromes. <i>Korean Circulation Journal</i> , 2012, 42, 449.	0.7	15
36	Elevated serum YKL-40 level predicts myocardial reperfusion and in-hospital MACE in patients with STEMI. <i>Herz</i> , 2013, 38, 202-209.	0.4	15

#	ARTICLE	IF	CITATIONS
37	Performance of Galactomannan Antigen, Beta-d-Glucan, and Aspergillus-Lateral-Flow Device for the Diagnosis of Invasive Aspergillosis. <i>Indian Journal of Hematology and Blood Transfusion</i> , 2017, 33, 87-92.	0.3	15
38	Fragmented QRS may predict postoperative atrial fibrillation in patients undergoing isolated coronary artery bypass graft surgery. <i>Anatolian Journal of Cardiology</i> , 2012, 12, 576-83.	0.4	14
39	Absence and Resolution of Fragmented QRS Predict Reversible Myocardial Ischemia With Higher Probability of ST Segment Resolution in Patients With ST Segment Elevation Myocardial Infarction. <i>Korean Circulation Journal</i> , 2012, 42, 674.	0.7	14
40	Increased serum resistin levels in patients with coronary slow-flow phenomenon. <i>Herz</i> , 2013, 38, 773-778.	0.4	14
41	ABO and Rh Blood Group Distribution in Kayseri Province in Turkey. <i>Turkish Journal of Haematology</i> , 2012, 29, 102-103.	0.2	13
42	Increased YKL-40 levels in patients with isolated coronary artery ectasia: an observational study. <i>Anatolian Journal of Cardiology</i> , 2013, 13, 465-70.	0.4	13
43	Ankle Brachial Index Intensifies the Diagnostic Accuracy of Epicardial Fat Thickness for the Prediction of Coronary Artery Disease Complexity. <i>Heart Lung and Circulation</i> , 2014, 23, 764-771.	0.2	13
44	Complexity of lower extremity peripheral artery disease reflects the complexity of coronary artery disease. <i>Vascular</i> , 2015, 23, 366-373.	0.4	13
45	A Case of Carbon Monoxide Poisoning Presenting with Supraventricular Tachycardia. <i>Internal Medicine</i> , 2011, 50, 2607-2609.	0.3	12
46	The particular interactions of the traditional cardiovascular risk factors with different circulating specific leukocyte subtype counts in blood: an observational study. <i>Anatolian Journal of Cardiology</i> , 2011, 11, 573-81.	0.4	12
47	Independent determinants of ascending aortic dilatation in hypertensive patients. <i>Blood Pressure Monitoring</i> , 2012, 17, 223-230.	0.4	12
48	Increased Epicardial Adipose Tissue in Patients with Isolated Coronary Artery Ectasia. <i>Internal Medicine</i> , 2012, 51, 833-838.	0.3	12
49	Relationship of fragmented QRS with prognostic markers and in-hospital MACE in patients undergoing CABG. <i>Scandinavian Cardiovascular Journal</i> , 2012, 46, 107-113.	0.4	12
50	Predictive value of elevated cystatin C in patients undergoing primary angioplasty for ST-elevation myocardial infarction. <i>Journal of Critical Care</i> , 2013, 28, 882.e13-882.e20.	1.0	12
51	Comparison of double dose plateletpheresis on the Fenwal Amicus, Fresenius COM.TEC and Trima Accel cell separators. <i>Transfusion and Apheresis Science</i> , 2014, 51, 193-196.	0.5	12
52	Comparison of plateletpheresis on the Fenwal Amicus, Fresenius COM.TEC, and Trima Accel Cell separators. <i>Journal of Clinical Apheresis</i> , 2015, 30, 171-175.	0.7	12
53	Ambulatory arterial stiffness index is associated with impaired left atrial mechanical functions in hypertensive diabetic patients: A speckle tracking study. <i>Anatolian Journal of Cardiology</i> , 2015, 15, 807-813.	0.5	12
54	Assessment of the association between the presence of fragmented QRS and the predicted risk score of sudden cardiac death at 5 years in patients with hypertrophic cardiomyopathy. <i>Anatolian Journal of Cardiology</i> , 2017, 18, 54-61.	0.5	12

#	ARTICLE	IF	CITATIONS
55	Increased Epicardial Adipose Tissue Thickness Is Correlated with Ascending Aortic Diameter. <i>Tohoku Journal of Experimental Medicine</i> , 2012, 226, 183-190.	0.5	11
56	Elevated serum osteoprotegerin levels predict in-hospital major adverse cardiac events in patients with ST elevation myocardial infarction. <i>Journal of Cardiology</i> , 2012, 60, 355-360.	0.8	11
57	Increased plasminogen activator inhibitor-1 levels in patients with isolated coronary artery ectasia. <i>Journal of Thrombosis and Thrombolysis</i> , 2012, 33, 120-123.	1.0	11
58	Aortic valve sclerosis is a high predictive marker of systemic endothelial dysfunction in hypertensive patients. <i>Herz</i> , 2013, 38, 915-921.	0.4	11
59	Increased Bone Marrow Iron Scores Are Strongly Correlated With Elevated Serum Ferritin Levels and Poorer Survival in Patients With Iron Overload That Underwent Allogeneic Hematopoietic Stem Cell Transplantation: A Single Center Experience. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2016, 16, 582-587.	0.2	11
60	The association of serum albumin with coronary slow flow. <i>Wiener Klinische Wochenschrift</i> , 2014, 126, 468-473.	1.0	10
61	Is atrial fibrillation a risk factor for contrast-induced nephropathy in patients with ST-elevation myocardial infarction?. <i>Journal of Cardiology</i> , 2016, 67, 327-330.	0.8	10
62	Endothelial dysfunction, subclinical atherosclerosis and LDL cholesterol are the independent predictors of left atrial functions in hypertension. <i>International Journal of Cardiovascular Imaging</i> , 2020, 36, 69-77.	0.7	10
63	The Relationship between Blood Viscosity and Isolated Coronary Artery Ectasia. <i>Acta Cardiologica Sinica</i> , 2019, 35, 20-26.	0.1	10
64	Increased circulating soluble CD40 levels in patients with slow coronary flow phenomenon: an observational study. <i>Anatolian Journal of Cardiology</i> , 2012, 13, 39-44.	0.4	9
65	Outcomes for haematological cancer patients admitted to an intensive care unit in a university hospital. <i>Australian Critical Care</i> , 2018, 31, 363-368.	0.6	9
66	A case of systemic lupus erythematosus presenting with hypereosinophilia and Loeffler endocarditis. <i>Acta Cardiologica</i> , 2010, 65, 571-3.	0.3	9
67	Preclinical Studies on Convalescent Human Immune Plasma-Derived Exosome: Omics and Antiviral Properties to SARS-CoV-2. <i>Frontiers in Immunology</i> , 2022, 13, 824378.	2.2	9
68	Relationship of fragmented QRS complexes with inadequate coronary collaterals in patients with chronic total occlusion. <i>Journal of Cardiovascular Medicine</i> , 2012, 13, 499-504.	0.6	8
69	Encephalitozoon intestinalis: A Rare Cause of Diarrhea in an Allogeneic Hematopoietic Stem Cell Transplantation (HSCT) Recipient Complicated by Albendazole-Related Hepatotoxicity. <i>Turkish Journal of Haematology</i> , 2013, 30, 204-208.	0.2	8
70	Management and outcomes of coronary artery perforations during percutaneous treatment of acute coronary syndromes. <i>Perfusion (United Kingdom)</i> , 2015, 30, 71-76.	0.5	8
71	Left ventricular hypertrophy is associated with increased sirtuin level in newly diagnosed hypertensive patients. <i>Clinical and Experimental Hypertension</i> , 2019, 41, 511-515.	0.5	8
72	Red Blood Cell Distribution Width (RDW) and its Association with Coronary Atherosclerotic Burden in Patients with Stable Angina Pectoris. <i>Electronic Journal of General Medicine</i> , 2012, 9, 7-13.	0.3	8

#	ARTICLE	IF	CITATIONS
73	Epicardial Adipose Tissue Predicts Carotid Intima-Media Thickness Independently of Body Mass Index and Waist Circumference. <i>Acta Cardiologica Sinica</i> , 2019, 35, 32-41.	0.1	8
74	The Effect of Mesenchymal Stromal Cells on the Mortality of Patients with Sepsis and Septic Shock: A Promising Therapy. <i>Emergency Medicine International</i> , 2022, 2022, 1-9.	0.3	8
75	Increased Epicardial Adipose Tissue Thickness on Transthoracic Echocardiography in Patients With BehÄŞet Disease. <i>Journal of Ultrasound in Medicine</i> , 2014, 33, 1393-1400.	0.8	7
76	Clinical staging in chronic heart failure associated with low vitamin D and elevated parathormone levels. <i>Acta Cardiologica</i> , 2014, 69, 665-671.	0.3	7
77	Is dipyridamole useful in improving left ventricular systolic and diastolic function in patients with coronary slow flow?. <i>Echocardiography</i> , 2016, 33, 1472-1478.	0.3	7
78	Acute Change in Left Ventricle End-diastolic Pressure after Primary Percutaneous Coronary Intervention in Patients with ST Segment Elevation Myocardial Infarction. <i>The American Heart Hospital Journal</i> , 2010, 8, 86.	0.2	7
79	Epicardial adipose tissue is independently associated with increased left ventricular mass in untreated hypertensive patients: an observational study. <i>Anatolian Journal of Cardiology</i> , 2013, 13, 320-7.	0.4	6
80	Increased pulse wave velocity and carotid intimaâ€™media thickness in patients with carpal tunnel syndrome. <i>Muscle and Nerve</i> , 2013, 47, 872-877.	1.0	6
81	Increased ventricular pacing threshold levels in patients with high serum uric acid levels. <i>Journal of Cardiology</i> , 2014, 64, 207-210.	0.8	6
82	Relationship between severity of pulmonary hypertension and coronary sinus diameter. <i>Revista Portuguesa De Cardiologia</i> , 2015, 34, 329-335.	0.2	6
83	Increased Hepatic Iron Content Predicts Poor Survival in Patients With Iron Overload Who Underwent Allogeneic Hematopoietic Stem Cell Transplantation. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2016, 16, S10-S18.	0.2	6
84	Epicardial adipose tissue predicted prolonged QTc interval in patients with arterial hypertension. <i>Clinical and Experimental Hypertension</i> , 2021, 43, 230-236.	0.5	6
85	Evaluation of left atrial mechanical function and atrial conduction abnormalities in Maras powder (smokeless tobacco) users and smokers: cardiovascular topic. <i>Cardiovascular Journal of Africa</i> , 2015, 26, 114-119.	0.2	6
86	Bone morphogenic protein-7 serum level decreases significantly in patients with contrast-induced nephropathy. <i>International Urology and Nephrology</i> , 2011, 43, 807-812.	0.6	5
87	Increased Carotid and Brachial Intimaâ€™Media Thickness is Related to Diffuse Coronary Involvement Rather Than Focal Lesions. <i>Angiology</i> , 2013, 64, 356-363.	0.8	5
88	Unusually located left ventricular outflow myxoma: a brief review of the literature. <i>BMJ Case Reports</i> , 2013, 2013, bcr2013009610-bcr2013009610.	0.2	5
89	Relationship of fragmented QRS with prognostic markers and long-term major adverse cardiac events in patients undergoing coronary artery bypass graft surgery. <i>Journal of Cardiovascular Medicine</i> , 2015, 16, 112-117.	0.6	5
90	The role of urotensin II and atherosclerotic risk factors in patients with slow coronary flow. <i>Interventional Medicine & Applied Science</i> , 2016, 8, 158-163.	0.2	5

#	ARTICLE	IF	CITATIONS
91	The Prognostic Significance of Elevated Serum Ferritin Levels Prior to Transplantation in Patients With Lymphoma Who Underwent Autologous Hematopoietic Stem Cell Transplantation (autoHSCT): Role of Iron Overload. <i>Clinical Lymphoma, Myeloma and Leukemia</i> , 2016, 16, S152-S158.	0.2	5
92	Epicardial adipose tissue is associated with increased systolic pulmonary artery pressure in patients with chronic obstructive pulmonary disease. <i>Clinical Respiratory Journal</i> , 2021, 15, 406-412.	0.6	5
93	Pericardial, pleural and peritoneal involvement in a patient with primary gastric mantle cell lymphoma. <i>Scottish Medical Journal</i> , 2015, 60, e21-e24.	0.7	4
94	Hematopoietic cell transplantation activity of Turkey in 2014: Ongoing increase in HCT rates. <i>Transfusion and Apheresis Science</i> , 2016, 54, 53-59.	0.5	4
95	The relationship between total atrial conduction time and left atrial global strain in patients with psoriasis vulgaris. <i>Archives of Medical Science</i> , 2019, 15, 865-871.	0.4	4
96	Elevated serum YKL40 level is a predictor of MACE during the long-term follow up in hypertensive patients. <i>Clinical and Experimental Hypertension</i> , 2020, 42, 271-274.	0.5	4
97	Flow cytometric analysis of platelet surface glycoproteins in the diagnosis of thirty-two Turkish patients with Glanzmann thrombasthenia: a multicenter experience. <i>Turkish Journal of Medical Sciences</i> , 2021, 51, 2135-2141.	0.4	4
98	Prognostic nutritional index is related to myocardial performance index in newly diagnosed nondiabetic hypertensive patients. <i>Clinical and Experimental Hypertension</i> , 2021, 43, 378-383.	0.5	4
99	Efeitos da Terapia com Anti-TNF alfa na Pressão Arterial em Pacientes com Hipertensão Resistente: Um Estudo Piloto Randomizado, Duplo-Cego e Controlado por Placebo. <i>Arquivos Brasileiros De Cardiologia</i> , 2021, 116, 443-451.	0.3	4
100	YKL-40 levels in patients with coronary artery ectasia. <i>Anatolian Journal of Cardiology</i> , 2013, 14, 97-8.	0.4	3
101	Relationship of platelet indices with acute stent thrombosis in patients with acute coronary syndrome. <i>Postępy W Kardiologii Interwencyjnej</i> , 2015, 3, 224-229.	0.1	3
102	Efficacy of striae gravidarum extension and localization on predicting intraperitoneal adhesion risk. <i>Journal of Obstetrics and Gynaecology Research</i> , 2019, 45, 2358-2363.	0.6	3
103	The Relationship Between Mean Platelet Volume and In-Hospital Mortality in Geriatric Patients with ST Segment Elevation Myocardial Infarction Who Underwent Primary Percutaneous Coronary Intervention. <i>Cardiovascular Innovations and Applications</i> , 2019, 4, .	0.1	3
104	Increased epicardial fat tissue thickness predicts advanced interatrial block among hypertensive patients. <i>Journal of Electrocardiology</i> , 2020, 61, 18-22.	0.4	3
105	The role of serum interleukin-37 levels, inflammation and blood pressure in patients with preeclampsia. <i>Clinical and Experimental Hypertension</i> , 2020, 42, 669-674.	0.5	3
106	Left atrial volume index and pulmonary arterial pressure predicted MACE among patients with STEMI during 8-year follow-up: experience from a tertiary center. <i>Herz</i> , 2020, 46, 367-374.	0.4	3
107	Secondary Hypogammaglobulinemia in Patients with Chronic Lymphocytic Leukemia Receiving Ibrutinib Therapy. <i>Indian Journal of Hematology and Blood Transfusion</i> , 2022, 38, 282-289.	0.3	3
108	Clinical staging in chronic heart failure associated with low vitamin D and elevated parathormone levels. <i>Acta Cardiologica</i> , 2014, 69, 665-71.	0.3	3

#	ARTICLE	IF	CITATIONS
109	The Coincidence of Familial Mediterranean Fever and Hypereosinophilia in a Patient with Hereditary Elliptocytosis. <i>Indian Journal of Hematology and Blood Transfusion</i> , 2014, 30, 138-141.	0.3	2
110	Acute inferior myocardial infarction in a young male patient associated with Behcet's disease and sildenafil. <i>BMJ Case Reports</i> , 2014, 2014, bcr2013201189-bcr2013201189.	0.2	2
111	Prolonged QRS duration on surface electrocardiogram is associated with left ventricular restrictive filling pattern. <i>Interventional Medicine & Applied Science</i> , 2017, 9, 9-14.	0.2	2
112	Endothelial dysfunction predicted increased left atrial volume index in newly diagnosed nondiabetic hypertensive patients. <i>Blood Pressure Monitoring</i> , 2020, 25, 75-81.	0.4	2
113	Blood urea nitrogen is associated with long-term all-cause mortality in stable angina pectoris patients: 8-year follow-up results. <i>Kardiologiya</i> , 2021, 61, 66-70.	0.3	2
114	Is Adiponectin Elevation Associated with Left Atrial Remodeling and Impaired Mechanical Functions? (a) <i>Tj ETQq0 0.0rgBT /Oyerlock 10</i>	0.3	2
115	Total Serum Protein Predicted Mortality in Patients with St-elevation Myocardial Infarction Who Underwent Primary Percutaneous Coronary Intervention: Results of 8-Year Follow-up. <i>E-Journal of Cardiovascular Medicine</i> , 2021, 9, 122-129.	0.1	2
116	A simple angiographic index to predict adverse clinical outcome associated with acute myocardial infarction. <i>Turk Kardiyoloji Dernegi Arsivi</i> , 2014, 42, 321-329.	0.6	2
117	Hair whitening and obesity are independently related to ascending aorta dilatation in young-middle aged men. <i>Åstanbul Kuzey Klinikleri</i> , 2017, 6, 33-39.	0.1	2
118	Preoperative blood urea nitrogen-to-left ventricular ejection fraction ratio is an independent predictor of long-term major adverse cardiac events in patients undergoing coronary artery bypass grafting surgery. <i>Journal of the Saudi Heart Association</i> , 2020, 32, 79-85.	0.2	2
119	Evaluation of left ventricular systolic and diastolic functions in bipolar patients during lithium therapy. <i>International Journal of Clinical and Experimental Medicine</i> , 2015, 8, 2917-22.	1.3	2
120	Primary percutaneous coronary intervention for the treatment of a patient with cardiogenic shock due to acute total occlusion of unprotected left main coronary artery without supportive collateral flow. <i>Future Cardiology</i> , 2011, 7, 749-755.	0.5	1
121	Value of IGF-I levels in the evaluation of response to treatment with levosimendan in patients with severe heart failure. <i>Anatolian Journal of Cardiology</i> , 2011, 11, 523-9.	0.4	1
122	YKL-40 as new cardiac biomarker. <i>Anatolian Journal of Cardiology</i> , 2013, 14, 98-9.	0.4	1
123	Relationship Between Mean Platelet Volume and Hypertrophic Cardiomyopathy. <i>Angiology</i> , 2014, 65, 851-851.	0.8	1
124	Detection of Acute Lymphoblastic Leukemia Involvement in Pleural Fluid in an Adult Patient with Ataxia Telangiectasia by Flow Cytometry Method. <i>Indian Journal of Hematology and Blood Transfusion</i> , 2014, 30, 73-76.	0.3	1
125	Mitral annular calcification predicted major cardiovascular events in patients presented with acute coronary syndrome and underwent percutaneous coronary intervention. <i>Acta Cardiologica</i> , 2020, 75, 767-773.	0.3	1
126	Serum osteoprotegerin level is independently related to subclinical left atrial mechanical function in patients with hypertension and diabetes. <i>Herz</i> , 2021, 46, 277-284.	0.4	1

#	ARTICLE	IF	CITATIONS
127	Assessment of cardio-ankle vascular index in patients with abdominal aortic aneurysm: An observational study. <i>Vascular</i> , 2021, 29, 190-195.	0.4	1
128	The effect of bilirubin on long-term mortality in patients with chronic total coronary occlusion. <i>Bratislava Medical Journal</i> , 2019, 120, 860-863.	0.4	1
129	Increased Epicardial Adipose Tissue is Associated with the Extent of Aortic Dissection. <i>Journal of the Saudi Heart Association</i> , 2020, 32, 415-420.	0.2	1
130	Prolonged QRS independently predicts long-term all-cause mortality in patients with narrow QRS complex undergoing coronary artery bypass grafting surgery (9-year follow-up results). <i>Kardiologia i Torakochirurgia Polska</i> , 2020, 17, 117-122.	0.1	1
131	Impact of Postdilation on Intervention Success and Long-Term Major Adverse Cardiovascular Events (MACE) among Patients with Acute Coronary Syndromes. <i>Cardiovascular Innovations and Applications</i> , 2020, 4, .	0.1	1
132	An Unusual Case of Löffler Endomyocarditis after Takotsubo Cardiomyopathy Induced by Deep Neck Infection. <i>Acta Cardiologica Sinica</i> , 2015, 31, 457-60.	0.1	1
133	The PR Interval Predicted Major Adverse Cardiovascular Events in Patients with Acute Coronary Syndrome Who Underwent Percutaneous Coronary Intervention: 3 Years Follow-up Results. <i>Journal of Academic Research in Medicine</i> , 2021, 11, 241-248.	0.1	1
134	A rare combination: Single coronary artery originating from the left sinus of valsalva coexisting with perimembranous ventricular septal defect. <i>Journal of Cardiology Cases</i> , 2012, 6, e39-e41.	0.2	0
135	Increased Platelet Indices in Acute Stent Thrombosis. <i>Angiology</i> , 2014, 65, 745-746.	0.8	0
136	Mediastinal hydatid cyst compressing the pulmonary artery and atrial septal defect: A rare association. <i>Revista Portuguesa De Cardiologia</i> , 2014, 33, 317-318.	0.2	0
137	Removal of the broken part of implantable cardioverter-defibrillator's electrode causing pulmonary embolism via femoral vein. <i>Indian Heart Journal</i> , 2015, 67, S88.e1-S88.e4.	0.2	0
138	PP-045 An Increased Epicardial Adipose Tissue Is Strongly Associated with Carotid-Intima Media Thickness and the Atherosclerotic Plaque, but LDL Only with the Plaque. <i>American Journal of Cardiology</i> , 2016, 117, S57-S58.	0.7	0
139	PP-097 Relation of Angiographic Thrombus Burden with Severity of Coronary Artery Disease in Patients with ST Segment Elevation Myocardial Infarction. <i>American Journal of Cardiology</i> , 2016, 117, S74.	0.7	0
140	Response to the Letter to the Editor: Critically ill Haematological Cancer Patients: How far the Severity Index Score can determine the outcome and duration of Aggressive Support?. <i>Australian Critical Care</i> , 2018, 31, 339.	0.6	0
141	Preoperative blood urea nitrogen-to-left ventricular ejection fraction ratio is an independent predictor of long-term major adverse cardiac events in patients undergoing coronary artery bypass grafting surgery: 10-year follow-up results. <i>Journal of the Saudi Heart Association</i> , 2019, .	0.2	0
142	Impaired left atrial mechanical functions as indicators for increased aortic root diameter in hypertensive and diabetic patients. <i>Herz</i> , 2021, 46, 272-279.	0.4	0
143	An echocardiographic predictor of contrast-induced nephropathy following percutaneous coronary intervention in acute coronary syndrome: aortic valve sclerosis. <i>Scandinavian Cardiovascular Journal</i> , 2020, 54, 227-231.	0.4	0
144	Paradoxical association between lipoprotein cholesterol levels and left atrial function in hypertensive diabetic patients: A speckle tracking study. <i>Journal of Clinical Ultrasound</i> , 2021, 49, 667-673.	0.4	0

#	ARTICLE	IF	CITATIONS
145	Predictors for the prolonged R wave peak time among patients with arterial hypertension. Turk Kardioloji Dernegi Arsivi, 2021, 49, 387-394.	0.6	0
146	FEV1 is independently related with impaired left atrial strain in chronic obstructive pulmonary disease patients: A speckle tracking study. Clinical Respiratory Journal, 2021, 15, 1359-1367.	0.6	0
147	Giant superior mesenteric artery aneurysm. Dicle Medical Journal, 2011, 38, 352-354.	0.2	0
148	A rare sign of ischemia during exercise ECG: PR interval lengthening in the recovery period. Turk Kardioloji Dernegi Arsivi, 2012, 40, 719-722.	0.6	0
149	Association between deletion polymorphism of angiotensin converting enzyme gene and pulmonary hypertension in pulmonary thromboembolism. Turkish Journal of Thoracic and Cardiovascular Surgery, 2012, , 281-286.	0.2	0
150	Myopericarditis Caused by H1N1 Virus Infection. Electronic Journal of General Medicine, 2013, 10, .	0.3	0
151	Carotid Artery Stenting in Symptomatic High Risk Patients: A Single Center Experience. KoÅŸuyolu Heart Journal, 2014, 17, 91-94.	0.0	0
152	Gastrointestinal Plasmoblastic Lymphoma Associated with Wiskott-Aldrich Syndrome: A Case Report. Blood, 2019, 134, 5359-5359.	0.6	0
153	A New Scoring System Detecting the Supraventricular Tachycardia Inducibility and Radiofrequency Ablation With High Specificity in Electrophysiological Study. Journal of the Saudi Heart Association, 2020, 32, 134-140.	0.2	0
154	Erken SaÅŸ BeyazlamasÄ± SaÅŸlÄ±klÄ± GenÅŸ ve Orta YaÅŸlÄ± Erkekler iÅŸin Hipertansiyonun Belirleyicisi midir?. Sakarya Medical Journal, 0, , .	0.1	0
155	Left atrial volume index to left ventricular ejection fraction ratio predicted major adverse cardiovascular event in ST-Elevated myocardial infarction patients during 8 years of follow-up. Journal of Cardiovascular Echography, 2021, 31, 227.	0.1	0
156	Fragmented QRS predicted major adverse cardiovascular events in patients with coronary artery disease and percutaneous coronary intervention, 10-years of follow-up. Kardiologiya, 2022, 62, 72-79.	0.3	0
157	Left ventricular noncompaction associated with a sinoatrial node artery originating from the posterolateral branch of the right coronary artery. Acta Cardiologica, 2011, 66, 83-5.	0.3	0