

Daniel Pineda-Tenor

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/642770/publications.pdf>

Version: 2024-02-01

25
papers

337
citations

933447

10
h-index

839539

18
g-index

26
all docs

26
docs citations

26
times ranked

693
citing authors

#	ARTICLE	IF	CITATIONS
1	The IL7RA rs6897932 polymorphism is associated with progression of liver fibrosis in patients with chronic hepatitis C: Repeated measurements design. PLoS ONE, 2018, 13, e0197115.	2.5	10
2	Mx1, OAS1 and OAS2 polymorphisms are associated with the severity of liver disease in HIV/HCV-coinfected patients: A cross-sectional study. Scientific Reports, 2017, 7, 41516.	3.3	22
3	<i>CXCL9</i> polymorphisms are associated with liver fibrosis in patients with chronic hepatitis C: a cross-sectional study. Clinical and Translational Medicine, 2017, 6, 26.	4.0	13
4	Soluble Adhesion Molecules in Patients Coinfected with HIV and HCV: A Predictor of Outcome. PLoS ONE, 2016, 11, e0148537.	2.5	8
5	<i>IL15</i> polymorphism is associated with advanced fibrosis, inflammation-related biomarkers and virological response in human immunodeficiency virus/hepatitis C virus coinfection. Liver International, 2016, 36, 1258-1266.	3.9	5
6	Short Communication: <i>CXCL12</i> rs1029153 Polymorphism Is Associated with the Sustained Virological Response in HIV/Hepatitis C Virus-Coinfected Patients on Hepatitis C Virus Therapy. AIDS Research and Human Retroviruses, 2016, 32, 226-231.	1.1	0
7	Relationship between ITPA polymorphisms and hemolytic anemia in HCV-infected patients after ribavirin-based therapy: a meta-analysis. Journal of Translational Medicine, 2015, 13, 320.	4.4	19
8	Association between IL7R polymorphisms and severe liver disease in HIV/HCV coinfecting patients: a cross-sectional study. Journal of Translational Medicine, 2015, 13, 206.	4.4	10
9	Reply. Hepatology, 2015, 62, 1643-1643.	7.3	2
10	<i>IL7</i> polymorphisms predict the <i>CD4</i> recovery in <i>HIV</i> patients on <i>ART</i>. European Journal of Clinical Investigation, 2015, 45, 1192-1199.	3.4	12
11	Single Nucleotide Polymorphisms of CXCL9-11 Chemokines Are Associated With Liver Fibrosis in HIV/HCV-Coinfected Patients. Journal of Acquired Immune Deficiency Syndromes (1999), 2015, 68, 386-395.	2.1	11
12	TLR3 polymorphisms are associated with virologic response to hepatitis C virus (HCV) treatment in HIV/HCV coinfecting patients. Journal of Clinical Virology, 2015, 65, 62-67.	3.1	6
13	Toll-like receptor 8 (TLR8) polymorphisms are associated with non-progression of chronic hepatitis C in HIV/HCV coinfecting patients. Infection, Genetics and Evolution, 2015, 36, 339-344.	2.3	6
14	rs7903146 Polymorphism at <i>Transcription Factor 7 Like 2</i> Gene Is Associated with Total Cholesterol and Lipoprotein Profile in HIV/Hepatitis C Virus-Coinfected Patients. AIDS Research and Human Retroviruses, 2015, 31, 326-334.	1.1	5
15	Relationship between European Mitochondrial Haplogroups and Chronic Renal Allograft Rejection in Patients with Kidney Transplant. International Journal of Medical Sciences, 2014, 11, 1129-1132.	2.5	3
16	FTOrs9939609 polymorphism is associated with metabolic disturbances and response to HCV therapy in HIV/HCV-coinfected patients. BMC Medicine, 2014, 12, 198.	5.5	4
17	Association of adiponectin (<i>ADIPOQ</i>) rs2241766 polymorphism and dyslipidemia in <i>HIV</i>/<i>HCV</i>-coinfecting patients. European Journal of Clinical Investigation, 2014, 44, 453-462.	3.4	12
18	SLC30A8 rs13266634 polymorphism is related to a favorable cardiometabolic lipid profile in HIV/hepatitis C virus-coinfected patients. Aids, 2014, 28, 1325-1332.	2.2	9

#	ARTICLE	IF	CITATIONS
19	Vitamin D deficiency is associated with severity of liver disease in HIV/HCV coinfecting patients. <i>Journal of Infection</i> , 2014, 68, 176-184.	3.3	28
20	CXCL9, CXCL10 and CXCL11 polymorphisms are associated with sustained virologic response in HIV/HCV-coinfecting patients. <i>Journal of Clinical Virology</i> , 2014, 61, 423-429.	3.1	13
21	Relationship of vitamin D status with advanced liver fibrosis and response to hepatitis C virus therapy: A meta-analysis. <i>Hepatology</i> , 2014, 60, 1541-1550.	7.3	68
22	PPAR β Pro12Ala polymorphism was associated with favorable cardiometabolic risk profile in HIV/HCV coinfecting patients: a cross-sectional study. <i>Journal of Translational Medicine</i> , 2014, 12, 235.	4.4	11
23	A proposed reference change value for an IgA anti-tissue transglutaminase immunoassay to improve interpretation of serial results in celiac patients. <i>Clinica Chimica Acta</i> , 2013, 421, 12-16.	1.1	1
24	Biological variation and reference change values of common clinical chemistry and haematologic laboratory analytes in the elderly population. <i>Clinical Chemistry and Laboratory Medicine</i> , 2013, 51, 851-862.	2.3	45
25	Extreme concentrations of high density lipoprotein cholesterol affect the calculation of low density lipoprotein cholesterol in the Friedewald formula and other proposed formulas. <i>Clinical Biochemistry</i> , 2011, 44, 1451-1456.	1.9	13