

# Daniel Pineda-Tenor

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/642770/publications.pdf>

Version: 2024-02-01

25  
papers

337  
citations

932766

10  
h-index

839053

18  
g-index

26  
all docs

26  
docs citations

26  
times ranked

693  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | The IL7RA rs6897932 polymorphism is associated with progression of liver fibrosis in patients with chronic hepatitis C: Repeated measurements design. PLoS ONE, 2018, 13, e0197115.   | 1.1 | 10        |
| 2  | Mx1, OAS1 and OAS2 polymorphisms are associated with the severity of liver disease in HIV/HCV-coinfected patients: A cross-sectional study. Scientific Reports, 2017, 7, 41516.   | 1.6 | 22        |
| 3  | <i>CXCL9</i> polymorphisms are associated with liver fibrosis in patients with chronic hepatitis C: a cross-sectional study. Clinical and Translational Medicine, 2017, 6, 26.  | 1.7 | 13        |
| 4  | Soluble Adhesion Molecules in Patients Coinfected with HIV and HCV: A Predictor of Outcome. PLoS ONE, 2016, 11, e0148537.   | 1.1 | 8         |
| 5  | <i>IL15</i> polymorphism is associated with advanced fibrosis, inflammation-related biomarkers and virological response in human immunodeficiency virus/hepatitis C virus coinfection. Liver International, 2016, 36, 1258-1266.                    | 1.9 | 5         |
| 6  | Short Communication: <i>CXCL12</i> rs1029153 Polymorphism Is Associated with the Sustained Virological Response in HIV/Hepatitis C Virus-Coinfected Patients on Hepatitis C Virus Therapy. AIDS Research and Human Retroviruses, 2016, 32, 226-231. | 0.5 | 0         |
| 7  | Relationship between ITPA polymorphisms and hemolytic anemia in HCV-infected patients after ribavirin-based therapy: a meta-analysis. Journal of Translational Medicine, 2015, 13, 320.   | 1.8 | 19        |
| 8  | Association between IL7R polymorphisms and severe liver disease in HIV/HCV coinfecting patients: a cross-sectional study. Journal of Translational Medicine, 2015, 13, 206.   | 1.8 | 10        |
| 9  | Reply. Hepatology, 2015, 62, 1643-1643.   | 3.6 | 2         |
| 10 | <i>IL7</i> polymorphisms predict the <i>CD4</i> recovery in HIV patients on <i>ART</i>. European Journal of Clinical Investigation, 2015, 45, 1192-1199.  | 1.7 | 12        |
| 11 | Single Nucleotide Polymorphisms of CXCL9-11 Chemokines Are Associated With Liver Fibrosis in HIV/HCV-Coinfected Patients. Journal of Acquired Immune Deficiency Syndromes (1999), 2015, 68, 386-395.  | 0.9 | 11        |
| 12 | TLR3 polymorphisms are associated with virologic response to hepatitis C virus (HCV) treatment in HIV/HCV coinfecting patients. Journal of Clinical Virology, 2015, 65, 62-67.  | 1.6 | 6         |
| 13 | Toll-like receptor 8 (TLR8) polymorphisms are associated with non-progression of chronic hepatitis C in HIV/HCV coinfecting patients. Infection, Genetics and Evolution, 2015, 36, 339-344.   | 1.0 | 6         |
| 14 | rs7903146 Polymorphism at <i>Transcription Factor 7 Like 2</i> Gene Is Associated with Total Cholesterol and Lipoprotein Profile in HIV/Hepatitis C Virus-Coinfected Patients. AIDS Research and Human Retroviruses, 2015, 31, 326-334.             | 0.5 | 5         |
| 15 | Relationship between European Mitochondrial Haplogroups and Chronic Renal Allograft Rejection in Patients with Kidney Transplant. International Journal of Medical Sciences, 2014, 11, 1129-1132.   | 1.1 | 3         |
| 16 | FTOrs9939609 polymorphism is associated with metabolic disturbances and response to HCV therapy in HIV/HCV-coinfected patients. BMC Medicine, 2014, 12, 198.  | 2.3 | 4         |
| 17 | Association of adiponectin (<i>ADIPOQ</i>) rs2241766 polymorphism and dyslipidemia in HIV/HCV-coinfected patients. European Journal of Clinical Investigation, 2014, 44, 453-462.   | 1.7 | 12        |
| 18 | SLC30A8 rs13266634 polymorphism is related to a favorable cardiometabolic lipid profile in HIV/hepatitis C virus-coinfected patients. Aids, 2014, 28, 1325-1332.  | 1.0 | 9         |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | Vitamin D deficiency is associated with severity of liver disease in HIV/HCV coinfecting patients. <i>Journal of Infection</i> , 2014, 68, 176-184.   | 1.7 | 28        |
| 20 | CXCL9, CXCL10 and CXCL11 polymorphisms are associated with sustained virologic response in HIV/HCV-coinfecting patients. <i>Journal of Clinical Virology</i> , 2014, 61, 423-429.   | 1.6 | 13        |
| 21 | Relationship of vitamin D status with advanced liver fibrosis and response to hepatitis C virus therapy: A meta-analysis. <i>Hepatology</i> , 2014, 60, 1541-1550.  | 3.6 | 68        |
| 22 | PPAR $\beta$ Pro12Ala polymorphism was associated with favorable cardiometabolic risk profile in HIV/HCV coinfecting patients: a cross-sectional study. <i>Journal of Translational Medicine</i> , 2014, 12, 235.                       | 1.8 | 11        |
| 23 | A proposed reference change value for an IgA anti-tissue transglutaminase immunoassay to improve interpretation of serial results in celiac patients. <i>Clinica Chimica Acta</i> , 2013, 421, 12-16.                                   | 0.5 | 1         |
| 24 | Biological variation and reference change values of common clinical chemistry and haematologic laboratory analytes in the elderly population. <i>Clinical Chemistry and Laboratory Medicine</i> , 2013, 51, 851-862.                    | 1.4 | 45        |
| 25 | Extreme concentrations of high density lipoprotein cholesterol affect the calculation of low density lipoprotein cholesterol in the Friedewald formula and other proposed formulas. <i>Clinical Biochemistry</i> , 2011, 44, 1451-1456. | 0.8 | 13        |